arXiv:hep-ph/0308049v1 5 Aug 2003

Group Theoretical Analysis of Light-Cone Dominated Hadronic Processes and Twist Decomposition of Nonlocal Operators in Quantum Chromodynamics

Markus Lazar

Fakultät für Physik und Geowissenschaften
der Universität Leipzig

Doctoral Thesis

June 17, 2002

Abstract

In the framework of nonlocal light-cone expansion of two current operators we construct bilocal as well as trilocal QCD light-cone operators with definite geometric twist. We are able to decompose uniquely the appearing QCD light-cone operators into all their twist parts on the light-cone.

The quark distribution functions and the vector meson distribution amplitudes which are defined as forward and skewed nucleonic matrix elements of nonlocal light-cone operators are classified with respect to geometric twist. Hereby we found well-kown and new Wandzura-Wilczek type relations between the quark distribution functions with dynamical twist and, additionally, between the vector meson distribution amplitudes with dynamical twist.

Using group theoretical methods we are able to construct all bilocal off-cone QCD operators with twist 2 and 3. Additionally, we calculate their nucleonic matrix elements (double distribution functions, scalar and vector meson distribution amplitudes). In some special cases we give also the twist decomposition of off-cone QCD operators from minimal up to infinite twist.

Contents

1 Introduction 1

2 Quantum field theoretical treatment of light-cone dominated hadronic processes 4
- 2.1 Light-cone dominated scattering processes and light-cone expansion in QCD . . . 4
  - 2.1.1 Phenomenology of hard scattering processes in QCD 4
  - 2.1.2 Light-cone expansion 7
  - 2.1.3 Distribution amplitudes 12
  - 2.1.4 Anomalous dimensions of QCD operators 15
- 2.2 The concept of geometric twist of nonlocal operators 18
  - 2.2.1 The procedure of twist decomposition of nonlocal operators 18
  - 2.2.2 Transformation properties of (non)local operators with definite geometric twist 22

3 Application of the twist decomposition to different aspects of the hadron phenomenology 24
- 3.1 Quark distribution functions of definite geometric twist 24
  - 3.1.1 Quark distribution functions of definite dynamical twist 24
  - 3.1.2 Twist decomposition of bilinear quark operators 26
  - 3.1.3 Forward matrix elements of LC–operators with definite geometric twist . 28
  - 3.1.4 Relations between quark distribution functions of dynamical and geometric twist 33
  - 3.1.5 Wandzura-Wilczek type relations 36
  - 3.1.6 Conclusions 40
- 3.2 Vector meson distribution amplitudes of definite geometric twist 42
  - 3.2.1 The ρ-meson distribution amplitudes of definite dynamical twist 42
  - 3.2.2 The ρ-meson matrix elements of LC–operators with definite geometric twist 44
  - 3.2.3 The relations between ρ-meson distribution amplitudes of dynamical and geometric twist 48
  - 3.2.4 Wandzura-Wilczek type relations 50
  - 3.2.5 Conclusions 55
- 3.3 Power corrections of non-forward quark distributions and harmonic operators with definite geometric
  - 3.3.1 Non-forward matrix elements of harmonic operators: The method 56
  - 3.3.2 Power Corrections of non-forward matrix elements 62
  - 3.3.3 Conclusions 70

4 The twist decomposition of nonlocal tensor operators 71
- 4.1 Irreducible tensor polynomials and harmonic extension in $2h$-dimensional spacetime – polynomial tec
- 4.2 Twist decomposition of a bilocal 2nd rank tensor operator 80
  - 4.2.1 Tensor operators of symmetry class (i) 82
  - 4.2.2 Tensor operators of symmetry class (ii) 86
  - 4.2.3 Tensor operators of symmetry class (iii) 94
  - 4.2.4 Tensor operators of symmetry class (iv) 96

4.2.5 Conclusions 101
4.3 Extension to trilocal tensor operators 102
4.3.1 General trilocal tensor operators: Quark–antiquark–gluon, four–quark and four–gluon operators103
4.3.2 Constrained trilocal operators: Shuryak-Vainshtein– and three–gluon operators106
4.3.3 Conclusions 111
4.4 Harmonic operators of definite geometric twist in terms of Gegenbauer polynomials and Bessel functions112
4.4.1 Totally symmetric harmonic operators of any geometric twist 112
4.4.2 Harmonic quark-antiquark operators of geometric twist 2 and 3 118

5 Conclusions and Outlook 124

A Symmetry classes and tensor representations 126

B Irreducible tensor representations of the Lorentz group 129

C The interior derivative on the light-cone and light-cone operators with definite geometric twist132

D Target mass corrections for a scalar theory in $2h$-dimensional spacetime 135

Chapter 1

Introduction

Quantum Chromodynamics (QCD) which is a relativistic local quantum field theory based on the non-Abelian colour gauge group $SU_{\mathrm{c}}(3)$ is well established nowadays as a microscopic theory of strong interaction. Because the gauge group is non-Abelian a gauge theory of this kind is called Yang-Mills theory. In a gauge theory exist fields which transform according to the adjoint representation (gauge fields) of the gauge group and spinor fields which transform as a fundamental representation (matter fields). The matter fields in QCD are "coloured" quarks. The interaction of quarks is mediated by the gauge bosons (gluons). Quarks and gluons form composite particles called hadrons (mesons and baryons). Due to presently unsolved bound state problem it is not possible to predict measurable observables from the quark and gluon fields. Although the high energy behaviour of QCD is believed to be described by the perturbation theory due to the asymptotically free nature of QCD, the cross sections are complicated combinations of short- and long-distance interactions and is not directly calculable. The standard approach to overcome these difficulties is based on the factorization theorem which allows to separate (factorize) the soft and hard part of the cross section in a systematic fashion. The soft part (long-distance part) can be described by expection values of QCD operators sandwiched between hadronic states. Due to the confinement this part can be determined only from experimental data or a suitable lattice calculation. On the other hand, the hard subprocess (short-distance part) and the evolution of the functions which are occurring in the soft part can be systematically calculated with the help of perturbation theory. The success of perturbative QCD in predicting the momentum dependence of hadronic observables serves an important argument for the correctness of the theory. So far all experimental data are consistent with the predictions of QCD.

The universal, non-perturbative parton distribution amplitudes parametrizing, modulo kinematical factors, the matrix elements of appropriate non-local quark-antiquark (as well as gluon) operators play a central role in phenomenological considerations. These operators occur in the quantum field theoretic description of light-cone dominated hadronic processes via the non-local light-cone expansion [6, 126, 106]. For deep inelastic lepton-hadron scattering and Drell-Yan processes the parton distributions are given as forward matrix elements of bilocal light-ray operators resulting from the time-ordered product of appropriate hadronic currents. For (deeply) virtual Compton scattering and hadron wave functions the so-called double distributions and hadron distribution amplitudes, respectively, are given by corresponding non-forward matrix elements. Thereby, various processes are governed by one and the same (set of) light-ray operators. The computation of the anomalous dimensions of these operators is one of the most important problems of the theory, since these quantities determine the deviation from the scaling behaviour of experimental observables.

The success of QCD as fundamental theory of strong interactions is intimately tied to its ability to describe hard exclusive and inclusive reactions, which has been tested in numerous experiments. In the corresponding kinematic regime, i.e., at large space-like virtualities, the relevant amplitudes are dominated by singularities on the light-cone, which, in the framework of a light-cone expansion, can be described in terms of contributions of definite twist. The notion of (geometric) twist, for local operators, has been introduced originally by Gross and Treiman [69] as twist = (canonical) dimension − (Lorentz) spin; it uses the irreducible representations of the orthochronous Lorentz group and, as such, is a Lorentz-invariant concept. Using the notion of twist, one is able to expand systematically the cross section in inverse powers of a characteristic large momentum scale Q^2, and extract the leading contributions. The higher twist effects describe the quark-gluon behaviour inside the nucleon, and thus contain information about the quark-gluon correlations. Possibly, in the near future the experimental precision data will allow for the determination of non-leading contributions and, therefore, require for a careful analysis of the various sub-dominant effects contributing to the physical processes. When considered beyond leading order, i.e., beyond lowest twist operators in tree approximation, one is confronted not only with radiative corrections but also with power corrections resulting from higher twist as well as target mass effects. Higher twist contributions are obtained by the decomposition of the operators with respect to (irreducible tensor) representations of the Lorentz group having definite twist τ. These representations are characterized by their symmetry type (under index permutations of the traceless tensors) which is determined by corresponding Young tableaux. With the growing accuracy of the experimental data a perturbative expansion with respect to the experimentally relevant variable, M^2/Q^2, M being a typical mass of the process, will be very useful.

An alternative approach to twist-counting is based on the light-cone quantization in the infinite momentum frame [93] with the decomposition of the quark fields into "good" and "bad" components, $\psi = \psi_+ + \psi_-$, with $\psi_\pm = \frac{1}{2}\gamma^\mp\gamma^\pm\psi$. As has been pointed out in [78, 79], a "bad" component introduces one unit of M/Q because these components are not dynamically independent and may be expressed through the equations of motion by the "good" ones times the above kinematical factor. Despite being conceptionally different this definition looks similar to the phenomenologically more convenient one which counts only powers of $1/Q$ in the infinite momentum frame [78, 79]. However, these power–counting concepts of "dynamical twist" are not Lorentz invariant and do not agree with the original definition. Furthermore, the same power of $1/Q$ may occur for different values of τ. The mismatch between dynamical and geometric twist becomes relevant once power-suppressed higher-twist contributions are included and leads to so-called Wandzura-Wilczek (WW) relations between matrix-elements of operators of different dynamical, but identical geometric twist, the prototype of which has been derived by Wandzura and Wilczek for the nucleon distribution functions (DFs) g_1 and g_2 [123].

In addition, the "dynamical twist" is *applicable only for matrix elements* of the operators and, therefore, is not directly related to the operators itself. Even more, if the matrix elements contain more than one momentum, as will be the case for the deeply virtual Compton scattering, the definition of Q is quite ambiguous and different definitions are related according to $Q \to Q(1 + f(p_\pm, q)M^2/Q^2 + \ldots)$ with some Lorentz invariant factor f. Therefore, from the point of view of renormalized quantum field theory only the original geometric definition of twist is a well-defined concept. The problem which remains is how this concept has to be generalized to the nonlocal operators appearing in the nonlocal light–cone expansion.

Because the (geometrical) twist is a straightforward scheme to decompose local operators and to classify generalized parton as well as skewed distribution amplitudes we generalize this notation for nonlocal operators, in principle, for any twist. The considered operators are necessary for the investigation of different phenomenological QCD processes. We perform this program for light-cone as well as off-cone QCD operators.

This thesis is organized as follows. The Section 2.1 is mainly introductory. It collects the necessary definitions, physical motivations and explains the basic ideas. In Section 2.2 we give a motivation why we use the concept of geometric twist for nonlocal operators. We collect and explain the steps for the procedure of twist decomposition of nonlocal operators. In Section 3.1 and 3.2 we give the straightforward classification with respect to (geometric) twist of quark distribution amplitudes and vector meson distribution amplitudes on the light-cone. These both classifications are quite similar because the same light-cone operators are related to different hadronic processes. A further problem, which we will solve, is how the (geometrical) twist decomposition of parton distributions which has been given in this work relates to the (dynamical) twist decomposition of parton distributions given, e.g., by Jaffe and Ji [78, 79] and to the decomposition of vector meson distribution amplitudes given by Ball and Braun [14, 15]. The interrelations between the dynamical and geometrical distribution amplitudes are calculated and used to derive Wandzura-Wilczek type relations between the dynamical twist distribution functions. Most of them are new and might be of phenomenological interest. The material presented in these two Sections was first published in Ref. [66] and in Refs. [97, 98]. In Section 3.3 we calculated the power corrections of double distribution amplitudes, and meson as well as pion distribution amplitudes with definite geometric twist. With the help of the harmonic extension we are able to resum the target mass or power corrections in general non-forward matrix elements. The material presented in this Section is based on Ref. [65].

The Section 4.1 contains all formulas of the so-called tensor polynomials which are relevant for the twist or spin decomposition in $2h$-dimensional spacetime in this work. The constructed tensor polynomials are irreducible with respect to $SO(1, 2h-1)$ and $SO(2h)$, respectively, on-cone as well as off-cone. They are given on-cone as well as off-cone.

In Section 4.2 we give systematically the complete twist decomposition of the various gluonic bilocal light–ray tensor operators of second rank being specified by different symmetry classes; their twist content ranges from 2 up to 6.

In Section 4.3 we extend these results with minor modifications to the trilocal light–ray operators: trilocal quark–gluon correlation operators (related to so-called Shuryak-Vainshtein operators) and four–fermion operator, as well as multilocal quark and gluon operators.

In Section 4.4 we present the general twist decomposition of off-cone operators. The procedure is very close to the so-called Nachtmann method for the mass corrections in inclusive processes. This approach is based on (irreducible) harmonics of $SO(1,3)$ and $SO(4)$, respectively. From the group theoretical point of view it also works for the case of non-forward processes. The power of that approach consists in the fact that one determines the twist decomposition at first for the local and nonlocal operators and only afterwards takes the matrix elements. For totally symmetric (local) operators we are able to study the decomposition up to infinite twist. Additionally we give the twist decomposition of operators with other symmetry types up to twist 2 and 3.

The technical details which are needed for the twist decomposition are collected in the Appendixes. In Appendix A a short exposition about the characterization of irreducible tensor representations through the Young tableaux is given and the Appendix B contains some basic notation with respect to the Lorentz group. Appendix C is devoted to the concept of an interior differential operator on the light-cone which was originally introduced by Bargmann and Todorov [19]. In Appendix D we investigate the target mass corrections for a scalar theory in $2h$-dimensional spacetime. In this framework we calculate the corresponding scattering amplitude.

Chapter 2

Quantum field theoretical treatment of light-cone dominated hadronic processes

2.1 Light-cone dominated scattering processes and light-cone expansion in QCD

In this Section we discuss the quantum field theoretical description of light-cone dominated scattering processes. In fact, we consider some scattering amplitudes, QCD-operators and corresponding nucleonic matrix elements. Unfortunately, so far it has not been possible to make an ab initio calculation of a QCD scattering amplitude on hadrons. The main reason is due to the fact that hadrons are bound states of both quarks and gluons (confinement). Anyway, until the confinement is not better understood, a way out to calculate QCD processes is the light-cone expansion together with the renormalization group equation.

2.1.1 Phenomenology of hard scattering processes in QCD

The scattering amplitudes of light-cone dominated hadronic processes, e.g. DIS, according to the factorization hypothesis – are usually represented by the convolution of the hard (process-dependent) scattering amplitude of the partons with appropriate soft (process-independent) parton distribution amplitudes. For many hard scattering processes in QCD which are dominated on the light-cone, the nonlocal light–cone (LC) expansion, together with the renormalization group equation, is a powerful tool to determine the dependence of the nonperturbative distribution amplitudes on the experimentally relevant momentum transfer Q^2. This is the case for, e.g., the parton distributions in deep inelastic scattering (DIS), the pion as well as vector meson distribution amplitudes and, as growing up more recently, the non–forward distribution functions in deeply virtual Compton scattering (DVCS).

First, we discuss the relation between the forward virtual Compton scattering and the DIS. The virtual Compton amplitude for the forward case is defined by

$$T_{\mu\nu}(P,Q) = \mathrm{i} \int \mathrm{d}^4x \, \mathrm{e}^{\mathrm{i}qx} \, \langle P,S \, | T(J_\mu(x/2)J_\nu(-x/2)) | \, P,S\rangle, \tag{2.1.1}$$

as shown graphically in Fig. 2.1. Here, $|P,S\rangle$ is the hadron state with momentum P and spin S. The imaginary part of the virtual Compton scattering amplitude, $T^{\mu\nu}(P,Q^2)$, is related to



Figure 2.1: Kinematic notation for the forward Compton amplitude.

the hadronic tensor $W^{\mu\nu}$ with the help of the optical theorem according to

$$W^{\mu\nu} = \frac{1}{2\pi}\,\mathrm{Im}\,T^{\mu\nu}. \tag{2.1.2}$$

The hadronic tensor is given by

$$W^{\mu\nu} = \frac{1}{4\pi}\int \mathrm{d}^4x\,\mathrm{e}^{\mathrm{i}qx}\langle P,S|\,[J^\mu(x/2),J^\nu(-x/2)]\,|P,S\rangle. \tag{2.1.3}$$

It describes the internal structure of the nucleon in deep inelastic scattering (DIS). DIS is a scattering process between leptons and a fixed hadronic target. The incoming lepton (l) with momentum k interacts with the hadron (N) target with momentum P through the exchange of a virtual photon of momentum $q = k - k'$ if we consider only the QED interaction between the lepton and the nucleon and keep only the lowest order (see Fig. 2.2). Thus, it is possible to calculate the hadronic tensor by means of the virtual Compton amplitude in the framework of OPE. Let S_α be the spin vector of the nucleon so that



Figure 2.2: The DIS process in Born approximation.

$$P^2 = M^2, \qquad S^2 = -M^2, \qquad (PS) = 0, \tag{2.1.4}$$

where M denotes the mass of the nucleon target.

On the other hand, using Lorentz covariance, gauge invariance, parity conservation and discrete symmetries of the strong interaction the hadronic tensor for the case of a nucleon target can be expressed in the Bjorken limit:

$$\nu = qP \longrightarrow \infty, \qquad Q^2 = -q^2 \longrightarrow \infty \tag{2.1.5}$$

with the scaling variable

$$\xi = \frac{Q^2}{2\nu} \qquad \text{fixed} \tag{2.1.6}$$

by the four scalar structure functions F_1, F_2, g_1 and g_2 according to

$$W^{\mu\nu} = \left(\frac{q^\mu q^\nu}{q^2} - g^{\mu\nu}\right) F_1(\xi, Q^2) + \left(P^\mu - \frac{(Pq)}{q^2} q^\mu\right) \left(P^\nu - \frac{(Pq)}{q^2} q^\nu\right) F_2(\xi, Q^2)$$
$$- \mathrm{i}\epsilon^{\mu\nu\alpha\beta} q_\alpha \left(\frac{S_\beta}{(Pq)} G_1(\xi, Q^2) + \frac{S_\beta (Pq) - P_\beta (Sq)}{(Pq)^2} G_2(\xi, Q^2)\right) . \tag{2.1.7}$$

The symmetric (antisymmetric) part of $W^{\mu\nu}$ is related to the unpolarized (polarized) scattering. The structure functions F_1 and F_2 can be measured by using an unpolarized beam and target, the structure functions g_2 and g_2 require both a polarized beam and a polarized target. The differential cross-section for inclusive scattering $l(k) + N(P) \to l(k') + X(P_X)$ (see Fig. 2.2) can be separated into the leptonic ($L^{\mu\nu}$) and the hadronic tensor. The leptonic tensor can be calculated completely in the framework of QED. Unlike the leptonic tensor, the hadronic tensor cannot be computed directly from QCD because of non-perturbative effects in the strong interaction dynamics.

In the Bjorken limit the structure functions are functions of ξ and independent of Q^2 to leading order, a property known as scaling. Radiative corrections in QCD produce a small logarithmic Q^2 dependence of the structure functions which is calculable for large Q^2 since QCD is asymptotically free. Accordingly, the scale invariance (dilation symmetry) in QCD is broken due to quantum corrections and introduces scaling violations (anomalous dimensions). The Q^2 evaluation of the structure functions or parton distribution functions is calculable in perturbative QCD. It is governed by the anomalous dimensions of the corresponding operators and the renormalization group equation.

Second, we discuss the off-forward virtual Compton scattering. The Compton amplitude for the scattering of a virtual photon off a hadron provides another possibility in QCD to understand the short-distance behaviour of the theory. The Compton amplitude for this general case of non-forward scattering is given by

$$T_{\mu\nu}(P_+, P_-, Q) = \mathrm{i} \int \mathrm{d}^4x \, \mathrm{e}^{\mathrm{i}qx} \, \langle P_2, S_2 \, | T(J_\mu(x/2) J_\nu(-x/2)) | \, P_1, S_1 \rangle. \tag{2.1.8}$$

Here,

$$P_+ = P_2 + P_1, \qquad P_- = P_2 - P_1 = q_1 - q_2, \tag{2.1.9}$$
$$q = (q_1 + q_2)/2, \qquad P_1 + q_1 = P_2 + q_2, \tag{2.1.10}$$

where $q_1(q_2)$ and $P_1(P_2)$ denote the four-momenta of the incoming(outgoing) photon and hadron, respectively, and S_1, S_2 are the spins of the initial- and final-state hadron. The kinematics of this process is depicted schematically in Fig. 2.3. To specify the asymptotics of virtual Compton



Figure 2.3: Kinematic notation for the off-forward Compton amplitude.

scattering the generalized Bjorken region will be defined as follows

$$\nu = \frac{qP_+}{2} \longrightarrow \infty, \qquad Q^2 = -q^2 \longrightarrow \infty \tag{2.1.11}$$

with the two independent scaling variables

$$\xi = \frac{Q^2}{qP_+}, \qquad \eta = \frac{qP_-}{qP_+} = \frac{q_1^2 - q_2^2}{2\nu}, \tag{2.1.12}$$

to be held fixed by the experimental setup. These variables which are not restricted to the interval $[-1, +1]$ are obvious generalizations of the usual Bjorken variable and the "skewedness" parameter. Of special experimental importance are the cases of deep inelastic scattering (DIS) described by the absorptive part of the forward Compton amplitude (see above), $\eta = 0$, and the deeply virtual Compton scattering (DVCS)[83] with one real outgoing photon $q_2^2 = 0$ corresponding to $\xi = -\eta$. In the generalized Bjorken region the amplitude (2.1.8) is dominated by the light-cone singularities which allows to apply the OPE. For DIS as well as DVCS the relevant information about the non-perturbative structure of the nucleon is contained in the expectation values of QCD operators in the nucleon, which can be interpreted as the (generalized) parton distributions and "skewed" parton distributions, respectively.

Similar considerations also hold for hard (light-cone dominated) exclusive processes in QCD. The amplitudes of those processes are expressed by the factorization formulas which separate the short-distance part from the long-distance one. Vector mesons ($V = \rho$, ω, K^*, ϕ) are produced in processes like exclusive semileptonic (or radiative) B decays ($B \to ve\nu$, $B \to V + \gamma$) and hard electroproduction of vector mesons ($\gamma^* + N \to V + N'$). The physical interest in this case arises from the fact that the following vertex function is directly observable in QCD:

$$T^V_{\mu\nu}(q, P; \lambda) = \mathrm{i} \int \mathrm{d}^4x \, \mathrm{e}^{\mathrm{i}qx} \langle 0| RT \left(J_\mu(x/2) J_\nu(-x/2) \mathcal{S}\right) |V(P, \lambda)\rangle, \tag{2.1.13}$$

where J_μ can be axial or vector currents and $|V(P, \lambda)\rangle$ is the vector meson state of momentum P and helicity λ. Here, the universal nonperturbative quantities, the meson distribution amplitudes (DAs), sometimes also called meson wave functions, are given by means of the vacuum-to-meson matrix elements of bi-local operators on the light-cone, $\langle 0|O_\Gamma(\tilde{x}, -\tilde{x})|V(P, \lambda)\rangle$, with $\Gamma = \{1, \gamma_5; \gamma_\alpha, \gamma_\alpha\gamma_5; \sigma_{\alpha\beta}, \sigma_{\alpha\beta}\gamma_5\}$ and describe the long-distance part.

2.1.2 Light-cone expansion

The light-cone expansion (LCE) is a straightforward scheme to treat different light-cone dominated scattering processes. It is possible to obtain asymptotically valid informations about these processes in this framework. There exist different versions of the LCE, e.g., the local and the nonlocal LCE.

The local LCE is a generalization of Wilson's short distance expansion [125]. For scalar currents $j(x) = \phi(x)\phi(x)$ the local OPE is given by [31]:

$$T(j(x)j(0)) \underset{x^2 \to 0}{\approx} \sum_{n=0}^{\infty} C_n(x^2) x^{\mu_1} \dots x^{\mu_n} \, O_{\mu_1 \dots \mu_n}(0) + \text{higher order terms.} \tag{2.1.14}$$

Here the considered operator product is represented as an infinite sum (tower) of local operators,

$$O_{\mu_1 \dots \mu_n}(0) = \phi(0) \partial_{\mu_1} \dots \partial_{\mu_n} \phi(0), \tag{2.1.15}$$

with canonical dimension d_n which is equal to the scale dimension[1] in free-field theory. Because this operator is completely symmetric and traceless ($x^2 = 0$) on the light-cone it has the maximal Lorentz-spin j. Usually local operators are classified with respect to their (geometrical) twist $\tau = d_n - j$ [69]. Here $C_n(x^2)$ are local singular c-number coefficient functions which can be classified according to their degree of singularity at $x^2 = 0$. Operators of the same singularity at $x^2 = 0$ are of the same importance. Coefficient functions with maximal singularity appear in leading order. A disadvantage of the local LCE is that Eq. (2.1.14) is not a true operator identity. It exists on a dense subspace of the Fock space only [126, 30].

The standard form of nonlocal light-cone expansion in scalar field theory was introduced by Anikin and Zavialov [6]. It is given by the following expression:

$$RT(j(x)j(0)S) \underset{x^2\to 0}{\approx} \int_0^1 \mathrm{d}\kappa_1 \int_0^1 \mathrm{d}\kappa_2\, C(x^2,\kappa_1,\kappa_2)\, O(\kappa_1\tilde{x},\kappa_2\tilde{x}) + \text{higher order terms}, \tag{2.1.16}$$

which represents a true operator identity. A rigorous proof of the nonlocal LCE has been given in [6] for the scalar field theory. This nonlocal OPE holds on the whole Hilbert space in contrast to local OPE which is only justified in a restricted region of the Hilbert space [126, 30]. Here $O(\kappa_1\tilde{x},\kappa_2\tilde{x})$ are bilocal light-ray operators on a straight light-like path between $\kappa_1\tilde{x}$ and $\kappa_2\tilde{x}$. The light-like vector

$$\tilde{x} = x + \rho\,\frac{(x\rho)}{\rho^2}\left(\sqrt{1-\frac{x^2\rho^2}{(x\rho)^2}} - 1\right), \quad \text{with} \quad \tilde{x}^2 = 0 \tag{2.1.17}$$

is related to x and a fixed non-null four-vector ρ whose dependence drops out in leading order expressions. The nonlocal LCE is in some sense an integral or summed up representation of the local LCE. The nonlocal and local operators are related through the following formulas:

$$\begin{aligned} O(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \sum_{n_1,n_2}^{\infty} \frac{\kappa_1^{n_1}}{n_1!}\frac{\kappa_2^{n_2}}{n_2!}\, O_{n_1,n_2}(\tilde{x}), \qquad (2.1.18)\\ O_{n_1,n_2}(\tilde{x}) &= \left(\frac{\partial}{\partial\kappa_1}\right)^{n_1}\left(\frac{\partial}{\partial\kappa_2}\right)^{n_2} O(\kappa_1\tilde{x},\kappa_2\tilde{x})|_{\kappa_1=\kappa_2=0}\,. \end{aligned}$$

In Refs. [84, 60, 106, 8, 9] some aspects of the nonlocal LCE for the case of QCD has been studied. Following the Refs. [60, 106] we use now the nonlocal OPE for the case of QCD. In this way we consider the renormalized (R) time-ordered (T) product of two electromagnetic currents

$$RT\big(J^\mu(x/2)J^\nu(-x/2)\mathcal{S}\big), \tag{2.1.19}$$

where the renormalized S-matrix is given by

$$\mathcal{S} = RT\,\exp\left(\mathrm{i}\int \mathrm{d}^4x\,\mathcal{L}_{\text{eff}}\right). \tag{2.1.20}$$

Here $\mathcal{L}_{\text{eff}}$ denotes the effective Lagrangian of QCD [74]

$$\mathcal{L}_{\text{eff}} = -\frac{1}{4}F^a_{\mu\nu}F^{a\mu\nu} + \bar{\psi}\left(\mathrm{i}\gamma^\mu D_\mu(A) - m\right)\psi - \frac{1}{2\alpha}\left(\partial^\mu A^a_\mu\right)^2 + \bar{c}^a\partial^\mu D^{ab}_\mu(A)c^b, \tag{2.1.21}$$

---

[1] The scale dimension of the local operator $O_{\mu_1\cdots\mu_n}(y)$ is given by the transformation behaviour with respect to dilations at the point $y = 0$.

which contains the gluon field $A^a_\mu(x)$, the ghost fields $c^a(x)$ and $\bar{c}^a(x)$, the quark field $\psi(x)$, and α is the gauge fixing parameter. The indices a, b are used for the field and the covariant derivative $D^{ab}_\mu(A)$ in the adjoint representation of the gauge group $SU(3)_{\text{colour}}$. On the other hand, the quark fields and the covariant derivative

$$D_\mu(A) = \partial_\mu + \mathrm{i}gA^a_\mu(x)t^a \tag{2.1.22}$$

are given in the fundamental representation of this gauge group.

The electromagnetic current of the hadrons reads

$$J^\mu(x) = \bar{\psi}(x)\gamma^\mu \frac{1}{2}\left(\lambda^3_f + \frac{1}{\sqrt{3}}\lambda^8_f\right)\psi(x), \qquad \text{for} \qquad SU(3)_{\text{flavour}}. \tag{2.1.23}$$

In lowest order in the coupling constant ($\mathcal{S} = 1$) we obtain for the T-product of the two vector currents

$$\begin{aligned} T\big(J^\mu(x/2)J^\nu(-x/2)\big) = -\bar{\psi}(x/2)\gamma^\mu \mathrm{i}S_F(x)\gamma^\nu c^a\lambda^a_f\psi(-x/2) - \bar{\psi}(-x/2)\gamma^\nu \mathrm{i}S_F(x)\gamma^\mu c^a\lambda^a_f\psi(x/2) \\ + \text{higher order terms,} \end{aligned} \tag{2.1.24}$$

where

$$S_F(x) \approx -\frac{\not{x}}{2\pi^2(x^2-\mathrm{i}0)^2} - \frac{\mathrm{i}m_q}{4\pi^2(x^2-\mathrm{i}0)} + \mathcal{O}(m_q^2) \tag{2.1.25}$$

is the free quark-propagator in the x-space near the light-cone with leading quark mass correction. Obviously, the first part in Eq. (2.1.25) is the massless quark-propagator. The $SU(3)_{\text{flavour}}$ vector

$$c^a = \left(\frac{2}{9}\delta^{a0} + \frac{1}{6}\delta^{a3} + \frac{1}{6\sqrt{3}}\delta^{a8}\right)e^2 \tag{2.1.26}$$

determines the flavour content and e denotes the electric charge. Here λ^a_f is a generator of the flavour group corresponding to the considered hadron. If we use the Chisholm-identity,

$$\gamma^\mu\gamma^\alpha\gamma^\nu = \left(g^{\mu\alpha}g^{\nu\beta} + g^{\mu\beta}g^{\nu\alpha} - g^{\mu\nu}g^{\alpha\beta}\right)\gamma_\beta - \mathrm{i}\epsilon^{\mu\nu\alpha\beta}\gamma^5\gamma_\beta, \tag{2.1.27}$$

where $\epsilon^{\mu\nu\alpha\beta}$ denotes the Levi-Civita symbol, we obtain for the massless part

$$\begin{aligned} T\big(J^\mu(x/2)J^\nu(-x/2)\big) = \frac{c^a x_\alpha}{2\pi^2(x^2-\mathrm{i}0)^2}\Big\{\epsilon^{\mu\nu\alpha\beta}\left(\bar{\psi}(x/2)\gamma^5\gamma_\beta\lambda^a_f\psi(-x/2) + \bar{\psi}(-x/2)\gamma^5\gamma_\beta\lambda^a_f\psi(x/2)\right) \\ + \mathrm{i}S^{\mu\nu\alpha\beta}\left(\bar{\psi}(x/2)\gamma_\beta\lambda^a_f\psi(-x/2) - \bar{\psi}(-x/2)\gamma_\beta\lambda^a_f\psi(x/2)\right)\Big\} + \ldots, \end{aligned} \tag{2.1.28}$$

The nonlocal LCE of QCD-operators is given by means of the T-product of two vector currents as an integral representation of two auxiliary variables κ_i:

$$\begin{aligned} T\big(J^\mu(x/2)J^\nu(-x/2)\big) \underset{x^2\to 0}{\approx} \int_{-1}^{1}\mathrm{d}\kappa_1\int_{-1}^{1}\mathrm{d}\kappa_2\Big(C^a(x^2,\kappa_1,\kappa_2,\mu^2)S^{\mu\nu\alpha\beta}\tilde{x}_\alpha(O^a_\beta(\kappa_1,\kappa_2) - O^a_\beta(\kappa_2,\kappa_1)) \\ - C^{a,5}(x^2,\kappa_1,\kappa_2,\mu^2)\mathrm{i}\epsilon^{\mu\nu\alpha\beta}\tilde{x}_\alpha(O^{a,5}_\beta(\kappa_1,\kappa_2) + O^{a,5}_\beta(\kappa_2,\kappa_1))\Big) + \ldots \end{aligned} \tag{2.1.29}$$

with the coefficient functions which in Born approximation read:

$$C^a(x^2,\kappa_1,\kappa_2,\mu^2) = \frac{\mathrm{i}c^a}{(2\pi)^2(x^2-\mathrm{i}0)^2}\left(\delta(\kappa_1-1/2)\delta(\kappa_2+1/2)-\delta(\kappa_2-1/2)\delta(\kappa_1+1/2)\right)$$
$$C^{a,5}(x^2,\kappa_1,\kappa_2,\mu^2) = \frac{\mathrm{i}c^a}{(2\pi)^2(x^2-\mathrm{i}0)^2}\left(\delta(\kappa_1-1/2)\delta(\kappa_2+1/2)+\delta(\kappa_2-1/2)\delta(\kappa_1+1/2)\right).$$

In the flavour non-singlet case $\lambda_f^a \neq 1$, the bilocal operators appearing in the operator-product expansion of two electromagnetic currents are the gauge-invariant (vector) quark operators:

$$O^a_\alpha(\kappa_1,\kappa_2) = \bar\psi(\kappa_1\tilde x)\gamma_\alpha\lambda^a U(\kappa_1\tilde x,\kappa_2\tilde x)\psi(\kappa_2\tilde x) \tag{2.1.30}$$
$$O^{a,5}_\alpha(\kappa_1,\kappa_2) = \bar\psi(\kappa_1\tilde x)\gamma^5\gamma_\alpha\lambda^a U(\kappa_1\tilde x,\kappa_2\tilde x)\psi(\kappa_2\tilde x), \tag{2.1.31}$$

where λ^a are the generators of the flavour group $SU_f(N)$. To guarantee the gauge invariance both operators contain the path-ordered phase factor along the straight line connecting the points $\kappa_1\tilde x$ and $\kappa_2\tilde x$

$$U(\kappa_1\tilde x,\kappa_2\tilde x) = P\exp\left\{\mathrm{i}g\int_{\kappa_1}^{\kappa_2}\mathrm{d}\tau\,\tilde x^\mu A_\mu(\tau\tilde x)\right\}, \tag{2.1.32}$$

where P denotes the path ordering, g is the strong coupling parameter and $A_\mu = A_\mu^a t^a$ with t^a being the generators of $SU(3)_{\mathrm{color}}$ in the fundamental representation spanned by the quark fields ψ.

In the flavour singlet case ($\lambda_f^0 \equiv 1$), the situation is much more complicated. The quark operators (2.1.30) mix with operators that are made of two gluon fields (bilocal gluon operators):

$$G_{\alpha\beta}(\kappa_1,\kappa_2) = F^{a\rho}_\alpha(\kappa_1\tilde x)U^{ab}(\kappa_1\tilde x,\kappa_2\tilde x)F^b_{\rho\beta}(\kappa_2\tilde x) \tag{2.1.33}$$
$$G^5_{\alpha\beta}(\kappa_1,\kappa_2) = F^{a\rho}_\alpha(\kappa_1\tilde x)U^{ab}(\kappa_1\tilde x,\kappa_2\tilde x)\widetilde F^b_{\rho\beta}(\kappa_2\tilde x). \tag{2.1.34}$$

Here

$$F_{\alpha\beta} \equiv F^a_{\alpha\beta}t^a = \partial_\alpha A_\beta - \partial_\beta A_\alpha - \mathrm{i}g[A_\alpha, A_\beta], \qquad \widetilde F_{\alpha\beta} = \frac{1}{2}\,\epsilon_{\alpha\beta\mu\nu}F^{\mu\nu} \tag{2.1.35}$$

is the field strength and the dual field strength of gluons. Additionally, $U^{ab}(\kappa_1\tilde x,\kappa_2\tilde x)$ is the phase factor (2.1.32) in the adjoint representation. It should be noted that these operators are a complete set of operators in the massless quark theory $m_q = 0$.

Now we calculate the part which is given by the second part of Eq. (2.1.25). It is known that for $m_q \neq 0$ there appears a further set of gauge invariant operators [88, 96]. If we use the relation

$$\begin{aligned}\gamma^\mu\gamma^\nu &= g^{\mu\nu} - \mathrm{i}\sigma^{\mu\nu}\\ &= g^{\mu\nu} + \frac{1}{2}\epsilon^{\mu\nu\alpha\beta}\gamma_5\sigma_{\alpha\beta},\end{aligned} \tag{2.1.36}$$

we obtain

$$\begin{aligned}T\big(J^\mu(x/2)J^\nu(-x/2)\big) &= (2.1.28)\\ &- \frac{m_q c^a}{8\pi^2(x^2-\mathrm{i}0)}\Big\{g^{\mu\nu}\left(\bar\psi(x/2)\lambda^a_f\psi(-x/2)+\bar\psi(-x/2)\lambda^a_f\psi(x/2)\right)\\ &- \epsilon^{\mu\nu\alpha\beta}\left(\bar\psi(x/2)\gamma^5\sigma_{\alpha\beta}\lambda^a_f\psi(-x/2)-\bar\psi(-x/2)\gamma^5\sigma_{\alpha\beta}\lambda^a_f\psi(x/2)\right)\Big\}+\dots,\end{aligned} \tag{2.1.37}$$

and in terms of the nonlocal LCE:

$$T\big(J^\mu(x/2)J^\nu(-x/2)\big) \underset{x^2\to 0}{\approx} (2.1.29)$$
$$- m_q \int_{-1}^{1} \mathrm{d}\kappa_1 \int_{-1}^{1} \mathrm{d}\kappa_2 \Bigg(C_m^a(x^2,\kappa_1,\kappa_2,\mu^2) g^{\mu\nu} (N^a(\kappa_1,\kappa_2) + N^a(\kappa_2,\kappa_1))$$
$$+ C_m^{a,5}(x^2,\kappa_1,\kappa_2,\mu^2) \mathrm{i}\epsilon^{\mu\nu\alpha\beta} (M^{a,5}_{[\alpha\beta]}(\kappa_1,\kappa_2) - M^{a,5}_{[\alpha\beta]}(\kappa_2,\kappa_1))\Bigg) + \ldots \tag{2.1.38}$$

with the coefficient functions:

$$C_m^a(x^2,\kappa_1,\kappa_2,\mu^2) = \frac{c^a}{(4\pi)^2(x^2-\mathrm{i}0)} \left(\delta(\kappa_1-1/2)\delta(\kappa_2+1/2) + \delta(\kappa_2-1/2)\delta(\kappa_1+1/2)\right)$$
$$C_m^{a,5}(x^2,\kappa_1,\kappa_2,\mu^2) = \frac{c^a}{(4\pi)^2(x^2-\mathrm{i}0)} \left(\delta(\kappa_1-1/2)\delta(\kappa_2+1/2) - \delta(\kappa_2-1/2)\delta(\kappa_1+1/2)\right).$$

The additional bilocal operators appearing in the operator-product expansion of two electromagnetic currents are the gauge-invariant quark operators:

$$N^a(\kappa_1,\kappa_2) = \bar\psi(\kappa_1\tilde x)\lambda^a U(\kappa_1\tilde x,\kappa_2\tilde x)\psi(\kappa_2\tilde x) \tag{2.1.39}$$
$$M^{a,5}_{[\alpha\beta]}(\kappa_1,\kappa_2) = \bar\psi(\kappa_1\tilde x)\mathrm{i}\gamma^5\sigma_{\alpha\beta}\lambda^a U(\kappa_1\tilde x,\kappa_2\tilde x)\psi(\kappa_2\tilde x). \tag{2.1.40}$$

In the flavour singlet case, the quark operators (2.1.39) and (2.1.40) which are multiplied by m_q may be mix with operators that are made of two gluon fields:

$$G^\mu{}_\mu(\kappa_1,\kappa_2) = F^{a\mu\nu}(\kappa_1\tilde x)U^{ab}(\kappa_1\tilde x,\kappa_2\tilde x)F^b_{\mu\nu}(\kappa_2\tilde x) \tag{2.1.41}$$
$$G^5_{[\alpha\beta]}(\kappa_1,\kappa_2) = F^{a\rho}{}_{[\alpha}(\kappa_1\tilde x)U^{ab}(\kappa_1\tilde x,\kappa_2\tilde x)\widetilde F^b_{\beta]\rho}(\kappa_2\tilde x). \tag{2.1.42}$$

In general, the coefficient functions $C^a(x^2,\kappa_1,\kappa_2,\mu^2)$ and $C^{a,5}(x^2,\kappa_1,\kappa_2,\mu^2)$ are generalized functions. They are determined perturbatively and can be ordered according to their degree of singularity at $x^2 = 0$. The Fourier transform $F_\Gamma(x^2, xq_i;\mu^2)$ of $C_\Gamma(x^2,\kappa_i;\mu^2)$ with respect to κ_i are an entire functions of the new arguments $\tilde x q_i$ [6] according to

$$C^a(x^2,\kappa_1,\kappa_2,\mu^2) = \int \mathrm{d}(\tilde xq_1)\mathrm{d}(\tilde xq_2)\, \mathrm{e}^{-\mathrm{i}\kappa_1(\tilde xq_1)-\mathrm{i}\kappa_2(\tilde xq_2)} F^a(x^2,(\tilde xq_1),(\tilde xq_2),\mu^2)$$
$$C^{a,5}(x^2,\kappa_1,\kappa_2,\mu^2) = \int \mathrm{d}(\tilde xq_1)\mathrm{d}(\tilde xq_2)\, \mathrm{e}^{-\mathrm{i}\kappa_1(\tilde xq_1)-\mathrm{i}\kappa_2(\tilde xq_2)} F^{a,5}(x^2,(\tilde xq_1),(\tilde xq_2),\mu^2). \tag{2.1.43}$$

In Born approximation we obtain (choice of the renormalization point of $\mu^2 = Q^2$):

$$F^a(x^2,(\tilde xq_1),(\tilde xq_2),\mu^2) = \frac{\mathrm{i}c^a}{(2\pi)^2(x^2-\mathrm{i}0)^2}\left(\mathrm{e}^{\mathrm{i}\kappa_1(\tilde xq_1)} - \mathrm{e}^{\mathrm{i}\kappa_2(\tilde xq_2)}\right)$$
$$F^{a,5}(x^2,(\tilde xq_1),(\tilde xq_2),\mu^2) = \frac{\mathrm{i}c^a}{(2\pi)^2(x^2-\mathrm{i}0)^2}\left(\mathrm{e}^{\mathrm{i}\kappa_1(\tilde xq_1)} + \mathrm{e}^{\mathrm{i}\kappa_2(\tilde xq_2)}\right). \tag{2.1.44}$$

The range of κ_i is restricted to $-1 \le \kappa_i \le +1$. Similar relations exist between $F_m^\Gamma(x^2, xq_i;\mu^2)$ of $C_m^\Gamma(x^2,\kappa_i;\mu^2)$.

Finally the Compton amplitude for the general case of non-forward scattering is given by means of the LCE as follows

$$T^{\mu\nu}(P_+,P_-,Q) \underset{x^2\to 0}{\approx} \frac{1}{(2\pi)^2}\int \mathrm{d}^4x \frac{c^a\mathrm{e}^{\mathrm{i}xq}}{(x^2-\mathrm{i}0)^2}\int_{-1}^{1}\mathrm{d}\kappa_1\int_{-1}^{1}\mathrm{d}\kappa_2 \tag{2.1.45}$$
$$\times \Big\{ \mathrm{i}\Delta_+(\kappa_1,\kappa_2)\epsilon^{\mu\nu\alpha\beta}\tilde x_\alpha \big\langle P_2,S_2|O^{a,5}_\beta(\kappa_1,\kappa_2) + O^{a,5}_\beta(\kappa_2,\kappa_1)|P_1,S_1\big\rangle$$
$$- \Delta_-(\kappa_1,\kappa_2)S^{\mu\nu\alpha\beta}\tilde x_\alpha \big\langle P_2,S_2|O^a_\beta(\kappa_1,\kappa_2) - O^a_\beta(\kappa_2,\kappa_1)|P_1,S_1\big\rangle \Big\} + \ldots,$$

where

$$\Delta_{\pm}(\kappa_1,\kappa_2) = \big[\delta(\kappa_1-1/2)\delta(\kappa_2+1/2) \pm \delta(\kappa_2-1/2)\delta(\kappa_2+1/2)\big]. \tag{2.1.46}$$

2.1.3 Distribution amplitudes

Distribution amplitudes of hadrons are the key ingredients of the QCD approach of exclusive, diffractive, and deep inelastic scattering processes. They provide the universal nonperturbative input in physical amplitudes. Accordingly, they have to be addressed using nonperturbative methods.

The parton distribution functions (PDFs) are the DAs in forward scattering processes (DIS) and are usually defined by the Fourier transformation of nucleon matrix elements of light-cone operators. We begin with the simple case – the chiral-even scalar operators on the light-cone[2]

$$O^a(\kappa\tilde{x},-\kappa\tilde{x}) \equiv \bar{\psi}(\kappa\tilde{x})(\tilde{x}\gamma)\lambda^a U(\kappa\tilde{x},-\kappa\tilde{x})\psi(-\kappa\tilde{x}), \tag{2.1.47}$$
$$O^{a,5}(\kappa\tilde{x},-\kappa\tilde{x}) \equiv \bar{\psi}(\kappa\tilde{x})\gamma^5(\tilde{x}\gamma)\lambda^a U(\kappa\tilde{x},-\kappa\tilde{x})\psi(-\kappa\tilde{x}). \tag{2.1.48}$$

In the unpolarized case, the forward matrix element of this operator is given by [78] (see also Section 3.1)

$$\begin{aligned}\langle P|O^a(\kappa\tilde{x},-\kappa\tilde{x})|P\rangle &= 2\,(\tilde{x}P)\int \mathrm{d}z\, \mathrm{e}^{2\mathrm{i}\kappa z(\tilde{x}P)}\, q^a(z,\mu^2)\\ &= 2\,(\tilde{x}P)\sum_{n=0}^{\infty}\frac{(2\mathrm{i}\kappa(\tilde{x}P))^n}{n!}\, q^a_n(\mu^2).\end{aligned} \tag{2.1.49}$$

Here $|P\rangle$ denotes the nucleon state with momentum P and $q^a(z,\mu^2)$ is a parton distribution function[3] which is related to a reduced matrix element $q^a_n(\mu^2)$ by means of a Mellin transformation,

$$q^a_n(\mu^2) = \int \mathrm{d}z\, z^n q^a(z,\mu^2), \tag{2.1.50}$$

with respect to the distribution parameter z with $-1 \leq z \leq 1$. This parton distribution function is defined through the inverse Fourier transformation of Eq. (2.1.49) and reads [106]

$$q^a(z,\mu^2) = \frac{1}{2}\int \frac{\mathrm{d}\kappa}{2\pi}\, \mathrm{e}^{-2\mathrm{i}\kappa z(\tilde{x}P)}\, \frac{|\tilde{x}P|}{(\tilde{x}P)}\, \langle P|O^a(\kappa\tilde{x},-\kappa\tilde{x})|P\rangle|_{\mu^2=Q^2}. \tag{2.1.51}$$

The factor $1/(\tilde{x}P)$ is due to the structure $\tilde{x}\gamma$ of the operator $O^a(\kappa\tilde{x},-\kappa\tilde{x})$ in the matrix element (2.1.49). Physically, this distribution function describes the probability to find quarks with the momentum fraction zP in an unpolarized nucleon. For $z > 0$ ($z < 0$) it can be interpreted as a quark (antiquark) distribution function. The (unpolarized) structure function F_1 is given by the linear combination of quark and antiquark distribution functions in the following way [80]

$$F_1(z,Q^2) = \frac{1}{2}\sum_{a=1}^{N_f} e_a^2\big(q^a(z,Q^2) - q^a(-z,Q^2)\big). \tag{2.1.52}$$

---

[2] In Section 3.1 we will prove that these scalar LC-operators are twist-2 operators.

[3] The PDF $q(z,\mu^2)$ is equal to the PDFs of twist-2 $f_1(z,\mu^2)$ and $F^{(2)}(z,\mu^2)$, see Eqs. (3.1.6) and (3.1.43).

In the polarized case, the forward matrix element of the axial chiral-even bilocal scalar operator is given by by [78]

$$\begin{aligned}\langle PS|O^{a,5}(\kappa\tilde{x},-\kappa\tilde{x})|PS\rangle &= 2\,(\tilde{x}S)\int \mathrm{d}z\, \mathrm{e}^{2\mathrm{i}\kappa z(\tilde{x}P)}\,\Delta q^a(z,\mu^2)\\ &= 2\,(\tilde{x}S)\sum_{n=0}^{\infty}\frac{(2\mathrm{i}\kappa(\tilde{x}P))^n}{n!}\,\Delta q^a_n(\mu^2),\end{aligned}\qquad (2.1.53)$$

where S is the spin vector of the nucleon. This parton distribution function[4] is defined through the inverse Fourier transformation according to

$$\Delta q^a(z,\mu^2) = \frac{1}{2}\int\frac{\mathrm{d}\kappa}{2\pi}\,\mathrm{e}^{-2\mathrm{i}\kappa z(\tilde{x}P)}\,\frac{|\tilde{x}P|}{(\tilde{x}S)}\,\langle P|O^a(\kappa\tilde{x},-\kappa\tilde{x})|P\rangle|_{\mu^2=Q^2}. \qquad (2.1.54)$$

The factor $1/(\tilde{x}S)$ is due to the structure $\gamma^5\tilde{x}\gamma$ of the operator $O^{a,5}(\kappa\tilde{x},-\kappa\tilde{x})$ in the matrix element (2.1.53). Physically, this distribution function describes the probability to find quarks with the momentum fraction zP in a polarized nucleon. The (polarized) structure function G_1 is given in terms of quark and antiquark distribution functions as follows [80]

$$G_1(z,Q^2) = \frac{1}{2}\sum_{a=1}^{N_f} e_a^2\big(\Delta q^a(z,Q^2) + \Delta q^a(-z,Q^2)\big). \qquad (2.1.55)$$

Analogously, we may define the scalar meson distribution amplitude $\hat{\phi}^a(\xi,\mu^2)$ by means of the vacuum to meson matrix element of the bilocal operator [106]:

$$\hat{\phi}^a(\xi,\mu^2) = \int\frac{\mathrm{d}\kappa}{2\pi}\,\mathrm{e}^{-\mathrm{i}\kappa\xi(\tilde{x}P)}\,\frac{|\tilde{x}P|}{(\tilde{x}P)}\,\langle 0|O^a(\kappa\tilde{x},-\kappa\tilde{x})|P\rangle|_{\mu^2=Q^2}. \qquad (2.1.56)$$

Here $|P\rangle$ is the one-particle state of a scalar meson and P denotes the momentum of the scalar meson. The meson distribution amplitude is the probability amplitude to find a quark-antiquark pair in the meson in dependence of the momentum fraction parameter ξ and the momentum transfer Q^2.

Consequently, in the unpolarized case, the nonforward matrix element of the scalar operator on the light-cone is given by [106]

$$\begin{aligned}\langle P_2|O^a(\kappa\tilde{x},-\kappa\tilde{x})|P_1\rangle &= (\tilde{x}P_+)\int \mathrm{d}z_+\int \mathrm{d}z_-\,\mathrm{e}^{\mathrm{i}\kappa\tilde{x}(P_+z_+ + P_-z_-)}\,f^a(z_+,z_-,\mu^2)\\ &= (\tilde{x}P_+)\sum_{n=0}^{\infty}\frac{(\mathrm{i}\kappa)^n}{n!}\sum_{m=0}^{n}\binom{n}{m}(\tilde{x}P_+)^{n-m}(\tilde{x}P_-)^m\,f^a_{nm}(\mu^2),\\ &= \sum_{n=0}^{\infty}\frac{(\mathrm{i}\kappa)^n}{n!}\,(\tilde{x}P_+)^{n+1}\sum_{m=0}^{n}\binom{n}{m}\tilde{\tau}^m\,f^a_{nm}(\mu^2),\end{aligned}\qquad (2.1.57)$$

where

$$\tilde{\tau} = \frac{\tilde{x}P_-}{\tilde{x}P_+},\quad P_+z_+ + P_-z_- = P_1z_1 + P_2z_2 \quad\text{with}\quad z_\pm = \frac{1}{2}(z_2 \pm z_1), \qquad (2.1.58)$$

---

[4] Here $\Delta q(z,\mu^2)$ is equivalent to the PDFs of twist-2 $g_1(z,\mu^2)$ and $G^{(2)}(z,\mu^2)$, see Eqs. (3.1.5) and (3.1.35).

and $\binom{n}{m} = n!/(n-m)!m!$ is the binomial coefficient. Here $f^a(z_+, z_-, \mu^2)$ is a double distribution function which is related to a reduced matrix element $f^a_{nm}(\mu^2)$ by the help of a twofold Mellin transformation,

$$f^a_{nm}(\mu^2) = \int \mathrm{d}z_+ \int \mathrm{d}z_- \, z_+^{n-m} \, z_-^m f^a(z_+, z_-, \mu^2), \tag{2.1.59}$$

with respect to the distribution parameters z_+ and z_- and the integration domain is: $-1 \leq z_+ \leq 1$, $-1 + |z_+| \leq z_- \leq 1 - |z_+|$. Hence, $f^a_{nm}(\mu^2)$ are double moments of the double distribution function $f^a(z_+, z_-, \mu^2)$ which only depend on the distribution parameters and not on kinematical factors like P_+, P_-, etc.. The double distributions are related to the ordinary parton distributions. By taking the matrix element (2.1.57) with $P_1 = P_2 = P$, we obtain [114, 65]

$$q^a(z, \mu^2) = \int \mathrm{d}z_- f^a(z_+ = z, z_-, \mu^2). \tag{2.1.60}$$

But on the other hand we may parametrize the matrix element in terms of a skewed parton distribution function [83, 114, 115, 22, 23]:

$$\begin{aligned}\langle P_2 | O^a(\kappa\tilde{x}, -\kappa\tilde{x}) | P_1 \rangle &= (\tilde{x}P_+) \int_{-1}^{1} \mathrm{d}z \, \mathrm{e}^{\mathrm{i}\kappa\tilde{x}P_+ z} \, H^a(z, \tilde{\tau}, \mu^2) \\ &= \sum_{n=0}^{\infty} \frac{(\mathrm{i}\kappa)^n}{n!} \, (\tilde{x}P_+)^{n+1} H^a_n(\tilde{\tau}, \mu^2),\end{aligned} \tag{2.1.61}$$

where $H^a(z, \tilde{\tau}, \mu^2)$ is the skewed parton distribution function[5] and $H^a_n(\tilde{\tau}, \mu^2)$ are the corresponding moments which depend on the variable $\tilde{\tau}$. By comparing Eqs. (2.1.57) and (2.1.61) we observe the following relation [114] between the skewed parton distribution $H^a(z, \tilde{\tau}, \mu^2)$ and the double distribution $f^a(z_+, z_-, \mu^2)$:

$$H^a_n(\tilde{\tau}, \mu^2) = \int_{-1}^{1} \mathrm{d}z \, z^n H^a(z, \tilde{\tau}, \mu^2) = \sum_{m=0}^{n} \binom{n}{m} \tilde{\tau}^m \int \mathrm{d}z_+ \int \mathrm{d}z_- \, z_+^{n-m} \, z_-^m f^a(z_+, z_-, \mu^2). \tag{2.1.62}$$

Thus, skewed parton distributions are obtained from double distributions after an appropriate integration (see Eq. (2.1.62)). Additionally, the skewed parton distributions are related to the ordinary parton distributions. In fact, we have

$$H^a(z, 0, \mu^2) = q^a(z, \mu^2). \tag{2.1.63}$$

Similar relations can be derived for the double distribution function $\Delta f^a(z_+, z_-, \mu^2)$, the skewed parton distribution $\widetilde{H}^a(z, \tilde{\tau}, \mu^2)$, and the parton distribution $\Delta q^a(z, \mu^2)$ which are corresponding to the axial operators $O^{a,5}(\kappa\tilde{x}, -\kappa\tilde{x})$.

Let me note that we will discuss double distribution amplitudes as off-cone nonforward matrix elements of more complicated operators with definite geometric twist in Section 3.3.

---

[5] Ji [83] originally introduced a skewed parton distribution $H(z, \tilde{\tau}, P_-^2)$ which depends on the kinematical factor P_-^2. Forming the first moment of this skewed distribution, one gets the sum rule [83]: $\int_{-1}^{1} \mathrm{d}z H(z, \tilde{\tau}, P_-^2) = F_1(P_-^2)$, where $F_1(P_-^2)$ is the Dirac form factor. In Section 3.3 we will see that the kinematical factors like P_-^2 are proportional to x^2 and can be generated through the twist projectors of geometric twist.

2.1.4 Anomalous dimensions of QCD operators

We discuss now some facts about the anomalous dimension of (twist-2) QCD-operators. Physically, the Q^2-evolution of DAs is governed by the anomalous dimensions, which are determined by the renormalization of the corresponding operators. The anomalous dimensions are necessary to connect and compare experimental data which are obtained at different values of Q^2.

For simplicity we symbolically use the notation $\mathcal{O}_n(\tilde{x})$ for any local operator on the light-cone (see Section 2.2). Here $\mathcal{O}_n^{\text{bare}}(\tilde{x})$ denotes the unrenormalized operator and the renormalized operator $\mathcal{O}_n(\tilde{x};\mu^2)$ is given by

$$\mathcal{O}_n(\tilde{x};\mu^2) = \sum_{n'=0}^{n} Z^{-1}_{nn'}(\mu^2)\mathcal{O}^{\text{bare}}_{n'}(\tilde{x}), \tag{2.1.64}$$

where $Z_{nn'}(\mu^2)$ is the renormalization matrix and μ^2 is the renormalization point. The invariance of the physics under the choice of the point of renormalization can be characterized by the so-called renormalization group equation

$$\mu\frac{\mathrm{d}}{\mathrm{d}\mu}\,\mathcal{O}_n(\tilde{x};\mu^2) = \sum_{n'=0}^{n} \gamma_{nn'}(g(\mu^2))\mathcal{O}_{n'}(\tilde{x};\mu^2), \tag{2.1.65}$$

where $\gamma_{nn'}(g(\mu^2))$ is the (local) anomalous dimension of the operator $\mathcal{O}_n(\tilde{x};\mu^2)$. It is defined by

$$\gamma_{nn'}(g(\mu^2)) = Z^{-1}_{nm}\,\mu\frac{\partial}{\partial\mu}\,Z_{mn'}(\mu^2). \tag{2.1.66}$$

In this connection, the β-function is introduced:

$$\beta(g(\mu^2)) = \mu\frac{\partial}{\partial\mu}\,g(\mu^2). \tag{2.1.67}$$

The scale dependence of the operator is obtained by the solution of the renormalization group equation as

$$\mathcal{O}_n(\tilde{x};Q^2) = \sum_{n'=0}^{n} \mathcal{O}_{n'}(\tilde{x};\mu^2)\,\exp\left[-\int_{g(\mu^2)}^{g(Q^2)} \mathrm{d}g\,\frac{\gamma_{nn'}(g)}{\beta(g)}\right]. \tag{2.1.68}$$

The anomalous dimension and the β-function are calculable using perturbation theory. They can be expanded as power series in g^2 as

$$\gamma_{nn'}(g) = \frac{g^2}{16\pi^2}\,\gamma^0_{nn'} + \frac{g^4}{(16\pi^2)^2}\,\gamma^1_{nn'} + O(g^6), \tag{2.1.69}$$

$$\beta(g) = -\frac{g^3}{16\pi^2}\,\beta^0 - \frac{g^5}{(16\pi^2)^2}\,\beta^1 + O(g^7), \tag{2.1.70}$$

where the coefficients of the β functions are:

$$\beta_0 = 11 - \frac{2}{3}\,N_f, \qquad \beta_1 = 102 - \frac{38}{3}\,N_f, \tag{2.1.71}$$

with the number of quark flavours N_f.

If we build the forward matrix element of the local operator, the renormalization group equation of the corresponding quark distribution is obtained as ($\gamma_n \equiv \gamma_{nn}$)

$$\mu \frac{\mathrm{d}}{\mathrm{d}\mu}\, q_n^a(\mu^2) = \gamma_n^{qq}(g(\mu^2))q_n^a(\mu^2). \tag{2.1.72}$$

In the flavour singlet case, the quark and gluon operators mix with each other under renormalization. The renormalization group equation for the local operators[6]

$$O_n(\tilde{x}) = \bar{\psi}(0)(\tilde{x}\gamma)(\tilde{x}\overleftrightarrow{D})^{n-1}\psi(0), \tag{2.1.73}$$

$$G_n(\tilde{x}) = \tilde{x}^\mu \tilde{x}^\nu F_\mu{}^\rho(0)(\tilde{x}\overleftrightarrow{D})^{n-2}F_{\rho\nu}(0) \tag{2.1.74}$$

is given by

$$\mu \frac{\mathrm{d}}{\mathrm{d}\mu} \begin{pmatrix} O_n(\tilde{x};\mu^2) \\ G_n(\tilde{x};\mu^2) \end{pmatrix} = \sum_{n'=0}^{n} \begin{pmatrix} \gamma_{nn'}^{qq} & \gamma_{nn'}^{qG} \\ \gamma_{nn'}^{Gq} & \gamma_{nn'}^{GG} \end{pmatrix} \begin{pmatrix} O_{n'}(\tilde{x};\mu^2) \\ G_{n'}(\tilde{x};\mu^2) \end{pmatrix} \tag{2.1.75}$$

where the local (one-loop) unpolarized anomalous dimensions are [34]:

$$\gamma_{nn'}^{qq} = \frac{g^2}{4\pi^2}\, C_F \bigg\{ \bigg[\frac{1}{2} - \frac{1}{(n+1)n} + 2\sum_{j=2}^{n} \frac{1}{j} \bigg] \delta_{nn'} - \bigg[\frac{1}{(n+1)n} + \frac{2}{n-n'}\frac{n'}{n} \bigg] \theta_{nn'} \bigg\} \tag{2.1.76}$$

$$\gamma_{nn'}^{qG} = -\frac{g^2}{4\pi^2}\, N_f T_R\, \frac{2}{(n+2)(n+1)n} \Big[(n^2+n+2) - (n-n')n \Big], \tag{2.1.77}$$

$$\gamma_{nn'}^{Gq} = -\frac{g^2}{4\pi^2}\, C_F\, \frac{1}{n(n^2-1)} \Big[(n^2+n+2)\delta_{nn'} + 2\theta_{nn'} \Big], \tag{2.1.78}$$

$$\begin{aligned} \gamma_{nn'}^{GG} = \frac{g^2}{4\pi^2}\, C_A \bigg\{ &\bigg[\frac{1}{6} - \frac{2}{n(n-1)} - \frac{2}{(n+2)(n+1)} + 2\sum_{j=2}^{n} \frac{1}{j} + \frac{2N_f T_R}{3C_A} \bigg] \delta_{nn'} \\ &+ \bigg[2\bigg(\frac{2n-1}{n(n-1)} - \frac{1}{n-n'} \bigg) - (n-n'+2)\bigg(\frac{1}{n(n-1)} + \frac{1}{(n+2)(n+1)} \bigg) \bigg] \theta_{nn'} \bigg\}, \end{aligned} \tag{2.1.79}$$

where $C_F = (N_c^2-1)/2N_c \equiv 4/3$, $T_R = 1/2$, $C_A = N_c \equiv 3$. We used the following notation

$$\theta_{n\,n'} = \begin{cases} 1 & \text{for } n' < n, \\ 0 & \text{otherwise .} \end{cases}$$

The local (axial) operators

$$O_n^5(\tilde{x}) = \bar{\psi}(0)\gamma^5(\tilde{x}\gamma)(\tilde{x}\overleftrightarrow{D})^{n-1}\psi(0), \tag{2.1.80}$$

$$G_n^5(\tilde{x}) = \tilde{x}^\mu \tilde{x}^\nu F_\mu{}^\rho(0)(\tilde{x}\overleftrightarrow{D})^{n-2}\widetilde{F}_{\rho\nu}(0) \tag{2.1.81}$$

satisfy the renormalization group equation

$$\mu \frac{\mathrm{d}}{\mathrm{d}\mu} \begin{pmatrix} O_n^5(\tilde{x};\mu^2) \\ G_n^5(\tilde{x};\mu^2) \end{pmatrix} = \sum_{n'=0}^{n} \begin{pmatrix} \Delta\gamma_{nn'}^{qq} & \Delta\gamma_{nn'}^{qG} \\ \Delta\gamma_{nn'}^{Gq} & \Delta\gamma_{nn'}^{GG} \end{pmatrix} \begin{pmatrix} O_{n'}^5(\tilde{x};\mu^2) \\ G_{n'}^5(\tilde{x};\mu^2) \end{pmatrix} \tag{2.1.82}$$

---

[6] The local operators are obtained after local expansion of the nonlocal ones (see Section 2.2).

with the local (one-loop) polarized anomalous dimensions [27]:

$$\Delta\gamma^{qq}_{nn'} = \gamma^{qq}_{nn'}, \tag{2.1.83}$$

$$\Delta\gamma^{qG}_{nn'} = -\frac{g^2}{4\pi^2}\, N_f T_R\, \frac{2(n'-1)}{(n+1)n}, \tag{2.1.84}$$

$$\Delta\gamma^{Gq}_{nn'} = \frac{g^2}{4\pi^2}\, C_F\, \frac{1}{(n+1)n}\Big[(n+2)\delta_{nn'} - \frac{2}{n-1}\,\theta_{nn'}\Big], \tag{2.1.85}$$

$$\begin{aligned}\Delta\gamma^{GG}_{nn'} = \frac{g^2}{4\pi^2}\, C_A\bigg\{&\bigg[\frac{1}{6} - \frac{4}{(n+1)n} + 2\sum_{j=2}^{n}\frac{1}{j} + \frac{2N_f T_R}{3C_A}\bigg]\delta_{nn'} \\ &+ \bigg[2\bigg(\frac{2n-1}{n(n-1)} - \frac{1}{n-n'}\bigg) - (n-n'+2)\,\frac{2}{(n+1)n}\bigg]\theta_{nn'}\bigg\}.\end{aligned} \tag{2.1.86}$$

The polarized and unpolarized forward anomalous dimensions [70, 57, 2] are obtained as the diagonal elements of Eqs. (2.1.76) – (2.1.79), (2.1.83) – (2.1.86).[7]

The nonlocal anomalous dimensions are obtained from the local ones by means of an inverse Mellin transformation. The corresponding nonlocal twist-2 anomalous dimensions calculated in the framework of nonlocal OPE are given in Refs. [34, 24, 27, 9, 12, 114].

---

[7] This (diagonal) form of the anomalous dimensions is also valid for the so-called conformal operators (see Refs. [50, 116, 102, 108, 44, 45]).

2.2 The concept of geometric twist of nonlocal operators

In this Section we present a straightforward scheme to discuss nonlocal operators with respect to definite geometric twist. This method is based by applying the representation theory of the Lorentz group. The advantages of this method are: First, the nonlocal operators of definite geometric twist are Lorentz covariant tensors (contrary to the phenomenological concept of twist). Second, this twist decomposition is unique, process– and model–independent. Furthermore, the twist decomposition is independent from the dimension $2h$ of spacetime. Up to now, no clear scheme for higher geometric twist was worked out in the literature.

On the other hand, Jaffe and Ji [78, 79] proposed the notion of *dynamical* twist (t) by counting powers Q^{2-t} which is directly related to the power by which the corresponding distributions contribute to the scattering amplitudes. Additionally, the dynamical twist is only defined for the *matrix elements* of operators and is not Lorentz invariant.

If one uses QCD operators with definite geometric twist, the different phenomenological distribution functions of definite geometric twist are given as matrix elements of these operators. It should be emphasized that the LC operators of definite geometric twist are to be preferred also from the point of view of renormalization theory. Therefore, we are using the concept of geometric twist instead of dynamical twist.

2.2.1 The procedure of twist decomposition of nonlocal operators

This Section is devoted to explain how the QCD operators may by decomposed according to their (geometric) twist which is defined [69]:

$$\text{twist}\,(\tau) \;=\; \text{canonical dimension}\,(d) - \text{Lorentz spin}\,(j).$$

We use the notion of twist because it is a straightforward scheme to classify DAs and, additionally, operators with equal twist mix with each other under renormalization.

The nonlocal operators which are relevant for the QCD processes (see previous Section) are obtained by the nonlocal LC expansion [6] of the (renormalized) time-ordered product of two electromagnetic hadronic currents:

$$RT\left(J^\mu(y+\tfrac{x}{2})J^\nu(y-\tfrac{x}{2})S\right) \underset{x^2\to 0}{\approx} \int \mathrm{d}^2\underline{\kappa}\, C_\Gamma^{\mu\nu}(x^2,\underline{\kappa}) RT\big(\mathcal{O}^\Gamma(y+\kappa_1\tilde{x}, y+\kappa_2\tilde{x})S\big) + \cdots\,, \qquad (2.2.1)$$

where $\Gamma = \{1, \gamma_\mu, \sigma_{\mu\nu}; \gamma_5, \gamma_\mu\gamma_5, \sigma_{\mu\nu}\gamma_5\}$ indicates the tensor structure of the nonlocal QCD operators. Strictly speaking, the expression in Eq. (2.2.1) is an operator valued distribution.

In general, the twist decomposition of an arbitrary nonlocal operator may be formulated for any $2h$-dimensional spacetime. Let us denote such operators for arbitrary values of x as follows:

$$\mathcal{O}^\Gamma(\kappa_1 x, \kappa_2 x) = \Phi'(\kappa_1 x)\Gamma U(\kappa_1 x, \kappa_2 x)\Phi(\kappa_2 x), \qquad (2.2.2)$$

where, suppressing any indices indicating the group representations, Φ generically denotes the various local fields with a scale dimension[8], e.g., scalar ($d_\varphi = h-1$), Dirac spinor ($d_\psi = h - \frac{1}{2}$) as well as gauge field strength ($d_F = h$). The gauge potential A_μ has dimension $d_A = h-1$. Furthermore, Γ labels the tensor structure as well as additional quantum numbers, if necessary. Now, the twist decomposition of operators (2.2.2) consists of the following steps:

---

[8] The scale dimension, d_Φ, of a field Φ is determined from the infinitesimal transformation law under dilations at the point $y=0$: $[\hat{D}, \Phi(y)]|_{y=0} = \mathrm{i} d_\Phi\, \Phi(0)$, where $\hat{D}$ is the dilation generator. For the unrenormalized operators the scale dimension agrees with the canonical dimension. For the renormalized operators the difference from the canonical one is called anomalous dimension.

(1) *Taylor expansion* of the nonlocal operators for *arbitrary* values of x at the point y into an infinite tower of local tensor operators having definite rank n and canonical dimension d. If we choose the expansion point $y = 0$, we obtain the expansion:

$$\begin{aligned}\mathcal{O}^\Gamma(\kappa_1 x, \kappa_2 x) &= \sum_{n=0}^{\infty} \frac{1}{n!} x^{\mu_1} \ldots x^{\mu_n} \Big[\Phi'(y)\Gamma \overset{\leftrightarrow}{D}_{\mu_1}(\kappa_1, \kappa_2) \ldots \overset{\leftrightarrow}{D}_{\mu_n}(\kappa_1, \kappa_2)\Phi(y)\Big]_{y=0} \\ &\equiv \sum_{n=0}^{\infty} \frac{1}{n!} \Big[\Phi'(y)\Gamma\big(x\overset{\leftrightarrow}{D}(\kappa_1, \kappa_2)\big)^n \Phi(y)\Big]_{y=0} \end{aligned} \tag{2.2.3}$$

with the generalized covariant derivatives

$$\begin{aligned} \overset{\leftrightarrow}{D}_\mu(\kappa_1, \kappa_2) \;&\equiv\; \kappa_1 \overset{\leftarrow}{D}_\mu + \kappa_2 \vec{D}_\mu\,, \\ \vec{D}_\mu = \overset{\rightarrow}{\partial}{}^y_\mu + \mathrm{i}gA_\mu(y), &\qquad \overset{\leftarrow}{D}_\mu = \overset{\leftarrow}{\partial}{}^y_\mu - \mathrm{i}gA_\mu(y)\,. \end{aligned} \tag{2.2.4}$$

We introduce, for later convenience, the variables

$$\kappa_\pm = \frac{1}{2}(\kappa_2 \pm \kappa_1) \quad \text{with} \quad \kappa_{1,2} = \kappa_+ \mp \kappa_-\,, \tag{2.2.5}$$

and the following derivatives

$$D^\mu_\pm = \vec{D}{}^\mu \pm \overset{\leftarrow}{D}{}^\mu, \tag{2.2.6}$$

with[9]

$$D^\mu_+ \equiv \partial^\mu_+ = \overset{\leftarrow}{\partial}{}^\mu_y + \overset{\rightarrow}{\partial}{}^\mu_y, \quad D^\mu_- \equiv \overset{\leftrightarrow}{D}{}^\mu = -\overset{\leftarrow}{\partial}{}^\mu_y + \overset{\rightarrow}{\partial}{}^\mu_y + 2\mathrm{i}gA^\mu(y). \tag{2.2.7}$$

Now we are able to rewrite Eq. (2.2.3) in terms of these variables and derivatives according to

$$\begin{aligned}\mathcal{O}^\Gamma(\kappa_1 x, \kappa_2 x) &= \sum_{n=0}^{\infty} \frac{1}{n!} \Big[\Phi'(y)\Gamma(\kappa_- xD_- + \kappa_+ xD_+)^n \Phi(y)\Big]_{y=0} \\ &= \sum_{n=0}^{\infty}\sum_{k=0}^{n} \frac{\kappa_+^k\, \kappa_-^{n-k}}{(n-k)!k!}\, \mathcal{O}^\Gamma_{nk}(y, x)|_{y=0}, \end{aligned} \tag{2.2.8}$$

where we introduced the following local operator

$$\mathcal{O}^\Gamma_{nk}(y, x) = \Phi'(y)\Gamma\big(xD_+\big)^k \big(xD_-\big)^{n-k}\Phi(y). \tag{2.2.9}$$

The local operators (2.2.9) mix under renormalization for the same value of k. Therefore, their anomalous are independent of k [34]. Another possibility of local expansion is to choose the expansion point $y = \kappa_+ x$. We get

$$\begin{aligned}\mathcal{O}^\Gamma(\kappa_1 x, \kappa_2 x) &\equiv \mathcal{O}^\Gamma(\kappa_+ x - \kappa_- x, \kappa_+ x + \kappa_- x) \\ &= \sum_{n=0}^{\infty} \frac{\kappa_-^n}{n!}\, \mathcal{O}^\Gamma_n(y, x)|_{y=\kappa_+ x}, \end{aligned} \tag{2.2.10}$$

---

[9] The derivative ∂_+ is sometimes called “total” derivative [9].

with the local operator:

$$\mathcal{O}_n^\Gamma(y,x) = \Phi'(y)\Gamma\big(xD_-\big)^n\Phi(y). \tag{2.2.11}$$

In this way the variables $\kappa_+ x$ and $2\kappa_- x$, respectively, are centre and the relative coordinates of the two points $\kappa_1 x$ and $\kappa_2 x$. Now we discuss the bilocal operator $\mathcal{O}^\Gamma(-\kappa x, \kappa x)$ which is centred around $y = 0$. For the local expansion of the "centred" operator we obtain the simpler formula

$$\mathcal{O}^\Gamma(-\kappa x, \kappa x) = \sum_{n=0}^{\infty} \frac{\kappa^n}{n!}\,\mathcal{O}_n^\Gamma(y,x)|_{y=0}, \tag{2.2.12}$$

with the local operator:

$$\mathcal{O}_n^\Gamma(y,x) = \Phi'(y)\Gamma\big(xD_-\big)^n\Phi(y). \tag{2.2.13}$$

Obviously, (2.2.12) is obtained from (2.2.10) with $\kappa_+ = 0$ and $\kappa_- = \kappa$. In the following we often use the abbreviation $\mathcal{O}_n^\Gamma(0,x) \equiv \mathcal{O}_n^\Gamma(x)$. Some remarks are necessary with respect to these relations. The local operators in Eqs. (2.2.3), (2.2.8), and (2.2.12) have to be infinitely differentiable with respect to the expansion point $y \in \mathbb{R}^{2h}$ and are homogeneous polynomials of degree n in $x \in \mathbb{R}^{2h}$. Additionally, the operators (2.2.9) and (2.2.13) have the same anomalous dimensions.

The Fourier transforms of the 'centred' operator $\mathcal{O}^\Gamma(-\kappa x, \kappa x)$ and its nth moments $\mathcal{O}_n^\Gamma(x)$, i.e., its Taylor coefficients w.r.t. κ, are related as follows:

$$\mathcal{O}^\Gamma(-\kappa x, \kappa x) = \int \mathrm{d}^{2h}p\,\mathcal{O}^\Gamma(p)\,\mathrm{e}^{-\mathrm{i}\kappa xp} = \sum_{n=0}^{\infty} \frac{(-\mathrm{i}\kappa)^n}{n!}\,\mathcal{O}_n^\Gamma(x), \tag{2.2.14}$$

$$\mathcal{O}_n^\Gamma(x) = \int \mathrm{d}^{2h}p\,\mathcal{O}^\Gamma(p)\,(xp)^n = (\mathrm{i})^n \frac{\partial^n}{\partial\kappa^n}\mathcal{O}^\Gamma(-\kappa x, \kappa x)\Big|_{\kappa=0}. \tag{2.2.15}$$

For notational simplicity we wrote the Fourier measure without the usual factor $1/(2\pi)^{2h}$.
(2) *Decomposition* of local operators with respect to *irreducible tensor representations* of the Lorentz group $SO(1, 2h-1)$ or, equivalently, the orthogonal group $SO(2h)$.[10] These representations are built up by traceless tensors of rank m ($m = n +$ number of indices of Γ) whose symmetry class is determined by some (normalized) Young operators $\mathcal{Y}_{[m]} = (f_{[m]}/m!)\mathcal{Q}\mathcal{P}$, where $[m] = (m_1, m_2, \dots m_r)$ with $m_1 \geq m_2 \geq \dots \geq m_r$ and $\sum_{i=1}^r m_i = m$ denotes the corresponding Young pattern. $\mathcal{P}$ and $\mathcal{Q}$, as usual, denote symmetrization and antisymmetrization with respect to that pattern. The allowed Young patterns for $SO(2h)$, which because of the tracelessness are restricted by $\ell_1 + \ell_2 \leq 2h$ (ℓ: length of *columns* of $[m]$), are for a fixed value of m:

(i)  $j = m, m-2, m-4, \dots$

(ii)  $j = m-1, m-2, m-3, \dots$

(iii)  $j = m-2, m-3, m-4, \dots$

---

[10] The corresponding representations are related through analytic continuation, cf. [20, 121, 63]

(iv)  $j = m-2, m-3, m-4, \ldots$

$\vdots$

(Here, we depicted only those patterns which appear for $h = 2$ where $\ell_1 + \ell_2 \leq 4$.) Loosely speaking, the lower spins in the above series correspond to the various trace terms which define, in general, reducible higher twist operators. These higher twist operators again have to be decomposed according to the considered symmetry classes. For more details see Appendices A and B.

Now, two remarks are in order. At first, this decomposition of the local operators appearing in (2.2.3) into irreducible components is independent of the special values of κ_1, κ_2. Furthermore, the totally symmetric indices of irreducible local tensor operators are multiplied by $x^{\mu_1} \ldots x^{\mu_n}$ in order to obtain harmonic (tensorial) polynomials of order n. Obviously, in this turn we use the vector x as a device for writing tensors with special Young symmetry in analytic form (see Section 4.1).

A local operator without definite twist and spin is given as a finite sum of local operator with definite geometric twist,

$$\mathcal{O}_{\Gamma\, n}(y,x) = \bigoplus_{\tau=\tau_{\min}}^{\tau_{\max}} c^{(\tau)\Gamma' n'}_{\Gamma n}(x)\, \mathcal{O}^{(\tau)}_{\Gamma'\, n'}(y,x), \tag{2.2.16}$$

where $c^{(\tau)\Gamma' n'}_{\Gamma n}(x)$ are coefficient functions depending on x^2 and x_α etc.. The local operators of geometric twist can be written

$$\mathcal{O}^{(\tau)}_{\Gamma\, n}(y,x) = \mathcal{P}^{(\tau)\Gamma' n'}_{\Gamma n}\, \mathcal{O}_{\Gamma'\, n}(y,x), \tag{2.2.17}$$

$$\big(\mathcal{P}^{(\tau)} \times \mathcal{P}^{(\tau')}\big)_{\Gamma n}^{\ \ \Gamma' n'} = \delta^{\tau\tau'} \mathcal{P}^{(\tau)\Gamma' n'}_{\Gamma n}, \tag{2.2.18}$$

where $\mathcal{P}^{(\tau)}_{\Gamma n \Gamma' n'}$ are orthogonal projectors of definite geometric twist and definite spin, respectively. These projectors are given in terms of x^2, $\Box$, x_α, ∂_α etc.. It is obvious that the twist decomposition corresponds to a Lorentz spin decomposition of the local operator.

(3) *Resummation* of the local operators $\mathcal{O}^{(\tau)}_{\Gamma\, n}(y,x)$ belonging to the same symmetry class (for any n) and having equal twist τ. That infinite tower of local operators (2.2.17) creates a *nonlocal* operator of definite twist,

$$\mathcal{O}^{(\tau)}_{\Gamma}(\kappa_1 x, \kappa_2 x) = \bigoplus_{n=0}^{\infty} \frac{\kappa_-^n}{n!}\, \mathcal{O}^{(\tau)}_{\Gamma\, n}(y,x)|_{y=\kappa_+ x}. \tag{2.2.19}$$

Obviously, from a group theoretical point of view, this nonlocal operator is built up as direct sum of irreducible local tensor operators which have the same twist but different Lorentz spin. In fact, the nonlocal operator (2.2.19) has no definite Lorentz spin because it is an infinite tower of local operators with definite Lorentz spin. Therfore, the notion of geometric twist is a well-defined concept to classify nonlocal operators. The nonlocal operators $\mathcal{O}^{(\tau)}_{\Gamma}(\kappa_1 x, \kappa_2 x)$ are harmonic tensor functions and they are given by

$$\mathcal{O}^{(\tau)}_{\Gamma}(\kappa_1 x, \kappa_2 x) = \mathcal{P}^{(\tau)\Gamma'}_{\Gamma}\, \mathcal{O}_{\Gamma'}(\kappa_1 x, \kappa_2 x), \tag{2.2.20}$$

$$\big(\mathcal{P}^{(\tau)} \times \mathcal{P}^{(\tau')}\big)_{\Gamma}^{\ \ \Gamma'} = \delta^{\tau\tau'} \mathcal{P}^{(\tau)\Gamma'}_{\Gamma}, \tag{2.2.21}$$

where $\mathcal{P}^{(\tau)}_{\Gamma\Gamma'}$ are orthogonal projection operators of definite geometric twist for the nonlocal operators.

Now, putting together all contributions, i.e., including the higher twist contributions resulting from the trace terms, we get the (infinite) twist decomposition of the nonlocal operator we started with:

$$\mathcal{O}_{\Gamma}(\kappa_1 x, \kappa_2 x) = \bigoplus_{\tau=\tau_{\min}}^{\infty} c_{\Gamma}^{(\tau)\Gamma'}(x)\, \mathcal{O}^{(\tau)}_{\Gamma'}(\kappa_1 x, \kappa_2 x)\,, \tag{2.2.22}$$

where $c_{\Gamma}^{(\tau)\Gamma'}(x)$ are the coefficient functions of the nonlocal operators with geometric twist which are obtained from the coefficients $c_{\Gamma n}^{(\tau)\Gamma' n'}(x)$.

(4) Finally, the *projection onto the light–cone*, $x \to \tilde{x}$, leads to the required light–cone operator with well defined geometric twist. However, since the harmonic tensor polynomials essentially depend on (infinite) sums of powers of x^2 and $\Box$ as well as some specific differential operators in front of it, in that limit only a finite number of terms survive:

$$\mathcal{O}_{\Gamma}(\kappa_1 \tilde{x}, \kappa_2 \tilde{x}) = \bigoplus_{\tau=\tau_{\min}}^{\tau_{\max}} c_{\Gamma}^{(\tau)\Gamma'}(\tilde{x})\, \mathcal{O}^{(\tau)}_{\Gamma'}(\kappa_1 \tilde{x}, \kappa_2 \tilde{x})\,. \tag{2.2.23}$$

The resulting light–ray operators of definite twist are tensor functions on the light–cone which fulfill another kind of tracelessness conditions to be formulated with the help of the interior (on the cone) derivative (see Appendix C). Let us remark that step (3) and (4) can be interchanged without changing the result.

2.2.2 Transformation properties of (non)local operators with definite geometric twist

Because the notion of geometric twist is related to (j, d) of the group $SO(1, 2h-1) \times \mathbb{R}_+$, where $\mathbb{R}_+$ is the group of dilations, it is useful to give the transformation rules of the (unrenormalized) nonlocal operators with respect to this group. We induce the transformation rules of the nonlocal operators from the transformation properties of local fields which are given in [47, 48]. Let $\mathcal{O}^{(\tau)}_{\Gamma n}(y, x)$ be the local operator of geometric twist τ.

The transformation properties of these local operators are [47, 48]

a) Poincaré transformations:

$$\hat{U}(a,\Lambda)\mathcal{O}^{(\tau)}_{\Gamma n}(y,x)\hat{U}^{-1}(a,\Lambda) = V_{\Gamma\Gamma'}(\Lambda)\mathcal{O}^{(\tau)}_{\Gamma' n}(\Lambda^{-1}(y-a), \Lambda^{-1}x), \quad a \in \mathbb{R}^{2h}, \Lambda \in SO(1, 2h-1), \tag{2.2.24}$$

where $V(\Lambda)$ is a finite dimensional representation of the Lorentz group.

b) Dilations:

$$\hat{U}(\rho)\mathcal{O}^{(\tau)}_{\Gamma n}(y,x)\hat{U}^{-1}(\rho) = \rho^d \mathcal{O}^{(\tau)}_{\Gamma n}(\rho y, x), \quad \rho > 0, \tag{2.2.25}$$

where d is the (scale) dimension of the local operator (in mass units). From group theoretical point of view, the anomalous dimensions do not destroy the invariance under dilations, they just change the representation with respect to the dilation group [45].

The transformation properties for the nonlocal operators are

a) Poincaré transformations:

$$\hat{U}(a,\Lambda)\mathcal{O}^{(\tau)}_{\Gamma}(y-x, y+x)\hat{U}^{-1}(a,\Lambda) = V_{\Gamma\Gamma'}(\Lambda)\mathcal{O}^{(\tau)}_{\Gamma'}(\Lambda^{-1}(y-a-x), \Lambda^{-1}(y-a+x)), \tag{2.2.26}$$

b) Dilations:

$$\hat{U}(\rho)\mathcal{O}_\Gamma^{(\tau)}(y-x,y+x)\hat{U}^{-1}(\rho)=\rho^d\mathcal{O}_\Gamma^{(\tau)}(\rho(y-x),\rho(y+x)). \tag{2.2.27}$$

Let us point out that in the case of nonlocal operators, i.e., for an infinite series of local ones, we deal with an infinite-dimensional representation of the dilation and the Lorentz group. Obviously, the translation $\hat{U}(a,0)=\mathrm{e}^{-\mathrm{i}a\cdot\hat{P}}$ with $\hat{P}_\mu=\mathrm{i}\frac{\partial}{\partial y^\mu}$ just shifts the expansion point y by the vector a. If we had chosen $a=x$, the twist of these operators would be not invariant under this translation, because the differential operators for definite twist do not commute with x. [11] For the non-forward matrix element of an uncentred operator with definite twist we obtain the relation to the corresponding centred operator as follows

$$\begin{aligned}\langle P_2|\mathcal{O}_\Gamma^{(\tau)}(\kappa_1 x,\kappa_2 x)|P_1\rangle &\equiv \langle P_2|\mathcal{O}_\Gamma^{(\tau)}(\kappa_+x-\kappa_-x,\kappa_+x+\kappa_-x)|P_1\rangle\\ &=\mathcal{P}_\Gamma^{(\tau)\Gamma'}\left\{\mathrm{e}^{\mathrm{i}\kappa_+(xP_-)}\langle P_2|\mathcal{O}_{\Gamma'}(-\kappa_-x,\kappa_-x)|P_1\rangle\right\}. \end{aligned}\tag{2.2.28}$$

We see that the twist projector acts on the factor $\exp(\mathrm{i}\kappa_+(xP_-))$ in Eq. (2.2.28). In the forward case ($P_1=P_2$) this factor gives no contribution due to $P_-=0$.

Let me mention that it is possible to construct conformal operators with geometric twist which carry elementary representations of the conformal group $SO(2,2h)$. The condition of conformal invariance is given by the infinitesimal transformation law under special conformal transformations at $y=0$:

$$[\hat{K}_\mu,\mathcal{O}_{\Gamma n}^{(\tau)}(y,x)]|_{y=0}=0, \tag{2.2.29}$$

where $\hat{K}_\mu$ is the generator for special conformal transformations [45, 85, 54]. The conformal LC-operators with minimal twist $\tau=2h-2$ are [45]:

$$\mathcal{O}_n^\phi(y,\tilde{x})=(\tilde{x}\partial_+)^n\phi(y)C_n^{h-3/2}(\tilde{x}D_-/\tilde{x}\partial_+)\phi(y), \tag{2.2.30}$$

$$\mathcal{O}_n^\psi(y,\tilde{x})=(\tilde{x}\partial_+)^{n-1}\bar{\psi}(y)(\tilde{x}\gamma)C_{n-1}^{h-1/2}(\tilde{x}D_-/\tilde{x}\partial_+)\psi(y), \tag{2.2.31}$$

$$\mathcal{O}_n^F(y,\tilde{x})=(\tilde{x}\partial_+)^{n-2}\tilde{x}^\mu F_\mu^{a\rho}(y)C_{n-2}^{h+1/2}(\tilde{x}D_-/\tilde{x}\partial_+)F_{\rho\nu}^a(y)\tilde{x}^\nu. \tag{2.2.32}$$

---

[11] The total derivative ∂_+ is related to translations in the following way: $[\hat{P},\mathcal{O}_{\Gamma n}^{(\tau)}(y,x)]=\mathrm{i}\partial_+\mathcal{O}_{\Gamma n}^{(\tau)}(y,x)$.

Chapter 3

Application of the twist decomposition to different aspects of the hadron phenomenology

3.1 Quark distribution functions of definite geometric twist

The aim of this Section is to present new *forward quark distribution functions* which are related to the nonlocal LC-operators of different geometric twist. In that framework, it is possible to investigate in a unique manner the contributions resulting from the traces of the operators having well-defined twist. In addition, the classification of the various distribution functions appears to be quite straightforward.

The definitions of geometric and dynamical twist do not coincide at higher orders. Thus, the definition of dynamical twist has a mismatch with the conventional definition of geometrical twist. This mismatch gives rise to relations of Wandzura-Wilczek type which show that dynamical twist distributions contain various parts of different geometric twist.

3.1.1 Quark distribution functions of definite dynamical twist

Quark distribution functions are usually defined as the forward matrix elements of bilocal quark-antiquark operators. Jaffe and Ji [78, 79] observed that one can generate all dynamical quark distribution functions up to (dynamical) twist-4 by decomposing the bilocal operators into independent tensor structures of their matrix elements. To implement the light-cone expansion in a systematic way, it is convenient to use light-like vectors. Using the relations (2.1.4), one may introduce light-like vectors p_α and $\tilde{x}_\alpha$ with

$$p^2 = 0, \qquad \tilde{x}^2 = 0, \tag{3.1.1}$$

according to (see Eq. (2.1.17) with $\rho_\alpha = P_\alpha/M$ and $\rho^2 = 1$)

$$\tilde{x}_\alpha = x_\alpha - \frac{P_\alpha}{M^2}\Big((xP) - \sqrt{(xP)^2 - x^2M^2}\Big), \qquad p_\alpha = P_\alpha - \frac{1}{2}\tilde{x}_\alpha \frac{M^2}{\tilde{x}P}. \tag{3.1.2}$$

The spin vector of the nucleon can be decomposed as

$$S_\alpha = S^\perp_\alpha + p_\alpha \frac{\tilde{x}S}{\tilde{x}P} - \frac{1}{2}\tilde{x}_\alpha \frac{\tilde{x}S}{(\tilde{x}P)^2} M^2, \tag{3.1.3}$$

where $S_\alpha^\perp$ is the transversal spin-polarization ($\tilde{x}S^\perp = 0$). Some useful scalar products are

$$\tilde{x}P = \tilde{x}p = \sqrt{(xP)^2 - x^2M^2}, \qquad pS = -\frac{M^2}{2(\tilde{x}p)}\,\tilde{x}S\,. \tag{3.1.4}$$

Accordingly, one finds nine independent quark distribution functions associating with each tensor structure. Using the Lorentz-tensor decomposition of the matrix elements of the gauge-invariant bilocal quark-antiquark operators (2.1.31), (2.1.30), (2.1.40), and (2.1.39), the definitions of the chiral-even quark distributions are:[1]

$$\begin{aligned}\langle PS|O_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle = 2\bigg[p_\alpha\frac{\tilde{x}S}{\tilde{x}P}\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}g_1(z,\mu^2) + S_\alpha^\perp\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}g_T(z,\mu^2)\\ + \tilde{x}_\alpha\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}g_3(z,\mu^2)\bigg]\end{aligned} \tag{3.1.5}$$

and

$$\langle P|O_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})|P\rangle = 2\left[p_\alpha\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}f_1(z,\mu^2) + \tilde{x}_\alpha\frac{M^2}{\tilde{x}P}\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}f_4(z,\mu^2)\right], \tag{3.1.6}$$

while the chiral-odd quark distributions are defined as

$$\langle P|N(\kappa_1\tilde{x},\kappa_2\tilde{x})|P\rangle = 2M\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}e(z,\mu^2), \tag{3.1.7}$$

and

$$\begin{aligned}\langle PS|M_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle = \frac{4}{M}\bigg[S^\perp_{[\alpha}p_{\beta]}\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}h_1(z,\mu^2) \qquad (3.1.8)\\ - \tilde{x}_{[\alpha}p_{\beta]}\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}h_L(z,\mu^2) - \tilde{x}_{[\alpha}S^\perp_{\beta]}\frac{M^2}{\tilde{x}P}\int_0^1 \mathrm{d}z\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}h_3(z,\mu^2)\bigg].\end{aligned}$$

These distribution functions are dimensionless functions of the Bjorken variable which we here denote by z. Additionally, they depend on the renormalization point μ^2 as well because the bilocal light-cone operators on the l.h.s. are renormalized at μ^2. We defined $\kappa = \kappa_1 - \kappa_2$.

In Tab. 3.1 the classification of these dynamical quark distribution functions according to their spin, dynamical twist and chirality and the relation to the geometric twist (see Section 3.1.4) are given. The parton distributions in the first row are spin-independent, those in the second and third row describe longitudinally ($S_\parallel$) and transversely ($S_\perp$) polarized nucleons, respectively. The underlined distributions are referred to as chiral-odd, because they correspond to chirality-violating Dirac matrix structures $\Gamma = \{\mathrm{i}\sigma_{\alpha\beta}\gamma_5,\, 1\}$. The other distributions are termed chiral-even, because of the chirality-conserving structures $\Gamma = \{\gamma_\alpha,\, \gamma_\alpha\gamma_5\}$.

The nucleon (spin-1/2 hadron) has thus three independent dynamical twist-2 quark distributions f_1, g_1, and h_1. These distributions have a simple parton model interpretation and appear as a leading contribution to various hard inclusive processes. For example, g_1 measures the quark helicity distribution in a longitudinally polarized nucleon and is called the *helicity* distribution. The both distributions f_1 and g_1 can be measured through deep inelastic lepton-nucleon scattering (DIS). The function h_1 appears as a leading contribution to the transversely polarised

---

[1] Here we suppress the flavour index.

Twist t	2	3	4
	$O(1)$	$O(1/Q)$	$O(1/Q^2)$
spin ave.	$f_1 = F^{(2)}$	$\underline{e} = E^{(3)}$	$f_4 = F^{(4)}/2 + F_f(F^{(2)}, F^{(4)})/2$
$S_\parallel$	$g_1 = G^{(2)}$	$\underline{h_L} = H^{(3)} + F_L(H^{(2)}, H^{(3)})$	$g_3 = -G^{(4)}/2 + F_g(G^{(2)}, G^{(3)}, G^{(4)})/2$
$S_\perp$	$\underline{h_1} = H^{(2)}$	$g_T = G^{(3)} + F_T(G^{(2)}, G^{(3)})$	$\underline{h_3} = H^{(4)}/2 + F_h(H^{(2)}, H^{(3)}, H^{(4)})/2$

Table 3.1: Spin, dynamical twist and chiral classification of the nucleon structure functions.

nucleon-nucleon Drell-Yan process. This distribution measures the probability to find a quark with spin polarized along the transverse spin of a nucleon minus the probability to find it polarized oppositely and was called the *transversity* distribution in Refs. [78, 79]. Additionally, the nucleon has three dynamical twist-3 distributions g_T, h_L, and e. Physically higher-twist distributions represent complicated quark-gluon correlations in the hadrons and cannot be interpreted in a simple parton model. Therefore, higher-twist or multiparton distributions are useful to probe quark-gluon dynamics as well as nonperturbative aspect of QCD. The distributions g_T and h_L are contributions to proper asymmetries in polarized DIS and Drell-Yan processes, respectively. The direct measurement of e seems very difficult [79]. The dynamical twist-4 distributions represent rather complicated two-quark-one-gluon, two-quark-two-gluon, and four-quark correlations so that they are very difficult to handle with. However, in principle, f_4 and h_3 are observable in unpolarized DIS and polarized Drell-Yan processes, respectively [79]. Up to now, no measurement of g_3 is known in the literature.

3.1.2 Twist decomposition of bilinear quark operators

In order to be able to classify the quark distributions with respect to geometrical twist, we have to use the corresponding operators with geometric twist. Now we apply the bilocal quark operators on the light-cone and the derived scalar and vector operators, $O_{(5)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = \tilde{x}^\alpha O_{(5)\alpha}(\kappa_1\tilde{x}, \kappa_2\tilde{x})$ and $M_{(5)\alpha}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = \tilde{x}^\beta M_{(5)[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x})$, respectively.

The resulting decomposition for the vector and skew tensor operators – which is independent of the presence or absence of γ_5 – reads:[2]

$$O(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = \tilde{x}^\alpha O^{\mathrm{tw2}}_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \tag{3.1.9}$$

$$O_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = O^{\mathrm{tw2}}_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + O^{\mathrm{tw3}}_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + O^{\mathrm{tw4}}_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \tag{3.1.10}$$

$$M_{[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = M^{\mathrm{tw2}}_{[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + M^{\mathrm{tw3}}_{[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + M^{\mathrm{tw4}}_{[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \tag{3.1.11}$$

$$M_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = M^{\mathrm{tw2}}_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + M^{\mathrm{tw3}}_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \tag{3.1.12}$$

$$N(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = N^{\mathrm{tw3}}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) \tag{3.1.13}$$

---

[2] The explicite twist decomposition of these light-cone operators will be computed in Section 4.2 (see also Ref. [63]). The twist decomposition of off-cone operators will be discussed in Sections 4.1 and 4.4.

with

$$O^{\mathrm{tw2}}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\Big(\partial_\alpha + \frac{1}{2}(\ln\lambda)\,x_\alpha\Box\Big)x^\mu O_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \tag{3.1.14}$$

$$O^{\mathrm{tw3}}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\Big(\delta^\mu_\alpha(x\partial) - x^\mu\partial_\alpha - (1+2\ln\lambda)x_\alpha\partial^\mu - (\ln\lambda)\,x_\alpha x^\mu\Box\Big)O_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \tag{3.1.15}$$

$$O^{\mathrm{tw4}}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \tilde{x}_\alpha\int_0^1 \mathrm{d}\lambda\Big((1+\ln\lambda)\partial^\mu + \frac{1}{2}(\ln\lambda)\,x^\mu\Box\Big)O_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \tag{3.1.16}$$

$$M^{\mathrm{tw2}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\Big\{2\lambda\,\partial_{[\beta}\delta^\mu_{\alpha]} - (1-\lambda)\big(2x_{[\alpha}\partial_{\beta]}\partial^\mu - x_{[\alpha}\delta^\mu_{\beta]}\Box\big)\Big\}x^\nu M_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \tag{3.1.17}$$

$$\begin{aligned} M^{\mathrm{tw3}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\Big\{&\lambda\big((x\partial)\delta^\nu_{[\beta} - 2x^\nu\partial_{[\beta}\big)\delta^\mu_{\alpha]} + \tfrac{1-\lambda^2}{\lambda}\Big(x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]} \\ &- x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box - x_{[\alpha}\partial_{\beta]}x^{[\mu}\partial^{\nu]}\Big)\Big\}M_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \end{aligned} \tag{3.1.18}$$

$$M^{\mathrm{tw4}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\tfrac{1-\lambda}{\lambda}\Big\{x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box - 2x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]}\Big\}M_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \tag{3.1.19}$$

$$M^{\mathrm{tw2}}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = M_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x}) - \tilde{x}_\alpha\partial^\mu\int_0^1 \mathrm{d}\lambda\,\lambda\,M_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \tag{3.1.20}$$

$$M^{\mathrm{tw3}}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \tilde{x}_\alpha\partial^\mu\int_0^1 \mathrm{d}\lambda\,\lambda M_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \tag{3.1.21}$$

Let us remark that the (axial) vector and skew tensor operators of twist τ, $O^{(\tau)}_{(5)\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ and $M^{(\tau)}_{(5)[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$, are obtained from the (undecomposed) operators $O_{(5)\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ and $M_{(5)[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ by the application of the twist projectors on the light-cone $\mathcal{P}^{(\tau)\mu}_{\alpha}$ and $\mathcal{P}^{(\tau)[\mu\nu]}_{[\alpha\beta]}$ (see also Eqs. (2.2.20) and (2.2.21)) defined by Eqs. (3.1.14) – (3.1.16) and (3.1.17) – (3.1.19), respectively:

$$O^{(\tau)}_{(5)\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = (\mathcal{P}^{(\tau)\mu}_{\alpha}O_{(5)\mu})(\kappa_1\tilde{x},\kappa_2\tilde{x}) \tag{3.1.22}$$

$$M^{(\tau)}_{(5)[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = (\mathcal{P}^{(\tau)[\mu\nu]}_{[\alpha\beta]}M_{(5)[\mu\nu]})(\kappa_1\tilde{x},\kappa_2\tilde{x}) \tag{3.1.23}$$

$$M^{(\tau)}_{(5)\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = (\mathcal{P}^{(\tau)\mu}_{(v)\alpha}M_{(5)\mu})(\kappa_1\tilde{x},\kappa_2\tilde{x}). \tag{3.1.24}$$

The twist–2 vector and skew tensor operators, Eqs. (3.1.14) and (3.1.17), are related to the corresponding scalar and vector operators, (3.1.9) and (3.1.20), respectively, in group theoretical way.

Here, we give the corresponding local expressions of the nonlocal operators, Eqs. (3.1.9) – (3.1.21). The relation between the local and nonlocal operators are

$$O_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \sum_{n=0}^{\infty}\frac{\kappa_-^n}{n!}\,O_{\alpha n}(y,\tilde{x})|_{y=\kappa_+\tilde{x}}, \qquad M_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \sum_{n=0}^{\infty}\frac{\kappa_-^n}{n!}\,M_{[\alpha\beta]n}(y,\tilde{x})|_{y=\kappa_+\tilde{x}}.$$

The expressions for the local operators are:

$$
\begin{aligned}
O^{\rm tw2}_{n+1}(y,\tilde x) &\equiv \tilde x^\mu O_{\mu n}(y,\tilde x) = \bar\psi(y)\gamma^5(\gamma\tilde x)(\tilde x\overset{\leftrightarrow}{D})^n\psi(y) && (3.1.25)\\
O^{\rm tw2}_{\alpha n}(y,\tilde x) &= \frac{1}{n+1}\Big(\partial_\alpha - \frac{1}{2(n+1)}\,x_\alpha\Box\Big)O_{n+1}(y,x)\big|_{x=\tilde x} && (3.1.26)\\
O^{\rm tw3}_{\alpha n}(y,\tilde x) &= \frac{1}{n+1}\Big(n\delta^\mu_\alpha - x^\mu\partial_\alpha - \frac{1}{n+1}x_\alpha\big((n-1)\partial^\mu - x^\mu\Box\big)\Big)O_{\mu n}(y,x)\big|_{x=\tilde x} && (3.1.27)\\
O^{\rm tw4}_{\alpha n}(y,\tilde x) &= \frac{1}{(n+1)^2}\,\tilde x_\alpha\Big(n\partial^\mu - \tfrac{1}{2}x^\mu\Box\Big)O_{\mu n}(y,x)\big|_{x=\tilde x} && (3.1.28)\\
M^{\rm tw2}_{[\alpha\beta]n}(y,\tilde x) &= \frac{1}{n+2}\Big\{2\partial_{[\beta}\delta^\mu_{\alpha]} - \frac{1}{n+1}\big(2x_{[\alpha}\partial_{\beta]}\partial^\mu - x_{[\alpha}\delta^\mu_{\beta]}\Box\big)\Big\}M_{\mu n+1}(y,x)\big|_{x=\tilde x} && (3.1.29)\\
M^{\rm tw3}_{[\alpha\beta]n}(y,\tilde x) &= \frac{1}{n+2}\Big\{\big(n\delta^\nu_{[\beta} - 2x^\nu\partial_{[\beta}\big)\delta^\mu_{\alpha]} + \frac{2}{n}\Big(x_{[\alpha}\big((n-1)\delta^{[\mu}_{\beta]} - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]} - x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box\\
&\qquad - x_{[\alpha}\partial_{\beta]}x^{[\mu}\partial^{\nu]}\Big)\Big\}M_{[\mu\nu]n}(y,x)\big|_{x=\tilde x} && (3.1.30)\\
M^{\rm tw4}_{[\alpha\beta]n}(y,\tilde x) &= \frac{1}{(n+1)n}\Big\{x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box - 2x_{[\alpha}\big((n-1)\delta^{[\mu}_{\beta]} - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]}\Big\}M_{[\mu\nu]n}(y,x)\big|_{x=\tilde x} && (3.1.31)\\
M^{\rm tw2}_{\alpha n+1}(y,\tilde x) &= M_{\alpha n+1}(\tilde x) - \frac{1}{n+2}\,\tilde x_\alpha\partial^\mu M_{\mu n+1}(y,x)\big|_{x=\tilde x} && (3.1.32)\\
M^{\rm tw3}_{\alpha n+1}(y,\tilde x) &= \frac{1}{n+2}\,\tilde x_\alpha\partial^\mu M_{\mu n+1}(y,x)\big|_{x=\tilde x} && (3.1.33)\\
N^{\rm tw3}_{n}(y,\tilde x) &= \bar\psi(y)(\tilde x\overset{\leftrightarrow}{D})^n\psi(y). && (3.1.34)
\end{aligned}
$$

The twist projectors of the nonlocal as well as local LC operators can be formulated in terms of *inner derivatives* on the light–cone (see Section 4.1 and Ref. [66]).

3.1.3 Forward matrix elements of LC–operators with definite geometric twist

Now, we define the (polarized) quark distribution functions for the bilinear quark operators with definite twist. As usual, the matrix elements of the nucleon targets, $\bar U(P,S)\gamma^\mu U(P,S) = 2P^\mu$ and $\bar U(P,S)\gamma^5\gamma^\mu U(P,S) = 2S^\mu$, are related to the nucleon momentum P_μ and nucleon spin vector S_μ, respectively. Here $U(P,S)$ denotes the free hadronic spinor.

Let us first consider the chiral-even *pseudo scalar operator* in the polarized case. The (forward) matrix element of this twist-2 operator, Eq. (3.1.9), taken between hadron states $|PS\rangle$ is trivially represented as

$$
\langle PS|O^{\rm tw2}_5(\kappa_1\tilde x,\kappa_2\tilde x)|PS\rangle = 2(\tilde xS)\int_0^1 {\rm d}z\,G^{(2)}(z,\mu^2)\,{\rm e}^{{\rm i}\kappa z(\tilde xP)} = 2(\tilde xS)\sum_{n=0}^\infty \frac{({\rm i}\kappa(\tilde xP))^n}{n!}\,G^{(2)}_n(\mu^2). \tag{3.1.35}
$$

Here, $G^{(2)}(z,\mu^2)$ is the twist-2 parton distribution function which by a Mellin transformation is obtained from the corresponding moments

$$
G^{(2)}_n(\mu^2) = \int_0^1 {\rm d}z\,z^n G^{(2)}(z,\mu^2).
$$

Now we consider the chiral-even *axial vector operator*. Using the projection properties (3.1.22), we introduce the parton distribution functions $G^{(\tau)}(z,\mu^2)$ of twist τ by

$$
\begin{aligned}
\langle PS|O^{(\tau)}_{5\alpha}(\kappa_1\tilde x,\kappa_2\tilde x)|PS\rangle &\equiv \langle PS|(\mathcal P^{(\tau)\beta}_\alpha O^{(\tau)}_{5\beta})(\kappa_1\tilde x,\kappa_2\tilde x)|PS\rangle\\
&= \mathcal P^{(\tau)\beta}_\alpha\Big(2S_\beta\int_0^1 {\rm d}z\,G^{(\tau)}(z,\mu^2)\,{\rm e}^{{\rm i}\kappa z(\tilde xP)}\Big). \qquad (3.1.36)
\end{aligned}
$$

For $\tau = 2$ this is consistent with (3.1.35), it reads

$$\langle PS|O^{\mathrm{tw}2}_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle = 2\int_0^1 \mathrm{d}\lambda \Big[\partial_\alpha + \frac{1}{2}\ln(\lambda)x_\alpha\Box\Big](xS)\int_0^1 \mathrm{d}z\, G^{(2)}(z,\mu^2)\,\mathrm{e}^{\mathrm{i}\kappa\lambda z(xP)}\big|_{x=\tilde{x}}\ .$$

Using the projection operators as they are determined by Eqs. (3.1.14) – (3.1.16) we obtain (from now on we suppress μ^2)

$$\begin{aligned}
\langle PS|O^{\mathrm{tw}2}_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= 2\int_0^1 \mathrm{d}\lambda\int_0^1 \mathrm{d}z\, G^{(2)}(z)\Big[S_\alpha + \mathrm{i}\kappa\lambda z P_\alpha(\tilde{x}S),\\
&\qquad\qquad + \frac{\tilde{x}_\alpha}{2}(\tilde{x}S)M^2(\mathrm{i}\kappa\lambda z)^2\big(\ln\lambda\big)\Big]\mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)}\\
&= 2\sum_{n=0}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}G^{(2)}_n\Big\{\frac{1}{n+1}\Big(S_\alpha + nP_\alpha\frac{\tilde{x}S}{\tilde{x}P}\Big) - \frac{n(n-1)}{2(n+1)^2}\tilde{x}_\alpha\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2\Big\},
\end{aligned} \tag{3.1.37}$$

$$\begin{aligned}
\langle PS|O^{\mathrm{tw}3}_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= 2\int_0^1 \mathrm{d}\lambda\int_0^1 \mathrm{d}z\, G^{(3)}(z)\Big[\Big(S_\alpha(\tilde{x}P) - P_\alpha(\tilde{x}S)\Big)\mathrm{i}\kappa\lambda z\\
&\qquad\qquad - \tilde{x}_\alpha M^2(\tilde{x}S)(\mathrm{i}\kappa\lambda z)^2\ln\lambda\Big]\mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)},\\
&= 2\sum_{n=1}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}G^{(3)}_n\Big\{\frac{n}{n+1}\Big(S_\alpha - P_\alpha\frac{\tilde{x}S}{\tilde{x}P}\Big) + \frac{n(n-1)}{(n+1)^2}\tilde{x}_\alpha\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2\Big\},
\end{aligned} \tag{3.1.38}$$

$$\begin{aligned}
\langle PS|O^{\mathrm{tw}4}_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= \int_0^1 \mathrm{d}\lambda\int_0^1 \mathrm{d}z\, G^{(4)}(z)\Big[\tilde{x}_\alpha(\tilde{x}S)M^2(\mathrm{i}\kappa\lambda z)^2\ln\lambda\Big]\mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)},\\
&= -\sum_{n=2}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}G^{(4)}_n\frac{n(n-1)}{(n+1)^2}\tilde{x}_\alpha\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2.
\end{aligned} \tag{3.1.39}$$

In the first expression of any equation we have written the nonlocal LC-operators of definite twist, and in the second line, after expanding the exponential, we introduced the moments of the structure functions. The *local* twist-2 and twist-3 matrix elements of traceless operators are given off–cone in Refs. [52, 105] for $n \leq 3$ and in Refs. [110, 26] for any n. Obviously, the trace terms which have been explicitly subtracted are proportional to M^2. According to the terminology of Jaffe and Ji, they contribute to *dynamical* twist-4. For the twist–2 operator we observe that the terms proportional to S_α, P_α and $\tilde{x}_\alpha$ have contributions starting with the zeroth, first and second moment, respectively. The twist–3 operator starts with the first moment, and the twist–4 operator starts with the second moment. Analogous statements also hold for the twist–τ operators below.

Putting together the different twist contributions we obtain, after replacing $\mathrm{i}\kappa\lambda z(\tilde{x}P)$ by $\lambda\partial/\partial\lambda$ and performing partial integrations, the following forward matrix element of the original operator ($\zeta = \kappa(\tilde{x}P)$)

$$\begin{aligned}
\langle PS|O_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= 2P_\alpha\frac{\tilde{x}S}{\tilde{x}P}\int_0^1 \mathrm{d}z\Big(G^{(2)}(z) - G^{(3)}(z)\Big)[e_0(\mathrm{i}\zeta z) - e_1(\mathrm{i}\zeta z)]\\
&\quad + 2S_\alpha\int_0^1 \mathrm{d}z\Big(G^{(2)}(z)e_1(\mathrm{i}\zeta z) + G^{(3)}(z)[e_0(\mathrm{i}\zeta z) - e_1(\mathrm{i}\zeta z)]\Big)\\
&\quad - \tilde{x}_\alpha\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2\int_0^1 \mathrm{d}z\Big(G^{(2)}(z) - 2G^{(3)}(z) + G^{(4)}(z)\Big)\\
&\qquad \times\Big[e_0(\mathrm{i}\zeta z) - 3e_1(\mathrm{i}\zeta z) + 2\int_0^1 \mathrm{d}\lambda e_1(\mathrm{i}\zeta\lambda z)\Big],
\end{aligned} \tag{3.1.40}$$

where we introduced the following "truncated exponentials"

$$e_0(\mathrm{i}\zeta z) = \mathrm{e}^{\mathrm{i}\zeta z}, \quad e_1(\mathrm{i}\zeta z) = \int_0^1 \mathrm{d}\lambda\, \mathrm{e}^{\mathrm{i}\zeta z\lambda} = \frac{\mathrm{e}^{\mathrm{i}\zeta z} - 1}{\mathrm{i}\zeta z}, \quad \cdots \quad , e_{n+1}(\mathrm{i}\zeta z) = \frac{(-1)^n}{n!} \int_0^1 \mathrm{d}\lambda\, \lambda^n\, \mathrm{e}^{\mathrm{i}\zeta z\lambda}. \tag{3.1.41}$$

As it should be the application of the projection operators $\mathcal{P}_\alpha^{(\tau)\beta}$ onto (3.1.40) reproduces the matrix elements (3.1.37) – (3.1.39). In comparison with the dynamical distribution functions we observe that the geometric distribution functions are accompanied not simply by the exponentials, $e_0(\mathrm{i}\zeta z)$, but by more involved combinations whose series expansion directly leads to the representations with the help of moments.

Now we consider the chiral-even *vector operator* $O_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x})$, which obeys relations Eqs. (3.1.9) – (3.1.16), as well as (3.1.22) with the *same* projection operator as the axial vector operator. Let us introduce the corresponding parton distribution functions $F^{(\tau)}(z)$ of twist τ by

$$\langle PS|O_\alpha^{(\tau)}(\kappa_1\tilde{x}, \kappa_2\tilde{x})|PS\rangle = \mathcal{P}_\alpha^{(\tau)\mu}\Big(2P_\mu \int_0^1 \mathrm{d}z\, F^{(\tau)}(z)\, \mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}\Big). \tag{3.1.42}$$

Again, this is consistent with the definition of the unpolarized distribution function $F^{(2)}(z)$ by the matrix element of the corresponding twist-2 *scalar operator*

$$\langle P|O^{\mathrm{tw}2}(\kappa_1\tilde{x}, \kappa_2\tilde{x})|P\rangle = 2(\tilde{x}P) \int_0^1 \mathrm{d}z\, F^{(2)}(z)\, \mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)} = 2(\tilde{x}P) \sum_{n=0}^{\infty} \frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!} F_n^{(2)}. \tag{3.1.43}$$

The forward matrix elements of the *vector operators* of twist τ are obtained as follows:

$$\begin{aligned}
\langle P|O_\alpha^{\mathrm{tw}2}(\kappa_1\tilde{x}, \kappa_2\tilde{x})|P\rangle &= 2\int_0^1 \mathrm{d}\lambda \int_0^1 \mathrm{d}z\, F^{(2)}(z) \Big[P_\alpha\big(1 + \mathrm{i}\kappa\lambda z(\tilde{x}P)\big) \\
&\qquad + \tilde{x}_\alpha M^2 \mathrm{i}\kappa\lambda z \ln\lambda \Big(1 + \frac{1}{2}\mathrm{i}\kappa\lambda z(\tilde{x}P)\Big)\Big] \mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)} \\
&= 2\sum_{n=0}^{\infty} \frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!} F_n^{(2)} \Big\{P_\alpha - \frac{n}{2(n+1)} \tilde{x}_\alpha \frac{M^2}{\tilde{x}P}\Big\}, \qquad (3.1.44) \\
\langle P|O_\alpha^{\mathrm{tw}3}(\kappa_1\tilde{x}, \kappa_2\tilde{x})|P\rangle &= -2\tilde{x}_\alpha M^2 \int_0^1 \mathrm{d}\lambda \int_0^1 \mathrm{d}z\, F^{(3)}(z) \mathrm{i}\kappa\lambda z \,\Big[\Big(1 + 2\ln\lambda\Big) \\
&\qquad + \mathrm{i}\kappa\lambda z\big(\ln\lambda\big)\,(\tilde{x}P)\Big] \mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)} \equiv 0\,, \qquad (3.1.45) \\
\langle P|O_\alpha^{\mathrm{tw}4}(\kappa_1\tilde{x}, \kappa_2\tilde{x})|P\rangle &= -2\tilde{x}_\alpha M^2 \int_0^1 \mathrm{d}\lambda \int_0^1 \mathrm{d}z\, F^{(4)}(z) \mathrm{i}\kappa\lambda z\, \big(\ln\lambda\big)\Big(1 + \frac{1}{2}\mathrm{i}\kappa\lambda z(\tilde{x}P)\Big) \mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)} \\
&= \sum_{n=1}^{\infty} \frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!} F_n^{(4)} \frac{n}{(n+1)} \tilde{x}_\alpha \frac{M^2}{\tilde{x}P}. \qquad (3.1.46)
\end{aligned}$$

The off–cone traceless local twist-2 matrix element has already been obtained in [71, 58]. The vanishing of the twist-3 distribution function $F^{(3)}(z)$ which follows here by partial integration proves in the *nonlocal* case have already proven for the *local* twist-3 operator by Jaffe and Soldate [75]. The trace terms of the twist-2 operator and the twist-4 operator itself both contribute to dynamical twist-4 and, consequently, to the same $1/Q^2$-behaviour in the cross section.

Putting together all the contributions of different twist and performing partial λ–integrations, we obtain the following representation for the forward matrix element of the original operator

$$\begin{aligned}\langle P|O_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})|P\rangle = 2P_\alpha\int_0^1 \mathrm{d}z\, F^{(2)}(z) e_0(\mathrm{i}\zeta z) - \tilde{x}_\alpha\frac{M^2}{\tilde{x}P}\int_0^1 \mathrm{d}z\Big(F^{(2)}(z)-F^{(4)}(z)\Big)&\\ \times\big[e_0(\mathrm{i}\zeta z)-e_1(\mathrm{i}\zeta z)\big].&\end{aligned}\tag{3.1.47}$$

Let us now consider the simplest bilocal *scalar operator*, $N(\kappa_1\tilde{x},\kappa_2\tilde{x})\equiv N^{\mathrm{tw3}}(\kappa_1\tilde{x},\kappa_2\tilde{x})$, which is trivially a twist-3 operator on the light-cone. Its matrix element arises as

$$\langle P|N(\kappa_1\tilde{x},\kappa_2\tilde{x})|P\rangle = 2M\int_0^1 \mathrm{d}z\, E^{(3)}(z)\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)} = 2M\sum_{n=0}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}E_n^{(3)},\tag{3.1.48}$$

where $E^{(3)}(z)$ is another spin-independent twist-3 structure function.

Now, we calculate the matrix elements of the chiral-odd *vector operators*. In fact they may be obtained from the chiral-odd skew-tensor operator by contraction with $\tilde{x}_\beta$ (see Eq. (3.1.12)). This also determines the corresponding structure functions which are introduced by

$$\langle PS|M^{(\tau)}_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle = \frac{2}{M}\mathcal{P}^{(\tau)[\mu\nu]}_{[\alpha\beta]}\Big((S_\mu P_\nu - S_\nu P_\mu)\int_0^1 \mathrm{d}z\, H^{(\tau)}(z)\,\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}\Big).\tag{3.1.49}$$

Applying the same procedure as above we obtain

$$\begin{aligned}\langle PS|M^{\mathrm{tw2}}_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= \frac{2}{M}\int_0^1 \mathrm{d}z\, H^{(2)}(z)\bigg[\Big(S_\alpha(\tilde{x}P)-P_\alpha(\tilde{x}S)\Big)\mathrm{e}^{\mathrm{i}\kappa z(\tilde{x}P)}\\ &\qquad + \tilde{x}_\alpha(\tilde{x}S)M^2\int_0^1 \mathrm{d}\lambda\,\mathrm{i}\kappa z\lambda^2\,\mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)}\bigg],\end{aligned}\tag{3.1.50}$$

$$= \frac{2}{M}\sum_{n=0}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}H_n^{(2)}\Big\{S_\alpha(\tilde{x}P)-P_\alpha(\tilde{x}S)+\frac{n}{n+2}\tilde{x}_\alpha\frac{\tilde{x}S}{\tilde{x}P}M^2\Big\},$$

$$\begin{aligned}\langle PS|M^{\mathrm{tw3}}_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= -2\int_0^1 \mathrm{d}\lambda\int_0^1 \mathrm{d}z\, H^{(3)}(z)\Big[\tilde{x}_\alpha(\tilde{x}S)M\mathrm{i}\kappa z\lambda^2\Big]\mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)},\\ &= -2\sum_{n=1}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}H_n^{(3)}\frac{n}{n+2}\tilde{x}_\alpha\frac{\tilde{x}S}{\tilde{x}P}M.\end{aligned}\tag{3.1.51}$$

The twist–2 part is in agreement with Jaffe and Ji's local expression (see Eq. (45) in [79]) and also with [94]. Adding up both local terms the matrix element of $M_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})$ coincides with the corresponding one of [79]; its nonlocal version reads:

$$\begin{aligned}\langle PS|M_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle =& \frac{2}{M}\Big(S_\alpha(\tilde{x}P)-P_\alpha(\tilde{x}S)\Big)\int_0^1 \mathrm{d}z H^{(2)}(z)\, e_0(\mathrm{i}\zeta z)\\ &+\tilde{x}_\alpha\frac{\tilde{x}S}{\tilde{x}P}M\int_0^1 \mathrm{d}z\Big(H^{(2)}(z)-H^{(3)}(z)\Big)\Big(e_0(\mathrm{i}\zeta z)+2e_2(\mathrm{i}\zeta z)\Big).\end{aligned}\tag{3.1.52}$$

Obviously, the twist-2 distribution function $H^{(2)}(z)$ and the twist-3 distribution function $H^{(3)}(z)$ contribute to the same $1/Q^2$-behaviour of the cross section. A similar *nonlocal* twist–2 matrix element (modulo correction of the trace term) has been given in [92].

Now, we consider the more complicated chiral-odd *skew tensor operator*. Let us note that Jaffe and Ji [79] have not considered the matrix elements of the traceless skew tensor operator.

They have just used the corresponding traceless vector operator in their investigations. We obtain the matrix elements of the skew tensor operators of twist τ by using the projectors determined by Eqs. (3.1.17) – (3.1.19) and Eq. (3.1.49):

$$
\begin{aligned}
\langle PS|M^{\rm tw2}_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= \frac{2}{M}\int_0^1 \mathrm{d}\lambda \int_0^1 \mathrm{d}z\, H^{(2)}(z)\Big[2\lambda\, S_{[\alpha}P_{\beta]}\big(2+\mathrm{i}\kappa\lambda z(\tilde{x}P)\big) \\
&\qquad + (1-\lambda)\, M^2\tilde{x}_{[\alpha}\Big\{4(\mathrm{i}\kappa\lambda z)S_{\beta]} + (\mathrm{i}\kappa\lambda z)^2\big(S_{\beta]}(\tilde{x}P)+P_{\beta]}(\tilde{x}S)\big)\Big\}\Big]\mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)}, \\
&= \frac{2}{M}\sum_{n=0}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}H_n^{(2)}\Big\{2S_{[\alpha}P_{\beta]} + \frac{4n}{(n+2)(n+1)}\frac{M^2}{\tilde{x}P}\tilde{x}_{[\alpha}S_{\beta]} \\
&\qquad + \frac{n(n-1)}{(n+2)(n+1)}\frac{M^2}{\tilde{x}P}\tilde{x}_{[\alpha}\Big(S_{\beta]}+P_{\beta]}\frac{\tilde{x}S}{\tilde{x}P}\Big)\Big\}, \qquad (3.1.53) \\
\langle PS|M^{\rm tw3}_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= -2\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda}\int_0^1 \mathrm{d}z\, \mathrm{i}\kappa\lambda z H^{(3)}(z)M\tilde{x}_{[\alpha}\Big\{S_{\beta]}+\mathrm{i}\kappa\lambda z P_{\beta]}(\tilde{x}S)\Big\} \\
&\qquad \times \mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)}, \qquad (3.1.54) \\
&= -2\sum_{n=1}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}H_n^{(3)}\frac{2}{n+2}\frac{M}{\tilde{x}P}\tilde{x}_{[\alpha}\Big(S_{\beta]}+(n-1)P_{\beta]}\frac{\tilde{x}S}{\tilde{x}P}\Big), \\
\langle PS|M^{\rm tw4}_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= -2\int_0^1 \mathrm{d}\lambda\,\frac{1-\lambda}{\lambda}\int_0^1 \mathrm{d}z\,(\mathrm{i}\lambda\kappa z)^2 H^{(4)}(z)M\tilde{x}_{[\alpha}\Big\{S_{\beta]}(\tilde{x}P)-P_{\beta]}(\tilde{x}S)\Big\} \\
&\qquad \times \mathrm{e}^{\mathrm{i}\kappa\lambda z(\tilde{x}P)}, \\
&= -2\sum_{n=2}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}H_n^{(4)}\frac{n-1}{n+1}\frac{M}{\tilde{x}P}\tilde{x}_{[\alpha}\Big(S_{\beta]}-P_{\beta]}\frac{\tilde{x}S}{\tilde{x}P}\Big). \qquad (3.1.55)
\end{aligned}
$$

Again, the moments of distributions functions of twist $\tau = 2, 3$ and 4 begin with $n = 0, 1$ and 2, respectively. In addition, we remark that only those terms of the operator (3.1.18) contribute to the twist–3 structure function which result from the trace terms of (3.1.17). Analogous to the vector case the forward matrix element of the 'true' twist–3 part of (3.1.18) vanishes. In Eq. (3.1.55) only the twist–4 operator contributes which result from the trace terms of (3.1.17). Let us also mention that after multiplication of (3.1.55) with $\tilde{x}_\alpha$ (or $\tilde{x}_\beta$) the matrix element vanishes because the corresponding vector operator does not contain any twist-4 contribution. This is a simple but important property which may be traced back to the fact that the corresponding Young pattern $[n+2] = (n, 1, 1)$ does allow only n symmetrizations; it is therefore characteristic for any twist–4 skew tensor operator.

The matrix element of the original skew tensor operator is obtained as

$$
\begin{aligned}
\langle PS|M_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= \frac{4}{M}S_{[\alpha}P_{\beta]}\int_0^1 \mathrm{d}z H^{(2)}(z)e_0(\mathrm{i}\zeta z) \qquad (3.1.56) \\
&+ 2\tilde{x}_{[\alpha}P_{\beta]}\frac{\tilde{x}S}{(\tilde{x}P)^2}M\int_0^1 \mathrm{d}z\Big\{H^{(2)}(z)\big[e_0(\mathrm{i}\zeta z)+2e_1(\mathrm{i}\zeta z)+6e_2(\mathrm{i}\zeta z)\big] \\
&- H^{(3)}(z)\big[1+2e_0(\mathrm{i}\zeta z)+6e_2(\mathrm{i}\zeta z)\big] + H^{(4)}(z)\big[1+e_0(\mathrm{i}\zeta z)-2e_1(\mathrm{i}\zeta z)\big]\Big\} \\
&+ 2\tilde{x}_{[\alpha}S_{\beta]}\frac{M}{\tilde{x}P}\int_0^1 \mathrm{d}z\Big\{H^{(2)}(z)\big[e_0(\mathrm{i}\zeta z)-2e_1(\mathrm{i}\zeta z)-2e_2(\mathrm{i}\zeta z)\big] \\
&+ H^{(3)}(z)\big[1+2e_2(\mathrm{i}\zeta z)\big] - H^{(4)}(z)\big[1+e_0(\mathrm{i}\zeta z)-2e_1(\mathrm{i}\zeta z)\big]\Big\}.
\end{aligned}
$$

3.1.4 Relations between quark distribution functions of dynamical and geometric twist

Obviously, since the geometric twist distribution functions are related to traceless operators they differ from the dynamical distribution functions. As far as the scalar LC–operators are concerned which definitely are of twist–2 the new and the old distributions functions coincide.

In order to be able to compare dynamical and geometrical distribution functions, we rewrite the matrix elements (3.1.40), (3.1.47) and (3.1.56) in terms of p_α and $S^\perp_\alpha$. Using this parametrization we obtain for the chiral-even quark distributions

$$\begin{aligned}\langle PS|O_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= 2p_\alpha\frac{\tilde{x}S}{\tilde{x}P}\int_0^1 \mathrm{d}z\, G^{(2)}(z)e_0(\mathrm{i}\zeta z) \qquad (3.1.57)\\ &+2S^\perp_\alpha\int_0^1\mathrm{d}z\Big\{G^{(2)}(z)e_1(\mathrm{i}\zeta z)+G^{(3)}(z)[e_0(\mathrm{i}\zeta z)-e_1(\mathrm{i}\zeta z)]\Big\}\\ &-\tilde{x}_\alpha\frac{\tilde{x}S}{(\tilde{x}P)^2}M^2\int_0^1\mathrm{d}z\Big\{G^{(4)}(z)\Big[e_0(\mathrm{i}\zeta z)-3e_1(\mathrm{i}\zeta z)+2\int_0^1\mathrm{d}\lambda\, e_1(\mathrm{i}\zeta\lambda z)\Big]\\ &-G^{(2)}(z)\Big[e_1(\mathrm{i}\zeta z)-2\int_0^1\mathrm{d}\lambda\, e_1(\mathrm{i}\zeta\lambda z)\Big]+4G^{(3)}(z)\Big[e_1(\mathrm{i}\zeta z)-\int_0^1\mathrm{d}\lambda\, e_1(\mathrm{i}\zeta\lambda z)\Big]\Big\},\end{aligned}$$

and

$$\begin{aligned}\langle P|O_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})|P\rangle &= 2p_\alpha\int_0^1\mathrm{d}z\, F^{(2)}(z)e_0(\mathrm{i}\zeta z)\\ &+\tilde{x}_\alpha\frac{M^2}{\tilde{x}P}\int_0^1\mathrm{d}z\Big\{F^{(4)}(z)e_0(\mathrm{i}\zeta z)+\big[F^{(2)}(z)-F^{(4)}(z)\big]e_1(\mathrm{i}\zeta z)\Big\}, \qquad (3.1.58)\end{aligned}$$

while the chiral-odd distributions are given by

$$\langle P|N(\kappa_1\tilde{x},\kappa_2\tilde{x})|P\rangle = 2M\int_0^1\mathrm{d}z\, E^{(3)}(z)e_0(\mathrm{i}\zeta z), \qquad (3.1.59)$$

and

$$\begin{aligned}\langle PS|M_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})|PS\rangle &= \frac{4}{M}S^\perp_{[\alpha}p_{\beta]}\int_0^1\mathrm{d}z\, H^{(2)}(z)e_0(\mathrm{i}\zeta z) \qquad (3.1.60)\\ &+2\tilde{x}_{[\alpha}p_{\beta]}\frac{\tilde{x}S}{(\tilde{x}P)^2}M\int_0^1\mathrm{d}z\Big\{4H^{(2)}(z)e_2(\mathrm{i}\zeta z)-2H^{(3)}(z)\big[e_0(\mathrm{i}\zeta z)+2e_2(\mathrm{i}\zeta z)\big]\Big\}\\ &-2\tilde{x}_{[\alpha}S^\perp_{\beta]}\frac{M}{\tilde{x}P}\int_0^1\mathrm{d}z\Big\{2H^{(2)}(z)\big[e_1(\mathrm{i}\zeta z)+e_2(\mathrm{i}\zeta z)\big]-H^{(3)}(z)\big[1+2e_2(\mathrm{i}\zeta z)\big]\\ &+H^{(4)}(z)\big[1+e_0(\mathrm{i}\zeta z)-2e_1(\mathrm{i}\zeta z)\big]\Big\}.\end{aligned}$$

Let us note that after multiplication of (3.1.60) by $\tilde{x}_\alpha$ (or $\tilde{x}_\beta$) the last part of the matrix element vanishes. This means that the dynamical twist-4 function $h_3(z)$ can be ignored in the case of the vector operator $M_{5\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ but not in the case of the more general tensor operator $M_{5[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$.

From the expressions (3.1.57) – (3.1.60) and (3.1.5) – (3.1.8) it is obvious that the conventional structure functions of twist $t\geq 3$ contain contributions also of lower geometric twist. By re-expressing the truncated exponentials and performing appropriate variable transformations or using the local expansion of the bilocal operators, we obtain the following relations, which give the relations between the distributions of geometric and dynamical twist. The dynamical twist distributions are given in terms of geometrical twist distributions as follows:

$$g_1(z) = G^{(2)}(z), \tag{3.1.61}$$

$$g_T(z) = G^{(3)}(z) + \int_z^1 \frac{\mathrm{d}y}{y}\Big(G^{(2)} - G^{(3)}\Big)(y)\,, \tag{3.1.62}$$

$$\begin{aligned} 2g_3(z) = -G^{(4)}(z) + \int_z^1 \frac{\mathrm{d}y}{y}\Big\{&\Big(G^{(2)} - 4G^{(3)} + 3G^{(4)}\Big)(y) \\ &+ 2\ln\Big(\frac{z}{y}\Big)\Big(G^{(2)} - 2G^{(3)} + G^{(4)}\Big)(y)\Big\}, \end{aligned} \tag{3.1.63}$$

$$f_1(z) = F^{(2)}(z), \tag{3.1.64}$$

$$2f_4(z) = F^{(4)}(z) + \int_z^1 \frac{\mathrm{d}y}{y}\Big(F^{(2)} - F^{(4)}\Big)(y)\,, \tag{3.1.65}$$

$$e(z) = E^{(3)}(z), \tag{3.1.66}$$

$$h_1(z) = H^{(2)}(z), \tag{3.1.67}$$

$$h_L(z) = H^{(3)}(z) + 2z\int_z^1 \frac{\mathrm{d}y}{y^2}\Big(H^{(2)} - H^{(3)}\Big)(y)\,, \tag{3.1.68}$$

$$\begin{aligned} 2h_3(z) = H^{(4)}(z) + \int_z^1 \frac{\mathrm{d}y}{y}\Big\{&2\Big(H^{(2)} - H^{(4)}\Big)(y) - 2\frac{z}{y}\Big(H^{(2)} - H^{(3)}\Big)(y) \\ &- \delta\Big(\frac{z}{y}\Big)\Big(H^{(3)} - H^{(4)}\Big)(y)\Big\}. \end{aligned} \tag{3.1.69}$$

The relations between the dynamical and geometric quark distribution functions hold for $1 \geq z \geq 0$; the corresponding antiquark distribution functions are obtained for $z \to -z$. We observe that both decompositions coincide in the leading twist, but differ at higher order. For instance, $g_1(z)$ and $h_1(z)$ are proper geometric twist-2 structure functions and $e(z)$ is a proper geometric twist-3 distribution. The distribution functions $g_T(z)$, $f_1(z)$ and $h_L(z)$ with dynamical twist $t = 3$ also contain contributions of geometrical twist $\tau = 2$ and 3. Additionally, the dynamical twist $t = 4$ functions $g_3(z)$ and $h_3(z)$ contain geometrical twist-2, twist-3, as well as twist-4 parts and the dynamical twist $t = 4$ function $f_4(z)$ contains geometrical twist-2 and twist-4.

These relations may be inverted in order to express the geometric twist distributions in terms of dynamical twist distributions. The nontrivial inverse relations are:

$$G^{(3)}(z) = g_T(z) + \frac{1}{z}\int_z^1 \mathrm{d}y\big(g_T - g_1\big)(y)\,, \tag{3.1.70}$$

$$\begin{aligned} G^{(4)}(z) = -\Big\{&2g_3(z) + \frac{1}{z^2}\int_z^1 \mathrm{d}y\, y\big(6g_3 + 4g_T - g_1\big)(y) \\ &+ \frac{1}{z^2}\int_z^1 \mathrm{d}y\, y\Big(1 - \frac{z}{y}\Big)\big(2g_3 + 4g_T - 3g_1\big)(y)\Big\}, \end{aligned} \tag{3.1.71}$$

$$F^{(4)}(z) = 2f_4(z) + \frac{1}{z}\int_z^1 \mathrm{d}y\big(2f_4 - f_1\big)(y)\,, \tag{3.1.72}$$

$$H^{(3)}(z) = h_L(z) + \frac{2}{z}\int_z^1 \mathrm{d}y\big(h_L - h_1\big)(y) \tag{3.1.73}$$

$$H^{(4)}(z) = 2\Big\{h_3(z) + \frac{1}{z^2}\int_z^1 \mathrm{d}y\, y\big(2h_3 - h_L\big)(y) - \frac{1}{z^2}\int_z^1 \mathrm{d}y\, y\Big(1 - \frac{z}{y}\Big)\big(h_L - h_1\big)(y)\Big\}. \tag{3.1.74}$$

The relations between the moments may be read off from Eqs. (3.1.57) – (3.1.60) as follows:

$$g_{1n} = G_n^{(2)}, \tag{3.1.75}$$

$$g_{Tn} = G_n^{(3)} + \frac{1}{n+1}\Big(G_n^{(2)} - G_n^{(3)}\Big), \tag{3.1.76}$$

$$2g_{3n} = -G_n^{(4)} + \frac{1}{n+1}\Big(G_n^{(2)} - 4G_n^{(3)} + 3G_n^{(4)}\Big) - \frac{2}{(n+1)^2}\Big(G_n^{(2)} - 2G_n^{(3)} + G_n^{(4)}\Big), \tag{3.1.77}$$

$$f_{1n} = F_n^{(2)}, \tag{3.1.78}$$

$$2f_{4n} = F_n^{(4)} + \frac{1}{n+1}\Big(F_n^{(2)} - F_n^{(4)}\Big), \tag{3.1.79}$$

$$e_n = E_n^{(3)}, \tag{3.1.80}$$

$$h_{1n} = H_n^{(2)}, \tag{3.1.81}$$

$$h_{Ln} = H_n^{(3)} + \frac{2}{n+2}\Big(H_n^{(2)} - H_n^{(3)}\Big), \tag{3.1.82}$$

$$2h_{3n} = H_n^{(4)} + \frac{2}{n+1}\Big(H_n^{(2)} - H_n^{(4)}\Big) - \frac{2}{n+2}\Big(H_n^{(2)} - H_n^{(3)}\Big) - \delta_{n0}\Big(H_n^{(3)} - H_n^{(4)}\Big). \tag{3.1.83}$$

In terms of the moments the relations between conventional and new distribution functions may be easily inverted. The nontrivial inverse relations are:

$$G_n^{(3)} = g_{Tn} + \frac{1}{n}\Big(g_{Tn} - g_{1n}\Big), \qquad n > 0 \tag{3.1.84}$$

$$G_n^{(4)} = -\Big\{2g_{3n} + \frac{1}{n-1}\Big(6g_{3n} + 4g_{Tn} - g_{1n}\Big) + \frac{1}{n(n-1)}\Big(2g_{3n} + 4g_{Tn} - 3g_{1n}\Big)\Big\}, \qquad n > 1 \tag{3.1.85}$$

$$F_n^{(4)} = 2f_{4n} + \frac{1}{n}\Big(2f_{4n} - f_{1n}\Big), \qquad n > 0 \tag{3.1.86}$$

$$H_n^{(3)} = h_{Ln} + \frac{2}{n}\Big(h_{Ln} - h_{1n}\Big), \qquad n > 0 \tag{3.1.87}$$

$$H_n^{(4)} = 2\Big\{h_{3n} + \frac{1}{n-1}\Big(2h_{3n} - h_{Ln}\Big) - \frac{1}{n(n-1)}\Big(h_{Ln} - h_{1n}\Big)\Big\}, \qquad n > 1. \tag{3.1.88}$$

The nontrivial relationships between the conventional and the new distribution functions are much simpler than to be assumed by a first glance at expressions such as (3.1.40), (3.1.47), and (3.1.56). They show that the conventional distribution functions are determined by the new ones of the same as well as lower geometrical twist, and vice versa (with respect to dynamical twist). Obviously, the same holds for their moments. In principle, this allows to determine, e.g., the new distribution functions from the experimental data if these are known for the conventional ones. At least, this should be possible for the lowest moments.

Another very important consequence is, if the dynamical and the geometrical distribution functions coincide then also their anomalous dimensions coincide. Accordingly, we are able to predict the anomalous dimensions of the geometric distribution functions $F^{(2)}(x, Q^2)$, $G^{(2)}(x, Q^2)$, $H^{(2)}(x, Q^2)$ from the anomalous dimensions of the dynamical ones. The leading order (LO) Q^2 evolution of $f_1(x, Q^2)$ and $g_1(x, Q^2)$ has been known for a long time [58, 70, 2] and the LO Q^2 evolution for $h_1(x, Q^2)$ was also studied in Refs. [116, 7, 94, 29] (in 1-loop calculation with $\mu^2 = Q^2$). For simplicity, we use the local anomalous dimensions. The local geometric twist-2 anomalous dimensions are given by[3] ($C_F = 4/3$)

$$\gamma_n^{F(2)} \equiv \gamma_n^{f_1} = \frac{g^2}{8\pi^2}\, C_F \left(1 - \frac{2}{(n+1)n} + 4\sum_{j=2}^{n}\frac{1}{j}\right), \tag{3.1.89}$$

$$\gamma_n^{G(2)} \equiv \gamma_n^{g_1} = \frac{g^2}{8\pi^2}\, C_F \left(1 - \frac{2}{(n+1)n} + 4\sum_{j=2}^{n}\frac{1}{j}\right), \tag{3.1.90}$$

$$\gamma_n^{H(2)} \equiv \gamma_n^{h_1} = \frac{g^2}{8\pi^2}\, C_F \left(1 + 4\sum_{j=2}^{n}\frac{1}{j}\right), \tag{3.1.91}$$

and hence $\gamma_n^{H(2)} > \gamma_n^{F(2)}$ for all n. The expressions for the anomalous dimensions of higher geometrical twist will be more complicated.

3.1.5 Wandzura-Wilczek type relations

As an immediate consequence of the interrelations (3.1.61) – (3.1.69) and (3.1.70) – (3.1.74) between the conventional structure functions and the new ones of proper geometric twist we are able to derive the decomposition of the conventional structure functions into its parts of genuine twist. Thereby, we also obtain new Wandzura-Wilczek–like relations for the conventional structure functions and also new sum rules of the type of the Burkhardt-Cottingham sum rule.

Let us use the decomposition of $g_T(z) = g_T^{(2)}(z) + g_T^{(3)}(z)$ into its parts $g_T^{(2)}(z)$ and $g_T^{(3)}(z)$ of genuine twist-2 and twist-3, respectively, parts of $g_2(z)$. Then, substituting (3.1.61) into (3.1.62), we get

$$g_T^{(2)}(z) = \int_z^1 \frac{\mathrm{d}y}{y}\, g_1(y) \tag{3.1.92}$$

$$g_T^{(3)}(z) = g_T(z) - \int_z^1 \frac{\mathrm{d}y}{y}\, g_1(y), \tag{3.1.93}$$

where (3.1.92) is the Wandzura-Wilczek relation[4] for the twist-2 part [123]. On the other hand, the analogous relation (3.1.93) for the twist-3 part, which is an immediate consequence of the above definition, has already been obtained in the framework of the local OPE [25] and the nonlocal OPE [96]. Wandzura and Wilczek noted in their original derivation that the twist-3 is small [123]. They obtained in the MIT bag model the approximation

$$\left|G_n^{(3)}/G_n^{(2)}\right| \simeq 0.10\,. \tag{3.1.94}$$

So far the SLAC data [1] agree very well with the Wandzura-Wilczek prediction. However the accuracy of the data is such that a relevant $g_T^{(3)}$-term would not yet be detectable [80]. Therefore, if more precise data of g_T become available, one can investigate the geometric twist-3 part of g_T in more detail.

Using QCD equations of motion operator relations (see [9, 14, 92]), one can decompose the geometric twist-3 part of $g_T(z)$, Eq. (3.1.93), into a quark mass dependent term and a quark-gluon correlation part [119]

$$g_T^{(3)}(z) = \frac{m}{M}\left[\frac{h_1(z)}{z} - \int_z^1 \frac{\mathrm{d}y}{y^2}\, h_1(y)\right] + \tilde{g}_T^{(3)}(z), \tag{3.1.95}$$

---

[3] The corresponding local operators are $O^{\mathrm{tw2}}_{(5)n}(\tilde{x})$ and $M^{\mathrm{tw2}}_{(5)\alpha n}(\tilde{x})$.

[4] With $g_T = g_2 + g_1$ this relation can be rewritten in its original form.

where $\tilde{g}_T^{(3)}(z)$ denotes the twist-3 quark-gluon correlation function of the so-called Shuryak-Vainshtein operator. Therefore, in the massless case $m = 0$ the function $g_T^{(3)}(z)$ measures the quark-gluon correlation in the nucleon.

For the moments we obtain

$$g_{Tn}^{(2)} = \frac{1}{n+1}\, g_{1n}, \tag{3.1.96}$$

$$g_{Tn}^{(3)} = g_{Tn} - \frac{1}{n+1}\, g_{1n}, \qquad n > 0. \tag{3.1.97}$$

Obviously, if we introduce the usual notation $g_T(z) \equiv g_1(z) + g_2(z)$, we get from the relation (3.1.96) for $n = 0$ the Burkhardt-Cottingham sum rule [40]:

$$\int_0^1 \mathrm{d}z\, g_2(z) = 0, \tag{3.1.98}$$

which is consistent with the SLAC data.

Of course, these relationships are well known. However, with the help of the results of the preceeding Section we are able to generalize these relationships also to the other nontrivial structure functions. Using the formulas (3.1.61), (3.1.62), (3.1.63), and (3.1.70), we obtain Wandzura-Wilczek–like integral relations for geometric twist parts of the function $g_3(z) = g_3^{(2)}(z) + g_3^{(3)}(z) + g_3^{(4)}(z)$ as follows:

$$g_3^{(2)}(z) = \int_z^1 \frac{\mathrm{d}y}{y}\Big\{\frac{g_1(y)}{2} + \ln\Big(\frac{z}{y}\Big) g_1(y)\Big\}, \tag{3.1.99}$$

$$g_3^{(3)}(z) = -2\int_z^1 \frac{\mathrm{d}y}{y}\Big\{g_T(y) + \ln\Big(\frac{z}{y}\Big) g_1(y)\Big\}, \tag{3.1.100}$$

$$g_3^{(4)}(z) = g_3(z) + \frac{1}{2}\int_z^1 \frac{\mathrm{d}y}{y}\Big\{\big(4g_T - g_1\big)(y) + 2\ln\Big(\frac{z}{y}\Big) g_1(y)\Big\}. \tag{3.1.101}$$

Due to the fact that $g_3(z)$ contains twist-2, twist-3 as well as twist-4 we have obtained three integral relations. For example, Eq. (3.1.99) demonstrates that the twist-2 part $g_3^{(2)}(z)$ can be expressed in terms of the twist-2 function $g_1(z)$. The relations for the moments are:

$$g_{3n}^{(2)} = \frac{1}{2(n+1)}\, g_{1n} - \frac{1}{(n+1)^2}\, g_{1n}, \tag{3.1.102}$$

$$g_{3n}^{(3)} = -\frac{2}{n+1}\, g_{Tn} + \frac{2}{(n+1)^2}\, g_{1n}, \quad n > 0 \tag{3.1.103}$$

$$g_{3n}^{(4)} = g_{3n} + \frac{1}{2(n+1)}\Big(4g_{Tn} - g_{1n}\Big) - \frac{1}{(n+1)^2}\, g_{1n}, \quad n > 1. \tag{3.1.104}$$

Using again the approximation (3.1.94), one should expect $g_{3n}^{(3)} \ll g_{3n}^{(2)}$. Additionally, we expect that $g_{3n}^{(4)}$ is small. Additionally, from (3.1.103) and (3.1.97) we observe the twist-3 relation

$$g_{3n}^{(3)} = -\frac{2}{n+1}\, g_{Tn}^{(3)}, \quad g_3^{(3)}(z) = -2\int_z^1 \frac{\mathrm{d}y}{y}\, g_T^{(3)}(y). \tag{3.1.105}$$

For $n = 0$ in (3.1.102) we find the sum rule

$$\int_0^1 \mathrm{d}z\, g_3(z) = -\frac{1}{2}\int_0^1 \mathrm{d}z\, g_1(z). \tag{3.1.106}$$

Additionally, we obtain for $n = 1$ in (3.1.102) and (3.1.103) the sum rules

$$\int_0^1 \mathrm{d}z\, z\, g_3^{(2)}(z) = 0, \quad \int_0^1 \mathrm{d}z\, z\, g_3^{(3)}(z) = -\frac{1}{2}\int_0^1 \mathrm{d}z\, z(2g_T - g_1)(z), \tag{3.1.107}$$

which give the following sum rule for $g_3(z)$

$$\int_0^1 \mathrm{d}z\, z\, g_3(z) = -\frac{1}{2}\int_0^1 \mathrm{d}z\, z(2g_T - g_1)(z). \tag{3.1.108}$$

If we use the Efremov-Leader-Teryaev sum rule

$$\int_0^1 \mathrm{d}z\, z(2g_T - g_1)(z) = 0, \tag{3.1.109}$$

which was discussed in [51] in the context of a QCD field theoretical framework, we discover the interesting sum rule from Eq. (3.1.108)

$$\int_0^1 \mathrm{d}z\, z\, g_3(z) = 0. \tag{3.1.110}$$

Let me note that the sum rules (3.1.106) and (3.1.108) should be experimentally verified because they give information about the dynamical twist-4 function $g_3(z)$. Although the function $g_3(z)$ does not contribute to either measurable distribution function including electromagnetic interaction [77], it gives contributions to the three polarized distribution functions in weak interaction processes which are introduced by Ji [82]. Therefore, the *new* sum rules and WW-type relations of $g_3(z)$ will be relevant in these processes.

Substituting (3.1.64) into (3.1.65), we get the integral relations for the twist-2 and twist-4 part of $f_4(z) = f_4^{(2)}(z) + f_4^{(4)}(z)$

$$f_4^{(2)}(z) = \frac{1}{2}\int_z^1 \frac{\mathrm{d}y}{y} f_1(y), \tag{3.1.111}$$

$$f_4^{(4)}(z) = f_4(z) - \frac{1}{2}\int_z^1 \frac{\mathrm{d}y}{y} f_1(y). \tag{3.1.112}$$

The corresponding relations for the moments read

$$f_{4n}^{(2)} = \frac{1}{2(n+1)}\, f_{1n}, \tag{3.1.113}$$

$$f_{4n}^{(4)} = f_{4n} - \frac{1}{2(n+1)}\, f_{1n}, \qquad n > 0. \tag{3.1.114}$$

For $n = 0, 1$ in (3.1.113) we observe the following sum rules

$$\int_0^1 \mathrm{d}z\, f_4(z) = \frac{1}{2}\int_0^1 \mathrm{d}z\, f_1(z), \qquad \int_0^1 \mathrm{d}z\, z\, f_4^{(2)}(z) = \frac{1}{4}\int_0^1 \mathrm{d}z\, z\, f_1(z). \tag{3.1.115}$$

The distributions $h_1(z)$, $h_L(z)$, and $h_3(z)$ can be treated in a similar fashion. Unfortunately, Jaffe and Ji used two different relations between $h_L(z)$ and $h_2(z)$. In their earlier paper [78] they used the notation $h_L(z) \equiv h_2(z) + h_1(z)/2$. But later [79] they introduced the relation $h_L(z) \equiv h_1(z) + h_2(z)/2$. In order to avoid any misunderstanding we use $h_L(z)$ instead of $h_2(z)$.

If we now substitute (3.1.67) into (3.1.68) and use $h_L(z) = h_L^{(2)}(z) + h_L^{(3)}(z)$, we derive

$$h_L^{(2)}(z) = 2z \int_z^1 \frac{\mathrm{d}y}{y^2}\, h_1(y), \tag{3.1.116}$$

$$h_L^{(3)}(z) = h_L(z) - 2z \int_z^1 \frac{\mathrm{d}y}{y^2}\, h_1(y), \tag{3.1.117}$$

where (3.1.116) is a twist-2 Wandzura-Wilczek–like relation which was obtained earlier by Jaffe and Ji [79] (using the notation $h_L(z) \equiv h_2(z) + h_1(z)/2$ we recover the Wandzura-Wilczek relation of the function $h_2(z)$ which was obtained in Ref. [78]). Obviously, (3.1.117) is the corresponding twist-3 relation which can be decomposed by means of QCD equations of motion operator relations as follows (see [79, 119, 92])

$$h_L^{(3)}(z) = \frac{m}{M}\left[\frac{g_1(z)}{z} - 2z \int_z^1 \frac{\mathrm{d}y}{y^3}\, g_1(y)\right] + \tilde{h}_L^{(3)}(z), \tag{3.1.118}$$

where $\tilde{h}_L^{(3)}(z)$ denotes the quark-gluon correlation function of the twist-3 part of the trilocal quark-gluon operator $V_{\alpha\beta}^{21}(x, tx, -x)$ (see Section 4.3).

For the moments we obtain

$$h_{Ln}^{(2)} = \frac{2}{n+2}\, h_{1n}, \tag{3.1.119}$$

$$h_{Ln}^{(3)} = h_{Ln} - \frac{2}{n+2}\, h_{1n}, \qquad n > 0. \tag{3.1.120}$$

For $n = 0, 1$ in (3.1.119) we observe the following sum rule

$$\int_0^1 \mathrm{d}z\, h_L(z) = \int_0^1 \mathrm{d}z\ h_1(z), \tag{3.1.121}$$

$$\int_0^1 \mathrm{d}z\, z\, h_L^{(2)}(z) = \frac{2}{3}\int_0^1 \mathrm{d}z\, z\, h_1(z). \tag{3.1.122}$$

The sum rule (3.1.121) was earlier derived in Ref. [41]. If we use the notation $h_L(z) \equiv h_1(z) + h_2(z)/2$, we recover in (3.1.121) a sum rule very similar to the Burkhardt-Cottingham sum rule, namely $\int_0^1 \mathrm{d}z\, h_2(z) = 0$, which was originally derived by Tangerman and Mulders [119].

Furthermore, using the formulas (3.1.67), (3.1.68) (3.1.69) and (3.1.73), we obtain the integral relations for the structure function $h_3(z) = h_3^{(2)}(z) + h_3^{(3)}(z) + h_3^{(4)}(z)$ as follows:

$$h_3^{(2)}(z) = \frac{1}{z}\int_z^1 \mathrm{d}y \Big(\frac{z}{y} - \frac{z^2}{y^2}\Big) h_1(y), \tag{3.1.123}$$

$$h_3^{(3)}(z) = \frac{1}{z}\int_z^1 \mathrm{d}y \Big\{h_L(y) - \Big(1 - \frac{z^2}{y^2}\Big) h_1(y)\Big\}, \tag{3.1.124}$$

$$h_3^{(4)}(z) = h_3(z) - \frac{1}{z}\int_z^1 \mathrm{d}y \Big\{h_L(y) - \Big(1 - \frac{z}{y}\Big) h_1(y)\Big\}, \tag{3.1.125}$$

and the relations for the corresponding moments

$$h_{3n}^{(2)} = \frac{1}{n+1}\, h_{1n} - \frac{1}{n+2}\, h_{1n}, \tag{3.1.126}$$

$$h_{3n}^{(3)} = \frac{1}{n}\, h_{Ln} - \frac{2}{(n+2)n}\, h_{1n}, \qquad n > 0 \tag{3.1.127}$$

$$h_{3n}^{(4)} = h_{3n} - \frac{1}{n}\, h_{Ln} + \frac{1}{(n+1)n}\, h_{1n}, \qquad n > 1. \tag{3.1.128}$$

From (3.1.127) and (3.1.120) we observe the following twist-3 relation

$$h^{(3)}_{3n} = \frac{1}{n}\, h^{(3)}_{Ln}, \quad h^{(3)}_3(z) = \frac{1}{z}\int_z^1 \mathrm{d}y\, h^{(3)}_L(y). \tag{3.1.129}$$

Therefore we conclude that $h_3^{(2)}$ has a simple parton interpretation like h_1, $h_3^{(3)}$ represent the quark-gluon correlation, like g_T, in the nucleon and last but not least the distribution $h_3^{(4)}$ is determined by the rather complicated multi quark-gluon and quark-quark correlation in the nucleon. Similar arguments are valid for the different parts of g_3.

Here we also obtain a interesting sum rule from (3.1.126) in the case $n = 0$, namely,

$$\int_0^1 \mathrm{d}z\, h_3(z) = \frac{1}{2}\int_0^1 \mathrm{d}z\, h_1(z), \tag{3.1.130}$$

which was also obtained in Ref. [41]. For $n = 1$ we get from Eqs. (3.1.126) and (3.1.127) the following sum rules

$$\int_0^1 \mathrm{d}z\, z\, h_3^{(2)}(z) = \frac{1}{6}\int_0^1 \mathrm{d}z\, z\, h_1(z), \tag{3.1.131}$$

$$\int_0^1 \mathrm{d}z\, z\, h_3^{(3)}(z) = \int_0^1 \mathrm{d}z\, z\Big(h_L(z) - \frac{2}{3}\, h_1(z)\Big), \tag{3.1.132}$$

which give the phenomenological interesting sum rule for $h_3(z)$:

$$\int_0^1 \mathrm{d}z\, z\, h_3(z) = \int_0^1 \mathrm{d}z\, z\Big(h_L(z) - \frac{1}{2}\, h_1(z)\Big). \tag{3.1.133}$$

The sum rules (3.1.130) and (3.1.133) give information about the dynamical twist-4 function $h_3(z)$ from the functions $h_1(z)$ and $h_2(z)$ and should be phenomenologically tested.

Let us note that Jaffe and Ji ignored in their investigations [79] the function $h_3(z)$ because they claimed that it is twist-4 and can be neglected. However, this assumption is only true in the case of the vector operator $M_{5\alpha}(\kappa_1\tilde{x}, \kappa_2\tilde{x})$ but not in the case of the tensor operator $M_{5[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x})$ which appear in polarized deep inelastic scattering by means of the massive quark propagator. As we have seen in Eq. (3.1.69), then the function $h_3(z)$ contains twist-2, twist-3 as well as twist-4 parts. Moreover, the twist-2 part $h_3^{(2)}(z)$ is given in terms of the twist-2 function $h_1(z)$ in (3.1.123). Therefore, the structure function $h_3(z)$ is experimentally relevant also at the twist-2 and twist-3 level and cannot be ignored in these cases. The dynamical twist-4 function $h_3(z)$ is, in principle, observable in the polarised Drell-Yan process [73].

3.1.6 Conclusions

Using the notion of geometric twist, we discussed the calculation of the forward matrix elements for those nonlocal LC-operators which correspond to the independent tensor structures in polarized nucleon matrix elements of the type $\langle PS|\bar{\psi}(\kappa_1\tilde{x})\Gamma U(\kappa_1\tilde{x}, \kappa_2\tilde{x})\psi(\kappa_2\tilde{x})|PS\rangle$. We have found nine independent forward distribution functions with well defined twist τ which, for twist $\tau \geq 3$, differ from the conventional ones. From the field theoretical point of view this Lorentz invariant classification is the most appropriate frame of introducing distribution functions since the separation of different (geometric) twist is unique and independent from the special kinematics of the process. Only the operators of definite geometric twist will have the correct mixing behaviour under renormalization. A very useful property of the nonlocal (and local) LC operators of definite twist τ is that they are obtained from the original, i.e., undecomposed LC–operators by the application of corresponding projection operators of geometric twist. An essential result of our calculations is the relation between the new parton distribution functions and those given by Jaffe and Ji [78, 79]. In addition we have also given the moments of all the distribution functions. These relations demonstrate the interrelations between the different twist definitions. One principal result are Wandzura-Wilczek–like relations between the dynamical twist distributions which we have obtained by means of interrelation rules in a natural manner. An advantage in our approach was that we used operators with geometric twist which allowed us to reveal the Wandzura-Wilczek–like relations between the dynamical twist distribution functions.

3.2 Vector meson distribution amplitudes of definite geometric twist

In this Section, we centre our discussion around the specific case of light vector-meson distribution amplitudes (DAs), which are relevant for describing light-cone dominated processes, e.g., the vector meson production off nucleons, $\gamma^* + N \to V + N$ and can be expressed in terms of matrix-elements of gauge-invariant non-local operators sandwiched between the vacuum and the meson state, $\langle 0|\bar{u}(x)\Gamma U(x,-x)d(-x)|\rho(P,\lambda)\rangle$. In particular, we investigate the (two-particle) *ρ-meson light-cone distribution amplitudes* of geometric twist. We will find eight meson distribution amplitudes of geometric twist.

On the other hand, Ball *et al.* used the notion of *dynamical twist t* for the classification of (two-particle) vector meson distribution amplitudes and vector meson distribution amplitudes, respectively [13, 14, 15]. They classified the meson distribution amplitudes which correspond to the independent tensor structure of the matrix elements of bilocal quark-antiquark operators. One key ingredient in their approach was the use of QCD equations of motion in order to obtain dynamical Wandzura-Wilczek-type relations for DAs that are not dynamically independent. The pion distribution functions were investigated in similar way [32, 33, 16].

Meson distribution amplitudes and form factors have been discussed in the framework of local operator product expansion and in the infinite momentum frame by [99, 36, 50, 42, 116, 44] and in the framework of nonlocal operator product expansion [60, 35, 106].

The aim of this Section is to present the (two-particle) *meson distribution amplitudes* which are related to the nonlocal LC-operators of different geometric twist. In the framework of geometric twist, it is possible to investigate the interrelations and the mismatch between the different twist definitions of meson distribution amplitudes as a kind of group theoretical relations. By means of this relations, we are able to obtain geometric Wandzura-Wilczek-type relations between the the dynamical twist function which differ from the dynamical Wandzura-Wilczek-type relations.

3.2.1 The ρ-meson distribution amplitudes of definite dynamical twist

Usually, the meson light-cone distributions are defined as the vacuum-to-meson matrix elements of quark-antiquark nonlocal gauge invariant operators on the light-cone. In this way, Ball *et al.* [13, 14, 15] found eight independent two-particle distributions. The classification of these dynamical twist ρ-meson distribution amplitudes with respect to spin, dynamical twist, chirality and the relations to geometrical twist (see Section 3.2.3) are summarized in Tab. 3.2. One distri-

Twist t	2	3	4
	$O(1)$	$O(1/Q)$	$O(1/Q^2)$
$e_\parallel$	$\hat{\phi}_\parallel \equiv \hat{\Phi}^{(2)}$	$\underline{\hat{h}_\parallel^{(t)}} = \hat{\Phi}^{(3)} + F_t(\hat{\Phi}^{(2)}, \hat{\Phi}^{(3)})$ $\underline{\hat{h}_\parallel^{(s)}} \equiv \hat{\Upsilon}^{(3)}$	$\hat{g}_3 = \hat{\Phi}^{(4)} + F_g(\hat{\Phi}^{(2)}, \hat{\Phi}^{(3)}, \hat{\Phi}^{(4)})$
$e_\perp$	$\underline{\hat{\phi}_\perp} \equiv \hat{\Psi}^{(2)}$	$\hat{g}_\perp^{(v)} = \hat{\Psi}^{(3)} + F_v(\hat{\Psi}^{(2)}, \hat{\Psi}^{(3)})$ $\hat{g}_\perp^{(a)} \equiv \hat{\Xi}^{(3)}$	$\underline{\hat{h}_3} = \hat{\Psi}^{(4)} + F_h(\hat{\Psi}^{(2)}, \hat{\Psi}^{(3)}, \hat{\Psi}^{(4)})$

Table 3.2: Spin, dynamical twist and chiral classification of the ρ-meson distribution amplitudes.

bution amplitude was obtained for longitudinally ($e_\parallel$) and transversely ($e_\perp$) polarized ρ-mesons of twist-2 and twist-4, respectively. Whereas, the number of twist-3 distribution amplitudes is doubled for each polarization. The higher twist distribution amplitudes contribute to a hard exclusive amplitude with additional powers of $1/Q$ compared to the leading twist-2 ones. The underlined distribution amplitudes are chiral-odd, the others chiral-even.

Using the Lorentz-tensor decomposition of the matrix elements of relevant bilocal operators,

$$O_\alpha(\tilde{x},-\tilde{x}) = \bar{u}(\tilde{x})\gamma_\alpha U(\tilde{x},-\tilde{x})d(-\tilde{x}), \tag{3.2.1}$$

$$O_{5\alpha}(\tilde{x},-\tilde{x}) = \bar{u}(\tilde{x})\gamma_\alpha\gamma_5 U(\tilde{x},-\tilde{x})d(-\tilde{x}), \tag{3.2.2}$$

$$M_{[\alpha\beta]}(\tilde{x},-\tilde{x}) = \bar{u}(\tilde{x})\sigma_{\alpha\beta} U(\tilde{x},-\tilde{x})d(-\tilde{x}), \tag{3.2.3}$$

$$N(\tilde{x},-\tilde{x}) = \bar{u}(\tilde{x}) U(\tilde{x},-\tilde{x})d(-\tilde{x}), \tag{3.2.4}$$

in analogy to nucleon structure functions, the explicit definitions of the chiral-even ρ-distributions are:

$$\begin{aligned}\langle 0|O_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = f_\rho m_\rho \Bigg[& p_\alpha \frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P}\int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{\phi}_\parallel(\xi,\mu^2) + e^{(\lambda)}_{\perp\alpha}\int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{g}^{(v)}_\perp(\xi,\mu^2) \\ & -\frac{1}{2}\tilde{x}_\alpha \frac{e^{(\lambda)}\tilde{x}}{(\tilde{x}P)^2} m_\rho^2 \int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{g}_3(\xi,\mu^2)\Bigg]\end{aligned} \tag{3.2.5}$$

and

$$\langle 0|O_{5\alpha}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = \frac{1}{2}\left(f_\rho - f_\rho^{\mathrm{T}}\frac{m_u+m_d}{m_\rho}\right) m_\rho \epsilon_\alpha{}^{\beta\mu\nu} e^{(\lambda)}_{\perp\beta} p_\mu \tilde{x}_\nu \int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{g}^{(a)}_\perp(\xi,\mu^2), \tag{3.2.6}$$

while the chiral-odd distributions are defined as

$$\begin{aligned}\langle 0|M_{[\alpha\beta]}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = \mathrm{i}f_\rho^{\mathrm{T}} \Bigg[& (e^{(\lambda)}_{\perp\alpha}p_\beta - e^{(\lambda)}_{\perp\beta}p_\alpha)\int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{\phi}_\perp(\xi,\mu^2) \\ & + (p_\alpha\tilde{x}_\beta - p_\beta\tilde{x}_\alpha)\frac{e^{(\lambda)}\tilde{x}}{(\tilde{x}P)^2}m_\rho^2 \int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{h}^{(t)}_\parallel(\xi,\mu^2) \\ & +\frac{1}{2}(e^{(\lambda)}_{\perp\alpha}\tilde{x}_\beta - e^{(\lambda)}_{\perp\beta}\tilde{x}_\alpha)\frac{m_\rho^2}{\tilde{x}P}\int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{h}_3(\xi,\mu^2)\Bigg]\end{aligned} \tag{3.2.7}$$

and

$$\langle 0|N(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = -\mathrm{i}\left(f_\rho^{\mathrm{T}} - f_\rho\frac{m_u+m_d}{m_\rho}\right)(e^{(\lambda)}\tilde{x})m_\rho^2\int_{-1}^{1}\mathrm{d}\xi\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}\hat{h}^{(s)}_\parallel(\xi,\mu^2). \tag{3.2.8}$$

In the above definitions we used the two light-cone vectors (see Eq. (2.1.17) with $\rho_\alpha = P_\alpha/m_\rho$)

$$\tilde{x}_\alpha = x_\alpha - \frac{P_\alpha}{m_\rho^2}\Big((xP) - \sqrt{(xP)^2 - x^2 m_\rho^2}\Big), \qquad p_\alpha = P_\alpha - \frac{1}{2}\tilde{x}_\alpha\frac{m_\rho^2}{\tilde{x}P}, \tag{3.2.9}$$

with $p^2 = 0$, $\tilde{x}^2 = 0$ and $p\tilde{x} = P\tilde{x}$. Here P_α is the ρ-meson momentum vector, so that $P^2 = m_\rho^2$, $e^{(\lambda)}e^{(\lambda)} = -1$, $Pe^{(\lambda)} = 0$, m_ρ denotes the ρ-meson mass. The polarization vector of the ρ-meson $e^{(\lambda)}_\alpha$ can be decomposed into projections onto the two light-like vectors and the orthogonal plane:

$$e^{(\lambda)}_\alpha = p_\alpha\frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P} - \frac{1}{2}\tilde{x}_\alpha\frac{e^{(\lambda)}\tilde{x}}{(\tilde{x}P)^2}m_\rho^2 + e^{(\lambda)}_{\perp\alpha}. \tag{3.2.10}$$

The distribution amplitudes are dimensionless functions of ξ and describe the probability amplitudes to find the ρ-meson in a state with minimal number of constituents (quark and anti-quark) which carry momentum fractions ξ. For example, the functions $\hat{\phi}_{\parallel}$ and $\hat{\phi}_{\perp}$ give the leading twist distributions in the fraction of total momentum carried by the quark in longitudinally and transversely polarized mesons, respectively [3]. The nonlocal operators are renormalized at scale μ, so that the distribution amplitudes depend on μ as well. This dependence can be calculated in perturbative QCD (from now on we suppress μ^2).

The vector and tensor decay constants f_ρ and f_ρ^{T} are defined as usually as

$$\begin{aligned}\langle 0|\bar{u}(0)\gamma_\alpha d(0)|\rho(P,\lambda)\rangle &= f_\rho m_\rho e_\alpha^{(\lambda)}, &(3.2.11)\\ \langle 0|\bar{u}(0)\sigma_{\alpha\beta} d(0)|\rho(P,\lambda)\rangle &= \mathrm{i} f_\rho^{\mathrm{T}}(e_\alpha^{(\lambda)}P_\beta - e_\beta^{(\lambda)}P_\alpha). &(3.2.12)\end{aligned}$$

All eight distributions $\hat{\phi} = \{\hat{\phi}_\parallel, \hat{\phi}_\perp, \hat{g}_\perp^{(v)}, \hat{g}_\perp^{(a)}, \hat{h}_\parallel^{(t)}, \hat{h}_\parallel^{(s)}, \hat{h}_3, \hat{g}_3\}$ are normalized as

$$\int_{-1}^{1} \mathrm{d}\xi\, \hat{\phi}(\xi) = 1. \qquad (3.2.13)$$

The distribution amplitude moments related to the reduced matric elements are given as

$$\phi_n = \int_{-1}^{1} \mathrm{d}\xi\, \xi^n \hat{\phi}(\xi) \equiv \int_{0}^{1} \mathrm{d}u\, \xi^n \phi(u), \qquad \xi = 2u-1 \qquad (3.2.14)$$

with both types of meson distribution amplitudes

$$\hat{\phi}(\xi) = \frac{1}{2}\,\phi\left(\frac{\xi+1}{2}\right), \qquad \phi(u) = 2\hat{\phi}(2u-1). \qquad (3.2.15)$$

Eq. (3.2.14) means, having information about the distribution amplitude moments, one can reconstruct the distribution amplitude itself.

3.2.2 The ρ-meson matrix elements of LC–operators with definite geometric twist

In this Section we define the meson distribution amplitudes for the bilinear LC-quark operators with definite twist sandwiched between the vacuum and the meson state. The twist projectors for the bilocal operators (3.2.1)–(3.2.4) are given in Eqs. (3.1.14) – (3.1.21). As usual, the matrix elements of the meson are related to the meson momentum P_μ and meson polarization vector $e_\mu^{(\lambda)}$, respectively and $\tilde{x}_\alpha$ is generated by the twist projectors. According to the above projection properties we are able to introduce one (and only one) distribution amplitude for any operator of definite twist.

We start with the chiral-even *scalar operator* $O^{\mathrm{tw}2}(\tilde{x},-\tilde{x}) = \bar{u}(\tilde{x})(\gamma\tilde{x})U(\tilde{x},-\tilde{x})d(-\tilde{x})$. The matrix element of this nonlocal twist-2 operator taken between the vacuum $\langle 0|$ and the meson state $|\rho(P,\lambda)\rangle$ reads

$$\langle 0|O^{\mathrm{tw}2}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = f_\rho m_\rho (e^{(\lambda)}\tilde{x}) \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Phi}^{(2)}(\xi)\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)} = f_\rho m_\rho (e^{(\lambda)}\tilde{x}) \sum_{n=0}^{\infty} \frac{(\mathrm{i}(\tilde{x}P))^n}{n!}\, \Phi_n^{(2)}. \qquad (3.2.16)$$

Here, $\hat{\Phi}^{(2)}(\xi)$ is the twist-2 meson distribution amplitude which is related to the corresponding moments

$$\Phi^{(2)}_n = \int_{-1}^{1} \mathrm{d}\xi\, \xi^n \hat{\Phi}^{(2)}(\xi).$$

Now we consider the chiral-even *vector operator*. Using the projection property (3.1.22), we introduce the meson distribution amplitudes $\hat{\Phi}^{(\tau)}(\xi)$ of geometric twist $\tau$by

$$\langle 0|O^{(\tau)}_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle \equiv \langle 0|(\mathcal{P}^{(\tau)\beta}_\alpha O^{(\tau)}_\beta)(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = f_\rho m_\rho \mathcal{P}^{(\tau)\beta}_\alpha \Big(e^{(\lambda)}_\beta \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Phi}^{(\tau)}(\xi)\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)} \Big)\ , \tag{3.2.17}$$

which, for $\tau = 2$, is consistent with (3.2.16). Using the twist projection operators as they are determined by Eqs. (3.1.14) – (3.1.16) we obtain for the twist-2 operator

$$\begin{aligned}
\langle 0|O^{\mathrm{tw2}}_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= f_\rho m_\rho \int_0^1 \mathrm{d}t \Big[\partial_\alpha + \frac{1}{2}\big(\ln t\big) x_\alpha \Box\Big](e^{(\lambda)}x) \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Phi}^{(2)}(\xi)\, \mathrm{e}^{\mathrm{i}\xi t(xP)}\big|_{x=\tilde{x}},\\
&= f_\rho m_\rho \int_0^1 \mathrm{d}t \int_{-1}^{1} \mathrm{d}\xi\, \Phi^{(2)}(\xi) \Big[e^{(\lambda)}_\alpha + \mathrm{i}\xi t P_\alpha (e^{(\lambda)}\tilde{x}) + \frac{\tilde{x}_\alpha}{2}(e^{(\lambda)}\tilde{x}) m^2_\rho (\mathrm{i}\xi t)^2 \big(\ln t\big)\Big] \mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)},\\
&= f_\rho m_\rho \sum_{n=0}^{\infty} \frac{(\mathrm{i}(\tilde{x}P))^n}{n!} \Phi^{(2)}_n \Big\{ \frac{1}{n+1}\Big(e^{(\lambda)}_\alpha + nP_\alpha \frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P}\Big) - \frac{n(n-1)}{2(n+1)^2} \tilde{x}_\alpha \frac{m^2_\rho (e^{(\lambda)}\tilde{x})}{(\tilde{x}P)^2} \Big\},
\end{aligned} \tag{3.2.18}$$

and for the higher twist operators

$$\begin{aligned}
\langle 0|O^{\mathrm{tw3}}_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= f_\rho m_\rho \int_0^1 \mathrm{d}t \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Phi}^{(3)}(\xi) \Big[\Big(e^{(\lambda)}_\alpha (\tilde{x}P) - P_\alpha (e^{(\lambda)}\tilde{x})\Big) \mathrm{i}\xi t\\
&\qquad\qquad - \tilde{x}_\alpha m^2_\rho (e^{(\lambda)}\tilde{x}) (\mathrm{i}\xi t)^2 \ln t\Big] \mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)},\\
&= f_\rho m_\rho \sum_{n=1}^{\infty} \frac{(\mathrm{i}(\tilde{x}P))^n}{n!} \Phi^{(3)}_n \Big\{ \frac{n}{n+1}\Big(e^{(\lambda)}_\alpha - P_\alpha \frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P}\Big) + \frac{n(n-1)}{(n+1)^2} \tilde{x}_\alpha \frac{m^2_\rho (e^{(\lambda)}\tilde{x})}{(\tilde{x}P)^2} \Big\},
\end{aligned} \tag{3.2.19}$$

$$\begin{aligned}
\langle 0|O^{\mathrm{tw4}}_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= \frac{1}{2} f_\rho m_\rho \tilde{x}_\alpha (e^{(\lambda)}\tilde{x}) m^2_\rho \int_0^1 \mathrm{d}t \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Phi}^{(4)}(\xi) (\mathrm{i}\xi t)^2 (\ln t)\, \mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)},\\
&= -f_\rho m_\rho \sum_{n=2}^{\infty} \frac{(\mathrm{i}(\tilde{x}P))^n}{n!} \Phi^{(4)}_n \frac{n(n-1)}{2(n+1)^2} \tilde{x}_\alpha \frac{m^2_\rho (e^{(\lambda)}\tilde{x})}{(\tilde{x}P)^2}.
\end{aligned} \tag{3.2.20}$$

Obviously, the trace terms which have been explicitly subtracted are proportional to m^2_ρ. They give rise to light-cone mass correction terms of mesons and contribute to *dynamical* twist-4. For the twist–2 operator we observe that the terms proportional to $e^{(\lambda)}_\alpha$, P_α and $\tilde{x}_\alpha$ have contributions starting with the zeroth, first and second moment, respectively. The twist–3 operator starts with the first moment, and the twist–4 operator starts with the second moment.

If we put together the different twist contributions we obtain, after replacing $\mathrm{i}\xi t(\tilde{x}P)$ by

$t\partial/\partial t$ and performing partial integrations, the following matrix element of the original operator

$$\begin{aligned}\langle 0|O_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= f_\rho m_\rho \Bigg[P_\alpha \frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P} \int_{-1}^{1} \mathrm{d}\xi \Big(\hat{\Phi}^{(2)}(\xi) - \hat{\Phi}^{(3)}(\xi) \Big) [e_0(\mathrm{i}\zeta\xi) - e_1(\mathrm{i}\zeta\xi)] \\ &+ e_\alpha^{(\lambda)} \int_{-1}^{1} \mathrm{d}\xi \Big(\hat{\Phi}^{(2)}(\xi) e_1(\mathrm{i}\zeta\xi) + \hat{\Phi}^{(3)}(\xi) [e_0(\mathrm{i}\zeta\xi) - e_1(\mathrm{i}\zeta\xi)] \Big) \\ &- \frac{1}{2} \tilde{x}_\alpha \frac{e^{(\lambda)}\tilde{x}}{(\tilde{x}P)^2} m_\rho^2 \int_{-1}^{1} \mathrm{d}\xi \Big(\hat{\Phi}^{(2)}(\xi) - 2\hat{\Phi}^{(3)}(\xi) + \hat{\Phi}^{(4)}(\xi) \Big) \Big[e_0(\mathrm{i}\zeta\xi) - 3e_1(\mathrm{i}\zeta\xi) + 2\int_0^1 \mathrm{d}t\, e_1(\mathrm{i}\zeta\xi t) \Big] \Bigg],\end{aligned} \tag{3.2.21}$$

where we used again the "truncated exponentials" and $\zeta = (\tilde{x}P)$. Let us mention that the twist-2 part of Eq. (3.2.21) and the local expression (3.2.18) are in agreement with Stoll's result [118] for the light-cone twist-2 operator.

Now we consider the chiral-even *axial vector operator* $O_{5\alpha}(\tilde{x},-\tilde{x})$, which obeys relations Eqs. (3.1.14) – (3.1.16), as well as (3.1.22) with the *same* projection operator as the vector operator. Let us introduce the corresponding meson distribution amplitudes $\Xi^{(\tau)}(z)$ of twist τ by

$$\langle 0|O_{5\alpha}^{(\tau)}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = \frac{1}{2}\Big(f_\rho - f_\rho^{\mathrm{T}} \frac{m_u+m_d}{m_\rho} \Big) m_\rho \mathcal{P}_\alpha^{(\tau)\beta} \Big(\epsilon_\beta{}^{\gamma\mu\nu} e_\gamma^{(\lambda)} P_\mu \tilde{x}_\nu \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Xi}^{(\tau)}(\xi)\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)} \Big), \tag{3.2.22}$$

where f_ρ^{T} denotes the tensor decay constant. The vacuum-to-meson matrix elements of these *vector operators* of geometric twist are obtained as follows:

$$\begin{aligned}\langle 0|O_{5\alpha}^{\mathrm{tw3}}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= \frac{1}{2}\Big(f_\rho - f_\rho^{\mathrm{T}} \frac{m_u+m_d}{m_\rho} \Big) m_\rho \epsilon_\alpha{}^{\beta\mu\nu} e_\beta^{(\lambda)} P_\mu \tilde{x}_\nu \int_0^1 \mathrm{d}t \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Xi}^{(3)}(\xi) \\ &\qquad \times \big(1 + \mathrm{i}\xi t(\tilde{x}P)\big) \mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)} \\ &= \frac{1}{2}\Big(f_\rho - f_\rho^{\mathrm{T}} \frac{m_u+m_d}{m_\rho} \Big) m_\rho \epsilon_\alpha{}^{\beta\mu\nu} e_\beta^{(\lambda)} P_\mu \tilde{x}_\nu \sum_{n=0}^{\infty} \frac{(\mathrm{i}(\tilde{x}P))^n}{n!}\, \Xi_n^{(3)}.\end{aligned} \tag{3.2.23}$$

We see that only the twist-3 operator gives a contribution and the twist-2 and twist-4 operator as well as all trace terms vanish. Thus, the matrix element of the original operator reads

$$\langle 0|O_{5\alpha}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = \frac{1}{2}\Big(f_\rho - f_\rho^{\mathrm{T}} \frac{m_u+m_d}{m_\rho} \Big) m_\rho \epsilon_\alpha{}^{\beta\mu\nu} e_\beta^{(\lambda)} P_\mu \tilde{x}_\nu \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Xi}^{(3)}(\xi) e_0(\mathrm{i}\zeta\xi). \tag{3.2.24}$$

The matrix element of the simplest bilocal *scalar operator* arises as

$$\begin{aligned}\langle 0|N(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= -\mathrm{i}\Big(f_\rho^{\mathrm{T}} - f_\rho \frac{m_u+m_d}{m_\rho} \Big) \big(e^{(\lambda)}\tilde{x}\big) m_\rho^2 \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Upsilon}^{(3)}(\xi) \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)}, \\ &= -\mathrm{i}\Big(f_\rho^{\mathrm{T}} - f_\rho \frac{m_u+m_d}{m_\rho} \Big) \big(e^{(\lambda)}\tilde{x}\big) m_\rho^2 \sum_{n=0}^{\infty} \frac{(\mathrm{i}(\tilde{x}P))^n}{n!}\, \Upsilon_n^{(3)},\end{aligned} \tag{3.2.25}$$

where $\hat{\Upsilon}^{(3)}(\xi)$ is another spin-independent twist-3 meson DA.

Now, we consider the matrix elements of the chiral-odd *skew tensor operators*. The corresponding distribution amplitudes are introduced by

$$\langle 0|M_{[\alpha\beta]}^{(\tau)}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = \mathrm{i} f_\rho^{\mathrm{T}} \mathcal{P}_{[\alpha\beta]}^{(\tau)[\mu\nu]} \Big((e_\mu^{(\lambda)} P_\nu - e_\nu^{(\lambda)} P_\mu) \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Psi}^{(\tau)}(\xi)\, \mathrm{e}^{\mathrm{i}\xi(\tilde{x}P)} \Big). \tag{3.2.26}$$

The matrix elements of the skew tensor operators of twist τ are obtained using the projectors determined by Eqs. (3.1.17) – (3.1.19):

$$
\begin{aligned}
\langle 0|M^{\mathrm{tw2}}_{[\alpha\beta]}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= \mathrm{i} f^{\mathrm{T}}_{\rho}\int_0^1 \mathrm{d}t \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Psi}^{(2)}(\xi)\Big[2t\, e^{(\lambda)}_{[\alpha} P_{\beta]}\big(2+\mathrm{i}\xi t(\tilde{x}P)\big) \qquad (3.2.27)\\
&\quad + (1-t)\, m_\rho^2 \tilde{x}_{[\alpha}\Big\{4(\mathrm{i}\xi t)e^{(\lambda)}_{\beta]} + (\mathrm{i}\xi t)^2\big(e^{(\lambda)}_{\beta]}(\tilde{x}P) + P_{\beta]}(e^{(\lambda)}\tilde{x})\big)\Big\}\Big]\mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)},\\
&= \mathrm{i} f^{\mathrm{T}}_{\rho}\sum_{n=0}^{\infty}\frac{(\mathrm{i}(\tilde{x}P))^n}{n!}\Psi^{(2)}_n\Big\{2e^{(\lambda)}_{[\alpha}P_{\beta]} + \frac{4n}{(n+2)(n+1)}\frac{m_\rho^2}{\tilde{x}P}\tilde{x}_{[\alpha}e^{(\lambda)}_{\beta]}\\
&\quad + \frac{n(n-1)}{(n+2)(n+1)}\frac{m_\rho^2}{\tilde{x}P}\tilde{x}_{[\alpha}\Big(e^{(\lambda)}_{\beta]} + P_{\beta]}\frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P}\Big)\Big\},\\
\langle 0|M^{\mathrm{tw3}}_{[\alpha\beta]}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= -\mathrm{i} f^{\mathrm{T}}_{\rho}\int_0^1 \mathrm{d}t\,\frac{1-t^2}{t}\int_{-1}^{1}\mathrm{d}\xi\,\mathrm{i}\xi t\,\hat{\Psi}^{(3)}(\xi)m_\rho^2\tilde{x}_{[\alpha}\Big\{e^{(\lambda)}_{\beta]} + \mathrm{i}\xi t P_{\beta]}(e^{(\lambda)}\tilde{x})\Big\}\\
&\quad \times \mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)}, \qquad (3.2.28)\\
&= -\mathrm{i} f^{\mathrm{T}}_{\rho}\sum_{n=1}^{\infty}\frac{(\mathrm{i}\kappa(\tilde{x}P))^n}{n!}\Psi^{(3)}_n\frac{2}{n+2}\frac{m_\rho^2}{\tilde{x}P}\tilde{x}_{[\alpha}\Big(e^{(\lambda)}_{\beta]} + (n-1)P_{\beta]}\frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P}\Big),\\
\langle 0|M^{\mathrm{tw4}}_{[\alpha\beta]}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= -\mathrm{i} f^{\mathrm{T}}_{\rho}\int_0^1 \mathrm{d}t\,\frac{1-t}{t}\int_{-1}^{1}\mathrm{d}\xi\,(\mathrm{i}t\xi)^2\hat{\Psi}^{(4)}(\xi)\, m_\rho^2\tilde{x}_{[\alpha}\Big\{e^{(\lambda)}_{\beta]}(\tilde{x}P) - P_{\beta]}(e^{(\lambda)}\tilde{x})\Big\}\\
&\quad \times \mathrm{e}^{\mathrm{i}\xi t(\tilde{x}P)},\\
&= -\mathrm{i} f^{\mathrm{T}}_{\rho}\sum_{n=2}^{\infty}\frac{(\mathrm{i}(\tilde{x}P))^n}{n!}\Psi^{(4)}_n\frac{n-1}{n+1}\frac{m_\rho^2}{\tilde{x}P}\tilde{x}_{[\alpha}\Big(e^{(\lambda)}_{\beta]} - P_{\beta]}\frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P}\Big). \qquad (3.2.29)
\end{aligned}
$$

Again, the moments of distribution amplitudes of twist $\tau = 2, 3$ and 4 start with $n = 0, 1$ and 2, respectively. In addition, we remark that only those terms of the operator (3.1.18) contribute to the twist–3 distribution amplitude which result from the trace terms of (3.1.17).

The matrix element of the original skew tensor operator is obtained as

$$
\begin{aligned}
\langle 0|M_{[\alpha\beta]}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= \mathrm{i} f^{\mathrm{T}}_{\rho}\Big[2e^{(\lambda)}_{[\alpha}P_{\beta]}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\Psi}^{(2)}(\xi)e_0(\mathrm{i}\zeta\xi) \qquad (3.2.30)\\
&\quad + \tilde{x}_{[\alpha}P_{\beta]}\frac{m_\rho^2(e^{(\lambda)}\tilde{x})}{(\tilde{x}P)^2}\int_{-1}^{1}\mathrm{d}\xi\Big\{\hat{\Psi}^{(2)}(\xi)\big[e_0(\mathrm{i}\zeta\xi) + 2e_1(\mathrm{i}\zeta\xi) + 6e_2(\mathrm{i}\zeta\xi)\big]\\
&\qquad - \hat{\Psi}^{(3)}(\xi)\big[1 + 2e_0(\mathrm{i}\zeta\xi) + 6e_2(\mathrm{i}\zeta\xi)\big] + \hat{\Psi}^{(4)}(\xi)\big[1 + e_0(\mathrm{i}\zeta\xi) - 2e_1(\mathrm{i}\zeta\xi)\big]\Big\}\\
&\quad + \tilde{x}_{[\alpha}e^{(\lambda)}_{\beta]}\frac{m_\rho^2}{\tilde{x}P}\int_{-1}^{1}\mathrm{d}\xi\Big\{\hat{\Psi}^{(2)}(\xi)\big[e_0(\mathrm{i}\zeta\xi) - 2e_1(\mathrm{i}\zeta\xi) - 2e_2(\mathrm{i}\zeta\xi)\big]\\
&\qquad + \hat{\Psi}^{(3)}(\xi)\big[1 + 2e_2(\mathrm{i}\zeta\xi)\big] - \hat{\Psi}^{(4)}(\xi)\big[1 + e_0(\mathrm{i}\zeta\xi) - 2e_1(\mathrm{i}\zeta\xi)\big]\Big\}\Big].
\end{aligned}
$$

Now we finish the determination of the eight meson wave functions which result from the nonlocal light–cone quark operators of definite twist.

3.2.3 The relations between ρ-meson distribution amplitudes of dynamical and geometric twist

By means of operators of geometric twist and using the vectors p_α and $e^{(\lambda)}_{\perp\alpha}$, the chiral-even ρ-distributions with definite (geometric) twist are:

$$
\begin{aligned}
\langle 0|O_\alpha(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= f_\rho m_\rho \Big[p_\alpha \frac{e^{(\lambda)}\tilde{x}}{\tilde{x}P} \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Phi}^{(2)}(\xi) e_0(\mathrm{i}\zeta\xi) \\
&+ e^{(\lambda)}_{\perp\alpha} \int_{-1}^{1} \mathrm{d}\xi \Big\{ \hat{\Phi}^{(2)}(\xi) e_1(\mathrm{i}\zeta\xi) + \hat{\Phi}^{(3)}(\xi)[e_0(\mathrm{i}\zeta\xi) - e_1(\mathrm{i}\zeta\xi)] \Big\} \\
&- \frac{1}{2}\tilde{x}_\alpha \frac{e^{(\lambda)}\tilde{x}}{(\tilde{x}P)^2} m_\rho^2 \int_{-1}^{1} \mathrm{d}\xi \Big\{ \hat{\Phi}^{(4)}(\xi) \Big[e_0(\mathrm{i}\zeta\xi) - 3e_1(\mathrm{i}\zeta\xi) + 2\int_0^1 \mathrm{d}t\, e_1(\mathrm{i}\zeta\xi t) \Big] \\
&\qquad - \hat{\Phi}^{(2)}(\xi) \Big[e_1(\mathrm{i}\zeta\xi) - 2\int_0^1 \mathrm{d}t\, e_1(\mathrm{i}\zeta\xi t) \Big] + 4\hat{\Phi}^{(3)} \Big[e_1(\mathrm{i}\zeta\xi) - \int_0^1 \mathrm{d}t\, e_1(\mathrm{i}\zeta\xi t) \Big] \Big\} \Big]
\end{aligned}
\tag{3.2.31}
$$

and

$$
\langle 0|O_{5\alpha}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = \frac{1}{2}\Big(f_\rho - f_\rho^{\mathrm{T}} \frac{m_u + m_d}{m_\rho} \Big) m_\rho \epsilon_\alpha^{\ \beta\mu\nu} e^{(\lambda)}_{\perp\beta} p_\mu \tilde{x}_\nu \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Xi}^{(3)}(\xi) e_0(\mathrm{i}\zeta\xi),
\tag{3.2.32}
$$

while the chiral-odd distributions are defined as

$$
\begin{aligned}
\langle 0|M_{[\alpha\beta]}(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle &= \mathrm{i} f_\rho^{\mathrm{T}} \Big[2e^{(\lambda)}_{\perp[\alpha} p_{\beta]} \int_{-1}^{1} \mathrm{d}\xi \hat{\Psi}^{(2)}(\xi) e_0(\mathrm{i}\zeta\xi) \\
&+ 2\tilde{x}_{[\alpha} p_{\beta]} \frac{e^{(\lambda)}\tilde{x}}{(\tilde{x}P)^2} m_\rho^2 \int_{-1}^{1} \mathrm{d}\xi \Big\{ 2\hat{\Psi}^{(2)}(\xi) e_2(\mathrm{i}\zeta\xi) - \hat{\Psi}^{(3)}(\xi) \big[e_0(\mathrm{i}\zeta\xi) + 2e_2(\mathrm{i}\zeta\xi) \big] \Big\} \\
&- \tilde{x}_{[\alpha} e^{(\lambda)}_{\beta]\perp} \frac{m_\rho^2}{\tilde{x}P} \int_{-1}^{1} \mathrm{d}\xi \Big\{ 2\hat{\Psi}^{(2)}(\xi) \big[e_1(\mathrm{i}\zeta\xi) + e_2(\mathrm{i}\zeta\xi) \big] - \hat{\Psi}^{(3)}(\xi) \big[1 + 2e_2(\mathrm{i}\zeta\xi) \big] \\
&\qquad\qquad + \hat{\Psi}^{(4)}(\xi) \big[1 + e_0(\mathrm{i}\zeta\xi) - 2e_1(\mathrm{i}\zeta\xi) \big] \Big\} \Big]
\end{aligned}
\tag{3.2.33}
$$

and

$$
\langle 0|N(\tilde{x},-\tilde{x})|\rho(P,\lambda)\rangle = -\mathrm{i}\Big(f_\rho^{\mathrm{T}} - f_\rho \frac{m_u + m_d}{m_\rho} \Big) \big(e^{(\lambda)}\tilde{x} \big) m_\rho^2 \int_{-1}^{1} \mathrm{d}\xi\, \hat{\Upsilon}^{(3)}(\xi) e_0(\mathrm{i}\zeta\xi).
\tag{3.2.34}
$$

Comparing these expressions (3.2.31)–(3.2.34) with the meson distribution amplitudes of dynamical twist (3.2.5) – (3.2.8), we observe that it is necessary to re-express the truncated exponentials and perform appropriate variable transformations. After such manipulations we obtain the following relations, which allow to reveal the interrelations between the different twist definitions of meson distribution amplitudes:

$$\hat{\phi}_{\parallel}(\xi) \equiv \hat{\Phi}^{(2)}(\xi), \tag{3.2.35}$$

$$\hat{g}_{\perp}^{(v)}(\xi) = \hat{\Phi}^{(3)}(\xi) + \int_{\xi}^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau}\Big(\hat{\Phi}^{(2)} - \hat{\Phi}^{(3)}\Big)(\tau)\,, \tag{3.2.36}$$

$$\begin{aligned}\hat{g}_3(\xi) = \hat{\Phi}^{(4)}(\xi) - \int_{\xi}^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau}\Big\{ &\Big(\hat{\Phi}^{(2)} - 4\hat{\Phi}^{(3)} + 3\hat{\Phi}^{(4)}\Big)(\tau) \\ &+ 2\ln\Big(\frac{\xi}{\tau}\Big)\Big(\hat{\Phi}^{(2)} - 2\hat{\Phi}^{(3)} + \hat{\Phi}^{(4)}\Big)(\tau)\Big\},\end{aligned} \tag{3.2.37}$$

$$\hat{g}_{\perp}^{(a)}(\xi) \equiv \hat{\Xi}^{(3)}(\xi), \tag{3.2.38}$$

$$\hat{\phi}_{\perp}(\xi) \equiv \hat{\Psi}^{(2)}(\xi), \tag{3.2.39}$$

$$\hat{h}_{\parallel}^{(t)}(\xi) = \hat{\Psi}^{(3)}(\xi) + 2\xi \int_{\xi}^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau^2}\Big(\hat{\Psi}^{(2)} - \hat{\Psi}^{(3)}\Big)(\tau)\,, \tag{3.2.40}$$

$$\begin{aligned}\hat{h}_3(\xi) = \hat{\Psi}^{(4)}(\xi) + \int_{\xi}^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau}\Big\{ &2\Big(\hat{\Psi}^{(2)} - \hat{\Psi}^{(4)}\Big)(\tau) - 2\frac{\xi}{\tau}\Big(\hat{\Psi}^{(2)} - \hat{\Psi}^{(3)}\Big)(\tau) \\ &- \delta\Big(\frac{\xi}{\tau}\Big)\Big(\hat{\Psi}^{(3)} - \hat{\Psi}^{(4)}\Big)(\tau)\Big\},\end{aligned} \tag{3.2.41}$$

$$\hat{h}_{\parallel}^{(s)}(\xi) \equiv \hat{\Upsilon}^{(3)}(\xi). \tag{3.2.42}$$

Obviously, both decompositions coincide in the leading terms, e.g., the meson distribution amplitudes $\hat{\phi}_{\parallel}(\xi)$ and $\hat{\phi}_{\perp}(\xi)$ are genuine geometric twist-2 and $\hat{g}_{\perp}^{(a)}(\xi)$ and $\hat{h}_{\parallel}^{(s)}(\xi)$ are genuine geometric twist-3 functions. But the different twist definitions of meson distribution amplitudes differ at higher twist. For instance, the meson distribution amplitudes $\hat{g}_{\perp}^{(v)}(\xi)$ and $\hat{h}_{\parallel}^{(t)}(\xi)$ with dynamical twist $t = 3$ contain contributions with geometrical twist $\tau = 2$ and 3. Additionally, dynamical twist $t = 4$ meson distribution amplitudes $\hat{g}_3(\xi)$ and $\hat{h}_3(\xi)$ contain contributions with geometrical twist $\tau = 2$, 3 as well as 4.

The conventional distribution amplitudes can be written in terms of the new ρ-meson distribution amplitudes. The nontrivial relations are:

$$\hat{\Phi}^{(3)}(\xi) = \hat{g}_{\perp}^{(v)}(\xi) + \frac{1}{\xi}\int_{\xi}^{\mathrm{sign}(\xi)} \mathrm{d}\tau\big(\hat{g}_{\perp}^{(v)} - \hat{\phi}_{\parallel}\big)(\tau)\,, \tag{3.2.43}$$

$$\begin{aligned}\hat{\Phi}^{(4)}(\xi) = \hat{g}_3(\xi) &+ \frac{1}{\xi^2}\int_{\xi}^{\mathrm{sign}(\xi)} \mathrm{d}\tau\,\tau\big(3\hat{g}_3 - 4\hat{g}_{\perp}^{(v)} + \hat{\phi}_{\parallel}\big)(\tau) \\ &+ \frac{1}{\xi^2}\int_{\xi}^{\mathrm{sign}(\xi)} \mathrm{d}\tau\,\tau\Big(1 - \frac{\xi}{\tau}\Big)\big(\hat{g}_3 - 4\hat{g}_{\perp}^{(v)} + 3\hat{\phi}_{\parallel}\big)(\tau)\,,\end{aligned} \tag{3.2.44}$$

$$\hat{\Psi}^{(3)}(\xi) = \hat{h}_{\parallel}^{(t)}(\xi) + \frac{2}{\xi}\int_{\xi}^{\mathrm{sign}(\xi)} \mathrm{d}\tau\big(\hat{h}_{\parallel}^{(t)} - \hat{\phi}_{\perp}\big)(\tau)\,, \tag{3.2.45}$$

$$\hat{\Psi}^{(4)}(\xi) = \hat{h}_3(\xi) + \frac{2}{\xi^2}\int_{\xi}^{\mathrm{sign}(\xi)} \mathrm{d}\tau\,\tau\big(\hat{h}_3 - \hat{h}_{\parallel}^{(t)}\big)(\tau) - \frac{2}{\xi^2}\int_{\xi}^{\mathrm{sign}(\xi)} \mathrm{d}\tau\,\tau\Big(1 - \frac{\xi}{\tau}\Big)\big(\hat{h}_{\parallel}^{(t)} - \hat{\phi}_{\perp}\big)(\tau)\,. \tag{3.2.46}$$

The relation between the moments may be read off from Eqs. (3.2.31) – (3.2.34) as follows:

$$\phi_{\|n} \equiv \Phi_n^{(2)}, \tag{3.2.47}$$

$$g_{\perp n}^{(v)} = \Phi_n^{(3)} + \frac{1}{n+1}\Big(\Phi_n^{(2)} - \Phi_n^{(3)}\Big), \tag{3.2.48}$$

$$g_{3n} = \Phi_n^{(4)} - \frac{1}{n+1}\Big(\Phi_n^{(2)} - 4\Phi_n^{(3)} + 3\Phi_n^{(4)}\Big) + \frac{2}{(n+1)^2}\Big(\Phi_n^{(2)} - 2\Phi_n^{(3)} + \Phi_n^{(4)}\Big), \tag{3.2.49}$$

$$g_{\perp n}^{(a)} \equiv \Xi_n^{(3)}, \tag{3.2.50}$$

$$\phi_{\perp n} \equiv \Psi_n^{(2)}, \tag{3.2.51}$$

$$h_{\|n}^{(t)} = \Psi_n^{(3)} + \frac{2}{n+2}\Big(\Psi_n^{(2)} - \Psi_n^{(3)}\Big), \tag{3.2.52}$$

$$h_{3n} = \Psi_n^{(4)} + \frac{2}{n+1}\Big(\Psi_n^{(2)} - \Psi_n^{(4)}\Big) - \frac{2}{n+2}\Big(\Psi_n^{(2)} - \Psi_n^{(3)}\Big) - \delta_{n0}\Big(\Psi_n^{(3)} - \Psi_n^{(4)}\Big), \tag{3.2.53}$$

$$h_{\|n}^{(s)} \equiv \Upsilon_n^{(3)}. \tag{3.2.54}$$

In terms of the moments the relations between old and new wave functions may be easily inverted; for the distribution amplitudes itself the expression of the new distribution amplitudes through the old ones is more involved. The inverse relations are:

$$\Phi_n^{(3)} = g_{\perp n}^{(v)} + \frac{1}{n}\Big(g_{\perp n}^{(v)} - \phi_{\|n}\Big), \qquad n > 0 \tag{3.2.55}$$

$$\Phi_n^{(4)} = g_{3n} + \frac{1}{n-1}\Big(3g_{3n} - 4g_{\perp n}^{(v)} + \phi_{\|n}\Big) + \frac{1}{n(n-1)}\Big(g_{3n} - 4g_{\perp n}^{(v)} + 3\phi_{\|n}\Big), \qquad n > 1 \tag{3.2.56}$$

$$\Psi_n^{(3)} = h_{\|n}^{(t)} + \frac{2}{n}\Big(h_{\|n}^{(t)} - \phi_{\perp n}\Big), \qquad n > 0 \tag{3.2.57}$$

$$\Psi_n^{(4)} = h_{3n} + \frac{2}{n-1}\Big(h_{3n} - h_{\|n}^{(t)}\Big) - \frac{2}{n(n-1)}\Big(h_{\|n}^{(t)} - \phi_{\perp n}\Big), \qquad n > 1. \tag{3.2.58}$$

The relations between the dynamical and geometrical twist meson DAs show that the dynamical distribution amplitudes can be used to determine the new geometrical ones and vice versa. Obviously, the same holds for their moments. In principle, this allows to determine, e.g., the new distribution amplitudes from the experimental data if these are known for the conventional ones. Thus, all of the physical and experimental correlations obtained by Ball *et al.* [13, 14, 15], e.g., the asymptotic wave functions, can be used for the geometrical twist distribution amplitudes.

3.2.4 Wandzura-Wilczek type relations

This Section is devoted to the general discussion of geometric Wandzura-Wilczek-type relations between the conventional distribution amplitudes which we have obtained by means of the distribution amplitudes with a definite geometric twist. These geometric Wandzura-Wilczek-type relations are independent from the QCD field equations and the corresponding operator relations. The mismatch between the definition of dynamical and geometric twist gives rise to relations of geometric Wandzura-Wilczek-type which show that the dynamical twist functions contain various parts of different geometric twist. Consequently, a genuine geometric twist distribution amplitude do not give rise to any geometric Wandzura-Wilczek-type relation. Thereby, we obtain new (geometric) Wandzura-Wilczek-type relations and sum rules for the meson distribution amplitudes of dynamical twist.

We use the notations $\hat g_\perp^{(v)}(\xi) = \hat g_\perp^{(v),\rm tw2}(\xi) + \hat g_\perp^{(v),\rm tw3}(\xi)$, where $\hat g_\perp^{(v),\rm tw2}(\xi)$ is the genuine twist-2 and $\hat g_\perp^{(v),\rm tw3}(\xi)$ the genuine twist-3 part of $\hat g_\perp^{(v)}(\xi)$. Substituting (3.2.35) into (3.2.36), we get

$$\hat g_\perp^{(v),\rm tw2}(\xi) = \int_\xi^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau}\, \hat\phi_\parallel(\tau), \tag{3.2.59}$$

$$\hat g_\perp^{(v),\rm tw3}(\xi) = \hat g_\perp^{(v)}(\xi) - \int_\xi^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau}\, \hat\phi_\parallel(\tau), \tag{3.2.60}$$

where (3.2.59) is really the analogue of the Wandzura-Wilczek relation for the twist-2 part [123] because they argued that the geometric twist-3 contribution should be small. On the other hand, Eq. (3.2.60) is the Wandzura-Wilczek-type relation of geometric twist-3 and gives the information how much is the physical contribution of the geometric twist-3 operator in this non-forward process. These relations show that the transverse distribution $\hat g_\perp^{(v)}(\xi)$ is related to the longitudinal distribution $\hat\phi_\parallel(\xi)$. Obviously, this analogy between the Wandzura-Wilczek relations of parton distribution amplitudes and of meson distribution amplitudes is not surprising because the operator structures are the same and the ρ-meson polarization vector formally substitutes the nucleon spin vector in the Lorentz structures. The generalization of the geometric Wandzura-Wilczek relation (3.2.59) for non-forward distribution amplitudes was discussed in Ref. [28]. Let me note that the Wandzura-Wilczek relations which are obtained in Refs. [3, 13, 14] for $g_\perp^{(v),\rm tw2}(u)$ are significantly different (see also [17]). The reason is that they used equations of motion operator relations in order to isolate the dynamical twist-3 part. Therefore, their Wandzura-Wilczek relations are dynamical relations (dynamical Wandzura-Wilczek-type relations). The corresponding geometric Wandzura-Wilczek-type relations for the moments are:

$$g_{\perp n}^{(v),\rm tw2} = \frac{1}{n+1}\, \phi_{\parallel n}, \tag{3.2.61}$$

$$g_{\perp n}^{(v),\rm tw3} = g_{\perp n}^{(v)} - \frac{1}{n+1}\, \phi_{\parallel n}, \qquad n > 0. \tag{3.2.62}$$

Note that Eq. (3.2.61) was earlier obtained in Ref. [13]. The reason for this agreement between the twist-2 part of the dynamical and geometric Wandzura-Wilczek relation is that the dynamical and geometric twist-2 meson distribution amplitude coincides. Obviously, from the relation (3.2.61) for $n = 0$ the condition of the normalization and for $n = 1$ the following sum rule follow:

$$\int_{-1}^{1} \mathrm{d}\xi\, \hat g_\perp^{(v)}(\xi) = \int_{-1}^{1} \mathrm{d}\xi\, \hat\phi_\parallel(\xi) = 1, \qquad \int_{-1}^{1} \mathrm{d}\xi\, \xi \hat g_\perp^{(v),\rm tw2}(\xi) = \frac{1}{2}\int_{-1}^{1} \mathrm{d}\xi\, \xi\hat\phi_\parallel(\xi). \tag{3.2.63}$$

Using the formulas (3.2.35), (3.2.36), (3.2.37) and (3.2.43), we obtain the integral relations for the function $\hat g_3(\xi) = \hat g_3^{\rm tw2}(\xi) + \hat g_3^{\rm tw3}(\xi) + \hat g_3^{\rm tw4}(\xi)$ as

$$\hat g_3^{\rm tw2}(\xi) = -\int_\xi^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau} \Big\{ \hat\phi_\parallel(\tau) + 2\ln\Big(\frac{\xi}{\tau}\Big)\hat\phi_\parallel(\tau)\Big\}, \tag{3.2.64}$$

$$\hat g_3^{\rm tw3}(\xi) = 4\int_\xi^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau} \Big\{ \hat g_\perp^{(v)}(\tau) + \ln\Big(\frac{\xi}{\tau}\Big)\hat\phi_\parallel(\tau)\Big\}, \tag{3.2.65}$$

$$\hat g_3^{\rm tw4}(\xi) = \hat g_3(\xi) - \int_\xi^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau} \Big\{ \big(4\hat g_\perp^{(v)} - \hat\phi_\parallel\big)(\tau) + 2\ln\Big(\frac{\xi}{\tau}\Big)\hat\phi_\parallel(\tau)\Big\}. \tag{3.2.66}$$

Due to the fact that $\hat g_3(\xi)$ contains twist-2, twist-3 as well as twist-4, we have obtained three integral relations. For example, Eq. (3.2.64) demonstrates that the twist-2 part $\hat g_3^{\rm tw2}(\xi)$ can be expressed in terms of the twist-2 function $\hat{\phi}_{\|}(\xi)$. Additionally, the twist-3 part $\hat{g}_3^{\mathrm{tw3}}(\xi)$ is given in Eq. (3.2.65) in terms of the functions $\hat{\phi}_{\|}(\xi)$ and $\hat{g}_{\perp}^{(v)}(\xi)$. Eq. (3.2.66) expresses how can be obtained the twist-4 part $\hat{g}_3^{\mathrm{tw4}}(\xi)$ from $\hat{g}_3(\xi)$. For the distribution amplitude moments we obtain:

$$g_{3n}^{\mathrm{tw2}} = -\frac{1}{n+1}\,\phi_{\|n} + \frac{2}{(n+1)^2}\,\phi_{\|n}, \tag{3.2.67}$$

$$g_{3n}^{\mathrm{tw3}} = \frac{4}{n+1}\,g_{\perp n}^{(v)} - \frac{4}{(n+1)^2}\,\phi_{\|n}, \qquad n>0 \tag{3.2.68}$$

$$g_{3n}^{\mathrm{tw4}} = g_{3n} - \frac{1}{n+1}\Big(4g_{\perp n}^{(v)} - \phi_{\|n}\Big) + \frac{2}{(n+1)^2}\,\phi_{\|n}, \qquad n>1. \tag{3.2.69}$$

For $n=1$ in (3.2.67) and (3.2.68) we find the sum rules

$$\int_{-1}^{1} \mathrm{d}\xi\,\xi\hat{g}_3^{\mathrm{tw2}}(\xi) = 0, \tag{3.2.70}$$

$$\int_{-1}^{1} \mathrm{d}\xi\,\xi\hat{g}_3^{\mathrm{tw3}}(\xi) = \int_{-1}^{1} \mathrm{d}\xi\,\xi\big(2\hat{g}_{\perp}^{(v)} - \hat{\phi}_{\|}\big)(\xi), \tag{3.2.71}$$

which give the interesting sum rule for the dynamical twist-4 function $\hat{g}_3(\xi)$:

$$\int_{-1}^{1} \mathrm{d}\xi\,\xi\hat{g}_3(\xi) = \int_{-1}^{1} \mathrm{d}\xi\,\xi\big(2\hat{g}_{\perp}^{(v)} - \hat{\phi}_{\|}\big)(\xi). \tag{3.2.72}$$

Let us now discuss the geometric Wandzura-Wilczek-type relations for the chiral-odd meson distribution amplitudes. If we substitute (3.2.39) into (3.2.40) and using $\hat{h}_{\|}^{(t)}(\xi) = \hat{h}_{\|}^{(t),\mathrm{tw2}}(\xi) + \hat{h}_{\|}^{(t),\mathrm{tw3}}(\xi)$ we derive

$$\hat{h}_{\|}^{(t),\mathrm{tw2}}(\xi) = 2\xi\int_{\xi}^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau^2}\,\hat{\phi}_{\perp}(\tau), \tag{3.2.73}$$

$$\hat{h}_{\|}^{(t),\mathrm{tw3}}(\xi) = \hat{h}_{\|}^{(t)}(\xi) - 2\xi\int_{\xi}^{\mathrm{sign}(\xi)} \frac{\mathrm{d}\tau}{\tau^2}\,\hat{\phi}_{\perp}(\tau), \tag{3.2.74}$$

where (3.2.73) is the analogue of the twist-2 Wandzura-Wilczek-type relation of the nucleon structure function $h_L(z)$ which was obtained by Jaffe and Ji [79]. These Wandzura Wilczek-type relations point out that the longitudinal distribution $\hat{h}_{\|}^{(t)}(\xi)$ is related to the transverse distribution $\hat{\phi}_{\perp}(\xi)$. Dynamical Wandzura-Wilczek relations were obtained in Ref. [14]. Obviously, (3.2.74) is the corresponding twist-3 relation. The relations for the moments read:

$$h_{\|n}^{(t),\mathrm{tw2}} = \frac{2}{n+2}\,\phi_{\perp n}, \tag{3.2.75}$$

$$h_{\|n}^{(t),\mathrm{tw3}} = h_{\|n}^{(t)} - \frac{2}{n+2}\,\phi_{\perp n}, \qquad n>0. \tag{3.2.76}$$

For $n=1$ in (3.2.75) we observe the following sum rule

$$\int_{-1}^{1} \mathrm{d}\xi\,\xi\hat{h}_{\|}^{(t),\mathrm{tw2}}(\xi) = \frac{2}{3}\int_{-1}^{1} \mathrm{d}\xi\,\xi\hat{\phi}_{\perp}(\xi). \tag{3.2.77}$$

Now, using the formulas (3.2.39), (3.2.40), (3.2.41) and (3.2.45), we obtain the integral relations for the function with $\hat{h}_3(\xi) = \hat{h}_3^{\rm tw2}(\xi) + \hat{h}_3^{\rm tw3}(\xi) + \hat{h}_3^{\rm tw4}(\xi)$ as

$$\hat{h}_3^{\rm tw2}(\xi) = \frac{2}{\xi}\int_\xi^{{\rm sign}(\xi)} {\rm d}\tau \Big(\frac{\xi}{\tau} - \frac{\xi^2}{\tau^2}\Big)\hat{\phi}_\perp(\tau), \tag{3.2.78}$$

$$\hat{h}_3^{\rm tw3}(\xi) = \frac{2}{\xi}\int_\xi^{{\rm sign}(\xi)} {\rm d}\tau \Big\{\hat{h}_\parallel^{(t)}(\tau) - \Big(1 - \frac{\xi^2}{\tau^2}\Big)\hat{\phi}_\perp(\tau)\Big\}, \tag{3.2.79}$$

$$\hat{h}_3^{\rm tw4}(\xi) = \hat{h}_3(\xi) - \frac{2}{\xi}\int_\xi^{{\rm sign}(\xi)} {\rm d}\tau \Big\{\hat{h}_\parallel^{(t)}(\tau) - \Big(1 - \frac{\xi}{\tau}\Big)\hat{\phi}_\perp(\tau)\Big\}. \tag{3.2.80}$$

and for the moments

$$h_{3n}^{\rm tw2} = \frac{2}{n+1}\,\phi_{\perp n} - \frac{2}{n+2}\,\phi_{\perp n}, \tag{3.2.81}$$

$$h_{3n}^{\rm tw3} = \frac{2}{n}\,h_{\parallel n}^{(t)} - \frac{4}{(n+2)n}\,\phi_{\perp n}, \qquad n > 0 \tag{3.2.82}$$

$$h_{3n}^{\rm tw4} = h_{3n} - \frac{2}{n}\,h_{\parallel n}^{(t)} + \frac{2}{(n+1)n}\,\phi_{\perp n}, \qquad n > 1. \tag{3.2.83}$$

Thus, Eq. (3.2.78) means that the twist-2 part $\hat{h}_3^{\rm tw2}(\xi)$ can be expressed in terms of the twist-2 function $\hat{\phi}_\perp(\xi)$. Additionally, the twist-3 part $\hat{h}_3^{\rm tw3}(\xi)$ is given in Eq. (3.2.79) in terms of the functions $\hat{\phi}_\perp(\xi)$ and $\hat{h}_\parallel^{(t)}(\xi)$. Eq. (3.2.80) gives the subtraction rule in order to obtain the twist-4 part $\hat{h}_3^{\rm tw4}(\xi)$ from the original function $\hat{h}_3(\xi)$. Obviously, the same is valid for the corresponding distribution amplitude moments. For $n = 1$ in (3.2.81) and (3.2.82) we observe the following sum rules

$$\int_{-1}^1 {\rm d}\xi\,\xi\hat{h}_3^{\rm tw2}(\xi) = \frac{1}{3}\int_{-1}^1 {\rm d}\xi\,\xi\hat{\phi}_\perp(\xi), \tag{3.2.84}$$

$$\int_{-1}^1 {\rm d}\xi\,\xi\hat{h}_3^{\rm tw3}(\xi) = \frac{1}{3}\int_{-1}^1 {\rm d}\xi\,\xi\big(6\hat{h}_\parallel^{(t)}(\xi) - 4\hat{\phi}_\perp(\xi)\big), \tag{3.2.85}$$

which give the sum rule for the dynamical twist-4 function $\hat{h}_3(\xi)$:

$$\int_{-1}^1 {\rm d}\xi\,\xi\hat{h}_3(\xi) = \int_{-1}^1 {\rm d}\xi\,\xi\big(2\hat{h}_\parallel^{(t)}(\xi) - \hat{\phi}_\perp(\xi)\big). \tag{3.2.86}$$

Because the functions $\hat{g}_\perp^{(a)}(\xi)$ and $\hat{h}_\parallel^{(s)}(\xi)$ are pure geometric as well as dynamical twist-3 functions, there is no mismatch between the dynamical and geometric twist and no geometric Wandzura-Wilczek-type relation occurs. However, dynamical Wandzura-Wilczek relations for $\hat{g}_\perp^{(a)}(\xi)$ and $\hat{h}_\parallel^{(s)}(\xi)$ were obtained by Ball *et al.* [13, 14] by using the QCD equations of motion. Here I am discussing the relation for $g_{\perp n}^{(a)}$. Neglecting quark masses and three-particle quark-antiquark-gluon operators of twist-3 one recovers the operator relation (see [9])

$$\big[\bar{u}(\tilde{x})\gamma_\alpha\gamma_5 d(-\tilde{x})\big]^{\rm tw3} = {\rm i}\epsilon_\alpha^{\ \beta\mu\nu}\tilde{x}_\beta \int_0^1 {\rm d}u\,u\,\hat{\partial}_\mu\big[\bar{u}(u\tilde{x})\gamma_\nu d(-u\tilde{x})\big] + \dots\,, \tag{3.2.87}$$

where $\hat{\partial}_\mu$ is the so-called total derivative which translates the expansion point y:

$$\hat{\partial}_\mu\big[\bar{u}(u\tilde{x})\gamma_\nu d(-u\tilde{x})\big] \equiv \frac{\partial}{\partial y^\mu}\big[\bar{u}(y+u\tilde{x})\gamma_\nu d(y-u\tilde{x})\big]\big|_{y\to 0}. \tag{3.2.88}$$

The total derivative on the r.h.s. of Eq. (3.2.87) is calculated as

$$\hat{\partial}_\mu\big[\bar{u}(u\tilde{x})\gamma_\nu d(-u\tilde{x})\big] = \bar{u}(u\tilde{x})\big(\overleftarrow{D}_\mu + \overrightarrow{D}_\mu\big)\gamma_\nu d(-u\tilde{x}) - \mathrm{i}\int_{-u}^{u}\mathrm{d}v\, \bar{u}(u\tilde{x})\tilde{x}^\rho F_{\rho\mu}(v\tilde{x})\gamma_\nu d(-u\tilde{x}), \tag{3.2.89}$$

where $F_{\mu\nu}$ is the gluon field strength. Obviously, the second operator on the r.h.s. of Eq. (3.2.89) is a Shuryak-Vainshtein type operator having minimal twist-3. The other operator on the r.h.s. of Eq. (3.2.89) has, in general, twist-2, 3 and 4. But, after multiplying by the Levi-Civita tensor $\epsilon_{\alpha\beta\mu\nu}$ only the twist-3 part survives and gives contributions which are not small. Therefore, the operator identity (3.2.87) is a genuine twist-3 relation. The twist-3 matrix element of the first operator on the r.h.s. of Eq. (3.2.89) is given by

$$\mathrm{i}\epsilon_\alpha{}^{\beta\mu\nu}\tilde{x}_\beta\langle 0|\bar{u}(\tilde{x})\big(\overleftarrow{D}_\mu + \overrightarrow{D}_\mu\big)\gamma_\nu d(-\tilde{x})|\rho(P,\lambda)\rangle = f_\rho m_\rho \epsilon_\alpha{}^{\beta\mu\nu} e^{(\lambda)}_{\perp\beta} p_\mu \tilde{x}_\nu \int_{-1}^{1}\mathrm{d}\xi\, \hat{\Omega}^{(3)}(\xi) e_0(\mathrm{i}\zeta\xi), \tag{3.2.90}$$

where $\hat{\Omega}^{(3)}(\xi)$ is the corresponding meson distribution amplitude of twist-3.

Taking the matrix element of the operators on both sides of Eq. (3.2.87), using Eqs. (3.2.32), (3.2.38), and (3.2.90), and neglecting quark masses and trilocal operators, one obtains a geometric relation between the meson distribution amplitudes $\hat{g}^{(a)}_\perp(\xi)$ and $\hat{\Omega}^{(3)}(\xi)$ imposed by the QCD equations of motion as

$$\hat{g}^{(a)}_\perp(\xi) = 2\xi\int_\xi^{\mathrm{sign}(\xi)}\frac{\mathrm{d}\tau}{\tau^2}\,\hat{\Omega}^{(3)}(\tau), \tag{3.2.91}$$

and for the meson distribution amplitude moments

$$g^{(a)}_{\perp n} = \frac{2}{n+2}\,\Omega^{(3)}_n, \tag{3.2.92}$$

where $\Omega^{(3)}_n$ are the geometric twist-3 distribution amplitude moments. This relation (3.2.92) reproduces the fact that the moments $g^{(a)}_{\perp n}$ have genuine geometric twist-3 (see Eq. (3.2.50)) and no geometric twist-2 contributions from the total derivative come into the game.

But this is not what Ball *et al.* [13, 14] did. They have calculated the r.h.s. of Eq. (3.2.87) by means of

$$\hat{\partial}_\mu\langle 0|\bar{u}(\tilde{x})\gamma_\nu d(-\tilde{x})|\rho(P,\lambda)\rangle \tag{3.2.93}$$

and Eq. (3.2.5). Eventually, they obtained the dynamical Wandzura-Wilczek-type relation [13]

$$\hat{g}^{(a)}_\perp(\xi) = 2\xi\int_\xi^{\mathrm{sign}(\xi)}\frac{\mathrm{d}\tau}{\tau^2}\,\hat{g}^{(v)}_\perp(\tau), \tag{3.2.94}$$

and for the moments

$$g^{(a)}_{\perp n} = \frac{2}{n+2}\,g^{(v)}_{\perp n}. \tag{3.2.95}$$

Obviously, Eqs. (3.2.94) and (3.2.95) are relations between equal dynamical twist. But, the l.h.s. of these relations has geometric twist-3 and the r.h.s. has geometric twist-2 and twist-3. Therefore, a dynamical Wandzura-Wilczek relation is a relation which is derived by means of the

QCD equations of motion in contrast to geometric one. Note that the Wandzura-Wilczek-like relations used in Refs. [4, 22, 87, 115, 5] are dynamical ones.

Similar arguments are valid for the other QCD equations of motion operator relations used in Ref. [14]. Thus, from the group theoretical point of view, it is a bit misleading to claim that an additional geometric twist-2 contribution is produced by geometric twist-3 operators with a total derivative and that the QCD equations of motion give a relation between distribution amplitudes of different geometric twist in exclusive processes and off-forward scattering. From this point of view, the geometric twist is a proper concept for classifying non-forward inclusive or general exclusive matrix-elements.

3.2.5 Conclusions

We discussed the model-independent classification of meson distribution amplitudes with respect to geometric twist. We gave the relations between the geometric twist and Ball and Braun's dynamical twist distribution amplitudes. These relations demonstrate the interrelations between the different twist definitions. Consequently, these relations are "transformation rules" between the dynamical and geometric twist distribution amplitudes.

Another results of this Section are geometric Wandzura-Wilczek-type relations between the dynamical twist distributions which we have obtained by means of our "transformation rules". The reason of these geometric Wandzura-Wilczek-type relations is the mismatch of the dynamical twist with respect to geometric twist. Because we have not used the QCD equations of motion, the geometric Wandzura-Wilczek-type relations are based on a no-dynamical level and are model-independent. Additionally, I have discussed the difference between dynamical and geometric Wandzura-Wilczek-type relations.

3.3 Power corrections of non-forward quark distributions and harmonic operators with definite geometric twist

In this Section we develop a formalism for the resummation of the target mass corrections for off-forward processes by means of harmonic off-cone operators of geometric twist. The harmonic operators of a given spin or twist have a fixed tensor structure and their matrix elements have an explicit target mass dependence, determined by the condition that the (local) operators have to be traceless.

We determine the twist decomposition of the following chiral-even (axial) vector operators, $O_\alpha(\kappa x, -\kappa x)$, and $O_{5\alpha}(\kappa x, -\kappa x)$, together with the corresponding (pseudo) scalar operators, $O_{(5)}(\kappa x, -\kappa x)$, and the chiral-odd scalar and skew tensor operator, $N(\kappa x, -\kappa x)$, and $M_{[\alpha\beta]}(\kappa x, -\kappa x)$ together with the vector and scalar operators, $M_\alpha(\kappa x, -\kappa x) = x^\beta M_{[\alpha\beta]}(\kappa x, -\kappa x)$, $M(\kappa x, -\kappa x) = x^\beta \partial^\alpha M_{[\alpha\beta]}(\kappa x, -\kappa x)$. The operators constitute a basis not only for the (usual) parton distributions as well as scalar and vector meson distribution amplitudes [66, 97, 98] but also for the consideration of double distribution amplitudes being relevant for the various light-cone dominated QCD processes under consideration. Here, their twist decomposition off the light-cone will be given up to twist 3 and, in the case of scalar operators, also for any twist. In fact, the 'external' operation of contracting with x^α or $x^\beta \partial^\alpha$ also influences the possible symmetry type of these operators and, therefore, of their twist decomposition.

These off-cone operators of definite twist are given for the local as well as the resummed nonlocal operators. The local operators given as the n-th Taylor coefficients of the non-local ones, are represented by (a finite series of) Gegenbauer polynomials $C_n^\nu(z)$, $\nu \geq 1$. The nonlocal operators, being obtained by resummation with respect to n, are represented by (a related series of) Bessel functions or, more exactly, either $J_{\nu-\frac{1}{2}}(z)$ or $I_{\nu-\frac{1}{2}}(z)$ depending on the values of their arguments.

The group theoretical method for the determination of target mass corrections in unpolarized deep inelastic scattering using harmonic scalar operators of definite spin and the corresponding matrix elements in terms of Gegenbauer polynomials has been used for the first time by Nachtmann [107] and later by Baluni and Eichten [18]. Using the same procedure, the target mass contributions for polarized deep inelastic scattering were studied in Refs. [124, 104, 86]. A key ingredient in this framework was that they obtained so-called "Nachtmann moments" for the unpolarized as well as polarized structure functions. A short review of Nachtmann's method in deep inelastic lepton-hadron scattering was given in Ref. [59]. All these investigations are based on the notion of geometric twist. A different method is due to Georgi and Politzer [58] for the target mass corrections in unpolarized deep inelastic scattering. This method has been applied also to the study of target mass corrections of polarized structure functions [110, 26]. In non-forward processes, the structure of mass corrections is more complicated than for deep-inelastic lepton-hadron scattering [16, 23].

Here, we generalize Nachtmann's procedure for the case of non-forward matrix elements applying it to off-cone quark-antiquark operators. Additionally, I extend the analysis of ρ-meson as well as pion DAs to include all meson-mass terms of harmonic twist-2 and twist-3 operators. Vector meson mass corrections of order x^2 were discussed by Ball and Braun [15, 16]. A resummed meson mass correction was just given in the case of a scalar theory by Ball [16] (see also Ref. [11]).

We calculate the operators and the matrix elements in the x-space. The corresponding matrix elements in the q-space which are needed to calculate the mass corrections in the Compton amplitude are given if we carry out the twist decomposition in the q-space.

3.3.1 Non-forward matrix elements of harmonic operators: The method

Parametrization of non-forward matrix elements by independent double distributions of definite twist

Now we consider the bilocal off-cone quark-antiquark operators i.e., operators *without* external operations which, generically, will be denoted by $\mathcal{O}_\Gamma(\kappa x, -\kappa x)$.

We will define the (off-cone) non-forward quark distributions of definite geometric twist as straightforward generalizations of the DAs discussed in Section 3.1 and 3.2 by writing the matrix elements of bilocal operators of definite geometric twist. Before discussing double distributions of geometric twist let us consider the most general parametrization of non-forward (off-cone) matrix elements:

$$\langle P_2, S_2|\mathcal{O}_\Gamma(\kappa x, -\kappa x)|P_1, S_1\rangle = \mathcal{K}^a_\Gamma(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, \mathrm{e}^{\mathrm{i}\kappa(x\mathbb{P})\mathbb{Z}}\, f_a(\mathbb{Z}, \mathbb{P}_i\mathbb{P}_j, x^2; \mu^2). \tag{3.3.1}$$

We use the notation $\mathbb{P} = \{P_+, P_-\}$ and $\mathbb{Z} = \{z_+, z_-\}$ with $P_\pm = P_2 \pm P_1$ and $z_\pm = \frac{1}{2}(z_2 \pm z_1)$ thereby defining some (2-dimensional) vector space with scalar product $\mathbb{P}\mathbb{Z} \equiv \sum P_i z_i = P_+ z_+ + P_- z_-$. In addition, the integration measure is defined by $\mathrm{D}\mathbb{Z} = dz_1 dz_2 \theta(1 - z_1)\theta(z_1 + 1)\theta(1 - z_2)\theta(z_2 + 1)$. The support restriction for the double distributions reads: $-1 \le z_i \le 1$. Here, $\mathcal{K}^a_\Gamma(\mathbb{P})$ denote the linear independent spin structures being defined by the help of the (free) hadron wave functions and governed by the $\Gamma-$structure of the corresponding nonlocal operator $\mathcal{O}_\Gamma$. For example, in the case of the virtual Compton scattering, there are two independent spin structures, the Dirac and the Pauli structure, $\mathcal{K}^1_\mu = \bar{U}(P_2, S_2)\gamma_\mu U(P_1, S_1)$ and

$\mathcal{K}^2_\mu = \bar{U}(P_2,S_2)\sigma_{\mu\nu}P^\nu_- U(P_1,S_1)/M$, respectively. Here $f_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2)$ are (renormalized) two-variable distribution amplitudes which have no definite geometric twist. The distributions are entire analytic functions with respect to $(x\mathbb{P}_i)$. In addition these distributions are, in principle, the (off-cone) generalizations of the distributions given in Ref. [106, 83].

Finally, let us comment on operators *with* external operations, i.e. when the (axial) vectors or the (skew) tensors are multiplied 'externally' by x^β and/or ∂^α. In these cases the 'external' vector x^β or tensor $x^\beta\partial^\alpha$ is assumed *not* to be Fourier transformed and the external operations have to be applied onto both sides of Eq. (3.3.1). Thereby the tensor structure of these external operations matches with the tensor structure of $\mathcal{K}^a_\Gamma$ which by itself are independent of the coordinates. Despite of this the external operations heavily influence the possible symmetry type of the local operators and their decomposition into irreducible tensor representations of the Lorentz group.

Now, let us take into account that the non-local operators $\mathcal{O}_\Gamma(\kappa x,-\kappa x)$, formally, are given by infinite series of operators of growing (geometric) twist τ, Eq. (2.2.22). By substituting the decomposition (2.2.22) into the representation (3.3.1) we get an analogous decomposition of the distributions $f_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2)$:

$$\mathcal{K}^a_\Gamma(\mathbb{P})f_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2) = \sum_{\tau=\tau_{\min}}^{\infty} c^{(\tau)\Gamma'}_{\Gamma}(x)\,\mathcal{K}^a_{\Gamma'}(\mathbb{P})f^{(\tau)}_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2), \tag{3.3.2}$$

where $f^{(\tau)}_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2)$ are distributions of geometric twist which still contain kinematical factors, like P^2_-. On the other side, the non-forward matrix elements of operators with geometric twist τ read:

$$\begin{aligned}\langle P_2,S_2|\mathcal{O}^{(\tau)}_\Gamma(\kappa x,-\kappa x)|P_1,S_1\rangle &= \mathcal{K}^a_\Gamma(\mathbb{P})\int \mathrm{D}\mathbb{Z}\,\mathrm{e}^{\mathrm{i}\kappa(x\mathbb{P})\mathbb{Z}} f^{(\tau)}_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2)\\ &= \mathcal{P}^{(\tau)\Gamma'}_{\Gamma}(x,\partial_x)\,\mathcal{K}^a_{\Gamma'}(\mathbb{P})\int \mathrm{D}\mathbb{Z}\,\mathrm{e}^{\mathrm{i}\kappa(x\mathbb{P})\mathbb{Z}}\; f^{(\tau)}_a(\mathbb{Z};\mu^2).\end{aligned} \tag{3.3.3}$$

In the second line of Eq. (3.3.3) we used the explicite twist decomposition by the help of orthogonal projectors (see Eq. (2.2.20). In this way one is able to introduce distributions of geometric twist $f^{(\tau)}_a(\mathbb{Z};\mu^2)$ which are independent form kinematical factors. Thus $f^{(\tau)}_a(\mathbb{Z};\mu^2)$ are already determined by the decomposition of the non-local matrix elements on the light-cone. Also their renormalization properties are only determined by the light-cone operators. These distribution amplitudes $f^{(\tau)}_a(\mathbb{Z};\mu^2)$ depend from the distribution parameters z_1 and z_2, only. Therefore, the power corrections of the distribution amplitudes $f^{(\tau)}_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2)$ are related to the distributions $f^{(\tau)}_a(\mathbb{Z};\mu^2)$ according to:

$$f^{(\tau)}_a(\mathbb{Z},\mathbb{P}_i\mathbb{P}_j,x^2;\mu^2) = \mathcal{F}^{(\tau)a'}_a\big((x\mathbb{P}\mathbb{Z}),x^2(\mathbb{P}\mathbb{Z})^2\big)f^{(\tau)}_{a'}(\mathbb{Z};\mu^2), \tag{3.3.4}$$

where $\mathcal{F}^{(\tau)a'}_a\big(((x\mathbb{P}\mathbb{Z}),x^2(\mathbb{P}\mathbb{Z})^2\big)$ is uniquely related to the twist decomposition of the off-cone operators and contains the information about the power corrections of the distributions amplitudes.

Let us now present the arguments leading to the r.h.s. of Eq. (3.3.3) more explicitly. Namely, the non-forward matrix elements after performing the twist projection of relations (2.2.14) are given as follows:

$$
\begin{aligned}
\langle P_2, S_2|\mathcal{O}^{(\tau)}_{\Gamma}(\kappa x, -\kappa x)|P_1, S_1\rangle &= \langle P_2, S_2|\mathcal{P}^{(\tau)\Gamma'}_{\Gamma}(x, \partial_x) \int \mathrm{d}^4 p\, \mathcal{O}_{\Gamma'}(p)\, \mathrm{e}^{\mathrm{i}\kappa xp}|P_1, S_1\rangle \\
&= \int \mathrm{d}^4 p \left(\mathcal{P}^{(\tau)\Gamma'}_{\Gamma}(x, \partial_x)\, \mathrm{e}^{\mathrm{i}\kappa xp}\right)\langle P_2, S_2|\mathcal{O}_{\Gamma'}(p)|P_1, S_1\rangle \qquad (3.3.5) \\
&= \int \mathrm{D}\mathbb{Z} \Big(\mathcal{P}^{(\tau)\Gamma'}_{\Gamma}(x, \partial_x)\, \mathrm{e}^{\mathrm{i}\kappa(x\mathbb{P})\mathbb{Z}}\Big)\, \mathcal{K}^{a}_{\Gamma'}(\mathbb{P})\, f^{(\tau)}_{a}(\mathbb{Z}; \mu^2).
\end{aligned}
$$

Here, in the second line we simply reordered the operations of integration, taking matrix elements and twist projections appropriately. In order to get the third line we observe the symmetry in x and p of the local twist projections, $\mathcal{P}^{(\tau)\Gamma'}_{n\ \Gamma}(x, \partial_x)(xp)^n \equiv \mathcal{P}^{(\tau)\Gamma'}_{n\ \Gamma}(p, \partial_p)(xp)^n$, from which one derives for the second line:

$$
\sum_{n=0}^{\infty} \frac{(\mathrm{i}\kappa)^n}{n!} \Big(\mathcal{P}^{(\tau)\Gamma'}_{n\ \Gamma}(x, \partial_x)\, x^{\mu_1} \dots x^{\mu_n}\Big) \int \mathrm{d}^4 p \left(\mathcal{P}^{(\tau)\Gamma''}_{n\ \Gamma'}(p, \partial_p)\, p_{\mu_1} \dots p_{\mu_n}\right)\langle P_2, S_2|\mathcal{O}_{\Gamma'' n}(p)|P_1, S_1\rangle.
$$

Here, the integrand contains the matrix elements of the irreducible local operators of definite twist $\mathcal{O}^{(\tau)}_{\Gamma\mu_1\dots\mu_n}(p)$. Their reduced matrix elements $f^{(\tau)}_{n_1 n_2}$, $n_1 + n_2 = n$, which are related to the decomposition of $(P_2 + P_1)^n$ into independent monomials may be represented by (double) moments of corresponding double distributions $f^{(\tau)}_{a}(z_1, z_2)$ multiplied by the kinematical factors $\mathcal{K}^{a}_{\Gamma}(P_1, P_2)$. Finally, since p reflects the dependence of the (local) operators on $\overleftrightarrow{D}$, after performing the integration and the resummation over n, p within the integrand simply has to be replaced by $\mathbb{P}\mathbb{Z}$.

Formally, in the various experimental situations, in any Fourier integrand having the structure of Eq. (3.3.5), we only have to perform the following replacements (denoted by $\doteq$):

(A) In the case of *non-forward scattering*, as just explained, we obtain

$$
\langle P_2, S_2|\mathcal{O}_{\Gamma}(p)|P_1, S_1\rangle \doteq \mathcal{K}^{a}_{\Gamma}(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, \delta^{(4)}(p - \mathbb{P}\mathbb{Z})\, f^{(\tau)}_{a}(\mathbb{Z}; \mu^2). \qquad (3.3.6)
$$

(B) In the case of *forward scattering*, i.e., for $P_1 = P_2 = P$, the situation changes into

$$
\langle P, S|\mathcal{O}_{\Gamma}(p)|P, S\rangle \doteq \mathcal{K}^{a}_{\Gamma}(P) \int \mathrm{d}z\, \delta^{(4)}(p - 2Pz)\, \hat{f}^{(\tau)}_{a}(z; \mu^2), \qquad (3.3.7)
$$

with $P_+ = 2P, P_- = 0; z_+ = z$. These distributions $\hat{f}^{(\tau)}_{a}(z, \mu^2)$ are obtained from the double distributions $\hat{f}^{(\tau)}_{a}(\mathbb{Z}, \mu^2)$ by integrating out the independent variable z_-, i.e.,

$$
\hat{f}^{(\tau)}_{a}(z, \mu^2) = \int \mathrm{d}z_- f^{(\tau)}_{a}(z_+ = z, z_-; \mu^2). \qquad (3.3.8)
$$

(C) In the case of *vacuum-to-hadron transition amplitudes*, e.g., for the meson distribution amplitudes one obtains

$$
\langle 0|\mathcal{O}_{\Gamma}(p)|V(P, \lambda)\rangle \doteq \widetilde{\mathcal{K}}^{a}_{\Gamma}(P) \int \mathrm{d}\xi\, \delta^{(4)}(p - P\xi)\, \tilde{f}^{(\tau)}_{a}(\xi; \mu^2), \qquad (3.3.9)
$$

with $P_2 = P; z_2 = \xi$. Obviously, the spin structures are different from the above cases and, by construction, only P and ξ occur.

From the foregoing discussion it is obvious that the target mass corrections to the various physical processes are completely determined by the twist structure of the bilocal operators $\mathcal{O}_\Gamma^{(\tau)}(\kappa x,-\kappa x)=\int \mathrm{d}^4p\,\mathcal{O}_\Gamma^{(\tau)}(p)\,\mathrm{e}^{\mathrm{i}\kappa xp}$. Performing matrix elements according to the replacements Eqs. (3.3.6) – (3.3.9) we obtain the related expressions for the S-matrix elements of the physical process under consideration. Then, a further Fourier transformation with respect to the coordinate x leads to a representation of the scattering or transition amplitudes in terms of the inverse momentum transfer depending, quite generally, on P_iP_j/Q^2.

The technique of harmonic polynomials

Now, we use the form of the harmonic extension in terms of Gegenbauer polynomials. This *polynomial technique* uses the vector $x\in\mathbb{R}^4$ as a device for writing tensors with special symmetries in analytic form [19, 48, 49]. It has the advantage to be directly related to the irreducible tensor representations of the Lorentz group. Its group theoretical background as far as it is related to totally symmetric tensors has been given in Ref. [19].

The local operator according to (2.2.15) is given by

$$O_{n+1}(x)\equiv\bar\psi(0)(x\gamma)(\mathrm{i}x\overleftrightarrow{D})^n\psi(0)=\int \mathrm{d}^4p\,\big(\bar\psi\gamma^\mu\psi\big)(p)\,x_\mu\,(xp)^n, \tag{3.3.10}$$

which shows that, in this connection, p formally replaces the covariant derivative $\overleftrightarrow{D}$ sandwiched between the quark operators. This local operator has a (finite) twist decomposition whose complete series will be given later on, cf., Eq. (4.4.15). The twist-2 part is given by the harmonic polynomials [121, 19] (see Section 4.1),

$$O^{\mathrm{tw2}}_{n+1}(x)=\sum_{k=0}^{[\frac{n+1}{2}]}\frac{(-1)^k(n+1-k)!}{4^kk!(n+1)!}\,x^{2k}\Box^kO_{n+1}(x)\equiv\mathcal{P}^{(2)}_{n+1}(x)\,O_{n+1}(x), \tag{3.3.11}$$

being characterized by traceless, totally symmetric tensors of rank $n+1$ whose indices are completely contracted by $x^{\mu_1}\cdots x^{\mu_{n+1}}$. They obey the condition of harmonicity: $\Box O^{\mathrm{tw2}}_{n+1}(x)=0$.

Performing the derivatives,

$$x^{2k}\Box^k(\gamma x)(px)^n=\frac{n!}{(n-2k)!}\,(\gamma x)(p^2x^2)^k(px)^{n-2k}+\frac{2k\,n!}{(n+1-2k)!}\,x^2(\gamma p)(p^2x^2)^{k-1}(px)^{n+1-2k}, \tag{3.3.12}$$

and using the series expansion of the Gegenbauer polynomials (see, e.g., Ref. [112], Appendix II.11),

$$C^\nu_n(z)=\frac{1}{(\nu-1)!}\sum_{k=0}^{[\frac{n}{2}]}\frac{(-1)^k(n-k+\nu-1)!}{k!(n-2k)!}\,(2z)^{n-2k}, \tag{3.3.13}$$

we obtain

$$\begin{aligned}O^{\mathrm{tw2}}_{n+1}(x)=\frac{1}{n+1}\int \mathrm{d}^4p\,\big(\bar\psi\gamma^\mu\psi\big)(p)\bigg\{&x_\mu\left(\frac12\sqrt{p^2x^2}\right)^nC^2_n\bigg(\frac{px}{\sqrt{p^2x^2}}\bigg)\\&-\frac12\,p_\mu\,x^2\left(\frac12\sqrt{p^2x^2}\right)^{n-1}C^2_{n-1}\bigg(\frac{px}{\sqrt{p^2x^2}}\bigg)\bigg\}.\end{aligned} \tag{3.3.14}$$

Eq. (3.3.14) is the analytic continuation from Euclidean spacetime — where the Gegenbauer polynomials obey the well-known orthonormality relations within the region $-1 \leq z \leq 1$ — to the Minkowski spacetime where the arguments of the square roots and of the Gegenbauer polynomials, depending on whether p^2 and/or x^2 are space-like or time-like, may be imaginary and, furthermore, could take values outside that region. This, however, has no influence on the validity of the above result since these polynomials being entire analytic functions are well defined on the whole complex plane.

Finally, according to the above mentioned interpretation of p as some kind of 'operator symbol' within the Fourier integral, Eq. (3.3.14) operationally has to be understood as if the polynomial inside the curly brackets had been written in terms of $\overset{\leftrightarrow}{D}$ instead of p and inserted into the operator $\bar{\psi}(0)\gamma^\mu\psi(0)$.

We now use the formula (see, e.g., Ref. [112], Eq. II.5.13.1.3),

$$\sum_{n=0}^{\infty} \frac{a^n}{(2\nu)_n} C_n^\nu(z) = \Gamma\left(\nu + \frac{1}{2}\right) \left(\frac{a}{2}\sqrt{1-z^2}\right)^{1/2-\nu} J_{\nu-1/2}\left(a\sqrt{1-z^2}\right) \mathrm{e}^{za}, \tag{3.3.15}$$

where $(2\nu)_n = 2\nu(2\nu{+}1)\ldots(2\nu{+}n{-}1) = \Gamma(n{+}2\nu)/\Gamma(2\nu)$ is the Pochhammer symbol. Choosing $z = (px)/\sqrt{p^2x^2}$ and $a = \mathrm{i}\sqrt{p^2x^2}/2$, Eq. (3.3.14) can be summed up according to Eq. (2.2.14) to the bilocal operator of twist-2:

$$\begin{aligned} O^{\text{tw2}}(\kappa x, -\kappa x) &= \sum_{n=0}^{\infty} \frac{(\mathrm{i}\kappa)^n}{n!} O_{n+1}^{\text{tw2}}(x) \\ &= \sqrt{\pi} \int \mathrm{d}^4 p \left(\bar{\psi}\gamma^\mu\psi\right)(p) \left\{ x_\mu \left(2 + x\partial\right) - \frac{1}{2}\mathrm{i}\kappa p_\mu x^2 \right\} (3 + x\partial) \\ &\qquad \times \left(\kappa\sqrt{(px)^2 - p^2x^2}\right)^{-3/2} J_{3/2}\left(\frac{\kappa}{2}\sqrt{(px)^2 - p^2x^2}\right) \mathrm{e}^{\mathrm{i}\kappa px/2}. \end{aligned} \tag{3.3.16}$$

The homogeneous derivations $(c + x\partial)$ are required to compensate the factors $(c + n)$ which are necessary in order to by able to introduce the Pochhammer symbols in the denominator. Obviously, for the second term in Eq. (3.3.14), after shifting $n \to n+1$ in the series over n, only one additional factor is required. — Also here p has to be considered as a symbol replacing the covariant derivatives sandwiched between the quark operators.

In order to show the main features of this formalism, we consider the forward as well as non-forward matrix elements. Let us now discuss the forward matrix elements of the twist-2 operators (3.3.14) and (3.3.16). The forward matrix elements are given by the reduced matrix element $f_n^{(2)}$ times the irreducible tensor written in terms of the momentum $P_+ = 2P$ and using $2P \equiv x_\mu \bar{U}(P)\gamma^\mu U(P)$. The final result can be written as

$$\begin{aligned} \langle P|O_{n+1}^{\text{tw2}}(x)|P\rangle &= f_n^{(2)}\, \mathcal{P}_{n+1}^{(2)}(x) \left\{ x_\mu \bar{U}(P)\gamma^\mu U(P)(2xP)^n \right\} \qquad (3.3.17) \\ &= \frac{f_n^{(2)}}{n+1}(2Px) \left\{ \left(\sqrt{P^2x^2}\right)^n C_n^2\!\left(\frac{Px}{\sqrt{P^2x^2}}\right) - \frac{P^2\,x^2}{(xP)} \left(\sqrt{P^2x^2}\right)^{n-1} C_{n-1}^2\!\left(\frac{Px}{\sqrt{P^2x^2}}\right) \right\}. \end{aligned}$$

The reduced matrix elements $f_n^{(2)}$ are given as the moments of a distribution $f^{(2)}(z)$ by means of a Mellin transformation,

$$f_n^{(2)} = \int_{-1}^{1} \mathrm{d}z z^n f^{(2)}(z). \tag{3.3.18}$$

We obtain, after summing up over n, the following expression of the forward matrix elements of the non-local twist-2 operators

$$\begin{aligned}\langle P|O^{\rm tw2}(\kappa x,-\kappa x)|P\rangle &= 2\sqrt{\pi}(Px)\int_{-1}^{1}{\rm d}z\, f^{(2)}(z)\left\{(2+x\partial)-{\rm i}\kappa z\frac{P^2x^2}{(xP)}\right\}(3+x\partial)\\ &\times\left(2\kappa z\sqrt{(Px)^2-P^2x^2}\right)^{-3/2}J_{3/2}\left(\kappa z\sqrt{(Px)^2-P^2x^2}\right){\rm e}^{{\rm i}\kappa zPx}.\end{aligned} \tag{3.3.19}$$

If this expression is restricted to the light-cone, $x^2=0$, the well-known light-cone parton distribution is obtained. Namely, using the Poisson integral for the Bessel functions (cf., Ref. [21], Eq. II.7.12.7),

$$\Gamma\left(\nu+\frac{1}{2}\right)J_\nu(z)=\frac{1}{\sqrt{\pi}}\Big(\frac{z}{2}\Big)^{\nu}\int_{-1}^{1}{\rm d}t\,(1-t^2)^{\nu-1/2}\,{\rm e}^{{\rm i}tz}\qquad\text{for}\qquad{\rm Re}\,\nu>-\frac{1}{2}, \tag{3.3.20}$$

we obtain ($\xi=\kappa z(\tilde{x}P)$)

$$\begin{aligned}\langle P|O^{\rm tw2}(\kappa\tilde{x},-\kappa\tilde{x})|P\rangle &= 2(\tilde{x}P)\sqrt{\pi}\int{\rm d}z\,f^{(2)}(z)\,(2+\xi\partial_\xi)(3+\xi\partial_\xi)\big(2\xi\big)^{-3/2}J_{3/2}(\xi)\,{\rm e}^{{\rm i}\xi}\\ &= 2(\tilde{x}P)\int{\rm d}z\,f^{(2)}(z)\int_0^1{\rm d}t\,t(1-t)(3+t\partial_t)(2+t\partial_t)\,{\rm e}^{2{\rm i}t\xi}\\ &= 2(\tilde{x}P)\int{\rm d}z\,f^{(2)}(z)\int_0^1{\rm d}t\,t(1+t\partial_t)\,{\rm e}^{2{\rm i}t\xi}\\ &= 2(\tilde{x}P)\int{\rm d}z\,f^{(2)}(z)\,{\rm e}^{2{\rm i}\kappa z(\tilde{x}P)};\end{aligned} \tag{3.3.21}$$

in the second line we introduced the Poisson integral after shifting the integration variable $t\to(t+1)/2$ and followed by changing $\xi\partial_\xi\to t\partial_t$, then we partially integrated two times retaining finally only the surface term at $t=1$. As a result we arrived at the twist-2 parton distribution as introduced in [66], $f^{(2)}(z)\equiv F^{(2)}(z)$, which coincides with $f_1(z)$ in the notation adopted by Jaffe and Ji [78].

More generally, when non-forward matrix elements are taken the argumentation is almost the same. But now, in the case of the local operators, there occur $n+1$ different reduced matrix elements, $f^{(2)}_{nm}$, related to the monomials $P_1^m\,P_2^{n-m}$, $0\leq m\leq n$ which are obtained from $(P_2+P_1)^n$, and the corresponding non-forward matrix element reads:

$$\langle P_2|O^{\rm tw2}_{n+1}(x)|P_1\rangle=\sum_{m=0}^{n}\binom{n}{m}f^{(2)}_{nm}\big(\bar{U}(P_2)\gamma_\mu U(P_1)\big)\,\mathcal{P}^{(2)}_{n+1}(x^2,\partial^2)\Big\{x^\mu(xP_1)^m(xP_2)^{n-m}\Big\}. \tag{3.3.22}$$

Now let us rewrite the reduced matrix elements $f^{(2)}_{nm}$ as the double moments of a double distribution $f^{(2)}(z_1,z_2)$,

$$f^{(2)}_{nm}=\int_{-1}^{1}{\rm d}z_1\int_{-1}^{1}{\rm d}z_2\,z_1^m z_2^{n-m}f^{(2)}(z_1,z_2). \tag{3.3.23}$$

After resumming w.r.t. n and rewriting $f^{(2)}(z_1,z_2)=F^{(2)}_D(z_+,z_-)$ we finally arrive at the expression for the non-forward matrix element of the nonlocal operator $O^{\rm tw2}(\kappa x,-\kappa x)$ as

$$\begin{aligned}\langle P_2|O^{\rm tw2}(\kappa x,-\kappa x)|P_1\rangle &= \sqrt{\pi}\,\bar{U}(P_2)\gamma^\mu U(P_1)\int{\rm D}\mathbb{Z}\,F^{(2)}_D(\mathbb{Z})\left\{x_\mu\,(2+x\partial)-\frac{1}{2}{\rm i}\kappa\,\mathbb{P}_\mu\mathbb{Z}\,x^2\right\}(3+x\partial)\\ &\times\left(\kappa\sqrt{(x\mathbb{P}\mathbb{Z})^2-x^2(\mathbb{P}\mathbb{Z})^2}\right)^{-3/2}J_{3/2}\Big(\frac{\kappa}{2}\sqrt{(x\mathbb{P}\mathbb{Z})^2-x^2(\mathbb{P}\mathbb{Z})^2}\Big)\,{\rm e}^{{\rm i}\kappa x\mathbb{P}\mathbb{Z}/2}.\end{aligned} \tag{3.3.24}$$

This holds in the case $(x\mathbb{P}\mathbb{Z})^2 - x^2\mathbb{P}\mathbb{Z}^2 \geq 0$, otherwise we have to change into $I_\nu(z)$.

From this approach it becomes also obvious that the mass corrections of different physical processes are related to each other if they can be traced back to the same operator content. In the following we use this polynomial technique to determine the local as well as nonlocal quark operators of definite twist. Thereby, we generalize Nachtmann's approach not only to non-forward matrix elements of (non)local operators. In addition we consider also the case of more general tensor operators having nontrivial symmetry types.

3.3.2 Power Corrections of non-forward matrix elements

In this Section I will give the resummed matrix elements of kinematical mass corrections in the x-space for the virtual Compton scattering and for exclusive processes with vector as well as (pseudo) scalar mesons by means of harmonic operators considered in the preceding Sections.

Double distributions from quark–antiquark operators with definite twist

In order to obtain the non-forward matrix elements for the quark–antiquark operators with definite twist τ we have only to take matrix elements of the expressions given in the last Section and to use formula (3.3.6). The spin-dependence of the hadronic states $|P_i, S_i\rangle, i = 1, 2$, is completely contained in the independent kinematical structures $\mathcal{K}^a_\Gamma(\mathbb{P})$ being relevant for the processes under consideration, e.g., the Dirac and the Pauli structure $\bar{U}(P_2, S_2)\gamma_\mu U(P_1, S_1)$ and $\bar{U}(P_2, S_2)\sigma_{\mu\nu}P^\nu_- U(P_1, S_1)/M$, respectively, in case of the virtual Compton scattering.

Let me now give the final expressions for matrix elements of nonlocal as well as local operators up to twist-3 which are relevant for the power corrections of the various processes. For the non-forward matrix element of the nonlocal twist-2 vector operator, Eqs. (4.4.48) with (4.4.44) obeying the replacement (3.3.6), one gets

$$\begin{aligned}
\langle P_2, S_2 | O^{\text{tw}2}_\alpha(\kappa x, -\kappa x) | P_1, S_1\rangle &= \mathcal{K}^a_\mu(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, F^{(2)}_a(\mathbb{Z})\, \partial_\alpha \int_0^1 \mathrm{d}t \left\{ x^\mu\, (2 + x\partial) - \frac{1}{2}\, \mathrm{i}\, \kappa t\, (\mathbb{P}^\mu\mathbb{Z})\, x^2 \right\} \\
&\qquad\qquad \times (3 + x\partial)\ \mathcal{H}_2(\mathbb{P}\mathbb{Z}|\kappa t x) \\
&= \mathcal{K}^a_\mu(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, F^{(2)}_a(\mathbb{Z}) \int_0^1 \mathrm{d}t\, (2 + x\partial) \Big\{ \Big[(3 + x\partial)\, \delta^\mu_\alpha - \mathrm{i}\, \kappa t\, (\mathbb{P}^\mu\mathbb{Z})\, x_\alpha \Big] \mathcal{H}_2(\mathbb{P}\mathbb{Z}|\kappa t x) \\
&\quad + \Big[(3 + x\partial) \Big((4 + x\partial)\, \mathrm{i}\, \kappa t\, (\mathbb{P}_\alpha\mathbb{Z}) x^\mu - \frac{1}{2} (\mathrm{i}\kappa t)^2 \big((\mathbb{P}\mathbb{Z})^2 x^\mu x_\alpha + (\mathbb{P}^\mu\mathbb{Z})(\mathbb{P}_\alpha\mathbb{Z})\, x^2 \big) \Big) \\
&\qquad + \frac{1}{4} (\mathrm{i}\kappa t)^3\, (\mathbb{P}^\mu\mathbb{Z})(\mathbb{P}\mathbb{Z})^2 x_\alpha x^2 \Big] \mathcal{H}_3(\mathbb{P}\mathbb{Z}|\kappa t x) \Big\}. \qquad (3.3.25)
\end{aligned}$$

The corresponding local matrix element of the twist-2 operator, Eq. (4.4.47), with (4.4.43) reads

$$\begin{aligned}
\langle P_2, S_2 | O^{\text{tw}2}_{\alpha n}(x) | P_1, S_1\rangle &= \frac{1}{(n+1)^2}\, \mathcal{K}^a_\mu(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, F^{(2)}_a(\mathbb{Z}) \Big\{ \delta^\mu_\alpha\, h^2_n(\mathbb{P}\mathbb{Z}|x) - (\mathbb{P}^\mu\mathbb{Z}) x_\alpha\, h^2_{n-1}(\mathbb{P}\mathbb{Z}|x) \\
&\quad + 2x^\mu (\mathbb{P}_\alpha\mathbb{Z})\, h^3_{n-1}(\mathbb{P}\mathbb{Z}|x) - \big(x^\mu x_\alpha (\mathbb{P}\mathbb{Z})^2 + (\mathbb{P}^\mu\mathbb{Z})(\mathbb{P}_\alpha\mathbb{Z}) x^2 \big)\, h^3_{n-2}(\mathbb{P}\mathbb{Z}|x) \\
&\qquad + \frac{1}{2}\, (\mathbb{P}^\mu\mathbb{Z}) x_\alpha\, x^2 (\mathbb{P}\mathbb{Z})^2\, h^3_{n-3}(\mathbb{P}\mathbb{Z}|x) \Big\}. \qquad (3.3.26)
\end{aligned}$$

For the matrix element of the nonlocal operator (4.4.50) and local one (4.4.49), respectively, we obtain

$$
\begin{aligned}
\langle P_2, S_2|O^{\rm tw3}_{\alpha}(\kappa x, -\kappa x)|P_1, S_1\rangle &= 2\mathcal{K}^a_\mu(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, F^{(3)}_a(\mathbb{Z}) \int_0^1 \mathrm{d}t\ x^\beta \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]}\,(1 + x\partial) - x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad \times \mathcal{H}_1(\mathbb{P}\mathbb{Z}|\kappa t x) \\
&= \mathcal{K}^a_\mu(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, F^{(3)}_a(\mathbb{Z})\ (2 + x\partial) \int_0^1 \mathrm{d}t\, x^\beta \Big\{ \Big[2\,\mathrm{i}\kappa t\, \delta^\mu_{[\alpha}(\mathbb{P}_{\beta]}\mathbb{Z})(2 + x\partial) - (\mathrm{i}\kappa t)^2 \delta^\mu_{[\alpha} x_{\beta]} (\mathbb{P}\mathbb{Z})^2 \Big] \\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad \times \mathcal{H}_2(\mathbb{P}\mathbb{Z}|\kappa t x) \\
&\quad - \mathrm{i}\kappa t\, x_{[\alpha}(\mathbb{P}_{\beta]}\mathbb{Z}) \Big[2\mathrm{i}\kappa t\, (\mathbb{P}^\mu\mathbb{Z})\, (5 + x\partial) - (\mathrm{i}\kappa t)^2 x^\mu (\mathbb{P}\mathbb{Z})^2 \Big]\, \mathcal{H}_3(\mathbb{P}\mathbb{Z}|\kappa t x) \Big\},
\end{aligned}
\tag{3.3.27}
$$

$$
\begin{aligned}
\langle P_2, S_2|O^{\rm tw3}_{\alpha n}(x)|P_1, S_1\rangle = \frac{2}{(n+1)^2}\, \mathcal{K}^a_\mu(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, F^{(3)}_a(\mathbb{Z})\, x^\beta \Big\{ & 2(n+1)\delta^\mu_{[\alpha}(\mathbb{P}_{\beta]}\mathbb{Z})\, h^2_{n-1}(\mathbb{P}\mathbb{Z}|x) \\
+ (n+2)\, x_{[\alpha}\delta^\mu_{\beta]}(\mathbb{P}\mathbb{Z})^2\, h^2_{n-2}(\mathbb{P}\mathbb{Z}|x) & - 4x_{[\alpha}(\mathbb{P}_{\beta]}\mathbb{Z})(\mathbb{P}^\mu\mathbb{Z})\, h^3_{n-2}(\mathbb{P}\mathbb{Z}|x) \\
& + 2x_{[\alpha}(\mathbb{P}_{\beta]}\mathbb{Z}) x^\mu (\mathbb{P}\mathbb{Z})^2\, h^3_{n-3}(\mathbb{P}\mathbb{Z}|x) \Big\}.
\end{aligned}
\tag{3.3.28}
$$

In the case of the axial vector operators $O^\tau_{5\alpha}(\kappa x, -\kappa x)$ the kinematical structures $\mathcal{K}_{5\mu}(\mathbb{P})$ contain an additional γ_5 and the double distributions should be denoted by $G^{(\tau)}_a(\mathbb{Z})$ but the formal structure will be the same.

The non-forward matrix elements of the nonlocal twist-3 chiral-odd scalar operator, Eq. (4.4.52), reads

$$
\langle P_2|N^{\rm tw3}(\kappa x, -\kappa x)|P_1\rangle = \mathcal{K}^a(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, E^{(3)}_a(\mathbb{Z})\ (1 + x\partial)\ \mathcal{H}_1(\mathbb{P}\mathbb{Z}|\kappa x),
\tag{3.3.29}
$$

and of the local one (4.4.51)

$$
\langle P_2|N^{\rm tw3}_n(x)|P_1\rangle = \mathcal{K}^a(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, E^{(3)}_a(\mathbb{Z})\, h^1_n(\mathbb{P}\mathbb{Z}|x).
\tag{3.3.30}
$$

The non-forward matrix element of the nonlocal twist-2 chiral-odd skew tensor operator, Eq. (4.4.54), is given by

$$
\begin{aligned}
\langle P_2, S_2|M^{\rm tw2}_{[\alpha\beta]}(\kappa x, -\kappa x)|P_1, S_1\rangle &= 2\mathcal{K}^a_{[\mu\nu]}(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, H^{(2)}_a(\mathbb{Z}) \int_0^1 \mathrm{d}t\, t \\
&\times \Big\{ (2 + x\partial)\delta^\mu_{[\alpha}\partial_{\beta]} - x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \Big[x^\nu\, (3 + x\partial) - \frac{1}{2}\,\mathrm{i}\kappa t\, (\mathbb{P}^\nu\mathbb{Z}) x^2 \Big] \mathcal{H}_2(\mathbb{P}\mathbb{Z}|\kappa t x),
\end{aligned}
\tag{3.3.31}
$$

and of the corresponding local operator (4.4.53) reads

$$
\begin{aligned}
\langle P_2, S_2|M^{\rm tw2}_{[\alpha\beta]n}(x)|P_1, S_1\rangle = \frac{2}{(n+2)(n+1)} & \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]} - \frac{1}{n+2}\, x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \mathcal{K}^a_{[\mu\nu]}(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, H^{(2)}_a(\mathbb{Z}) \\
& \times \Big\{ x^\nu h^2_n(\mathbb{P}\mathbb{Z}|x) - \frac{1}{2}\, (\mathbb{P}^\nu\mathbb{Z}) x^2 h^2_{n-1}(\mathbb{P}\mathbb{Z}|x) \Big\}.
\end{aligned}
\tag{3.3.32}
$$

The related nonlocal twist-3 operator, Eq. (4.4.56), and local one (4.4.55), respectively, reads

$$
\begin{aligned}
\langle P_2, S_2|M^{\rm tw3}_{[\alpha\beta]}(\kappa x, -\kappa x)|P_1, S_1\rangle = 2\mathcal{K}^a_{[\mu\nu]}(\mathbb{P}) & \int \mathrm{D}\mathbb{Z}\, H^{(3)}_a(\mathbb{Z})\, x_{[\alpha}\partial_{\beta]}\, \mathrm{i}\kappa\, (\mathbb{P}^\mu\mathbb{Z}) x^\nu (2 + x\partial) \\
& \times \int_0^1 \mathrm{d}t\, \mathcal{H}_2(\mathbb{P}\mathbb{Z}|\kappa t x),
\end{aligned}
\tag{3.3.33}
$$

$$\langle P_2, S_2|M^{\mathrm{tw3}}_{[\alpha\beta]n}(x)|P_1, S_1\rangle = \frac{2}{(n+2)n}\, x_{[\alpha}\partial_{\beta]}\, \mathcal{K}^a_{[\mu\nu]}(\mathbb{P}) \int \mathrm{D}\mathbb{Z}\, H^{(3)}_a(\mathbb{Z})\, (\mathbb{P}^\mu\mathbb{Z})x^\nu h^2_{n-1}(\mathbb{P}\mathbb{Z}|x). \tag{3.3.34}$$

Additionally, the non-forward matrix element of the independent nonlocal twist-3 skew tensor, Eq. (4.4.58), and local operator (4.4.57), respectively, reads

$$\begin{aligned}\langle P_2, S_2|\widetilde{M}^{\mathrm{tw3}}_{[\alpha\beta]}(\kappa x, -\kappa x)|P_1, S_1\rangle = 3\widetilde{\mathcal{K}}^a_{[\mu\nu]}(\mathbb{P})\int \mathrm{D}\mathbb{Z}\, \widetilde{H}^{(3)}_a(\mathbb{Z})\, x^\gamma \Big\{\delta^\mu_{[\alpha}\delta^\nu_\beta\partial_{\gamma]}x\partial - 2\,\delta^\mu_{[\alpha}x_\beta\partial_{\gamma]}\partial^\nu\Big\}\\ \times(1 + x\partial)\int_0^1 \mathrm{d}t\, \frac{1-t^2}{2t}\, \mathcal{H}_1(\mathbb{P}\mathbb{Z}|\kappa t x),\end{aligned} \tag{3.3.35}$$

$$\langle P_2, S_2|\widetilde{M}^{\mathrm{tw3}}_{[\alpha\beta]n}(x)|P_1, S_1\rangle = \frac{3x^\gamma}{n+2}\left\{\delta^\mu_{[\alpha}\delta^\nu_\beta\partial_{\gamma]}x\partial - \frac{2}{n}\,\delta^\mu_{[\alpha}x_\beta\partial_{\gamma]}\partial^\nu\right\}\widetilde{\mathcal{K}}^a_{[\mu\nu]}(\mathbb{P})\int \mathrm{D}\mathbb{Z}\, \widetilde{H}^{(3)}_a(\mathbb{Z})\, h^1_n(\mathbb{P}\mathbb{Z}|x). \tag{3.3.36}$$

Now, some remarks are in order. First, as indicated, different tensorial structures of the operators lead to different kinematical structures. In the forward case they simplify or eventually disappear, e.g., the Dirac structures are to be replaced by P_μ and S_μ for the vector and axial vector case, respectively, whereas the Pauli structures vanish. In the forward case we obtain from Eqs. (3.3.26), (3.3.28), and (3.3.30) and the replacement $x \to q$ the matrix elements of the local operators in the q-space $\langle PS|O^{\mathrm{tw2}}_{\alpha n}(q)|PS\rangle$, $\langle PS|O^{\mathrm{tw3}}_{\alpha n}(q)|PS\rangle$, and $\langle PS|N^{\mathrm{tw3}}_n(q)|PS\rangle$ which are in agreement with Wandzura's forward matrix elements in the q-space (see Eqs. (C.6), (C.9), and (C.1) in Ref. [124]).

Second, going on-cone the double distributions remain the same. Therefore, their evolution is determined by the anomalous dimensions resulting from the renormalization group equation of the corresponding light-cone operators of geometric twist.

Furthermore, we have to mention that – contrary to the case when the non-forward matrix elements are restricted to the light-cone where the decomposition of the non-local quark-antiquark operators into operators of definite twist terminates at finite values $\tau_{\max}$ – the off-cone decomposition results in an infinite series of any twist.

Mass corrections of vector meson distributions

In this Subsection I extend the analysis of the ρ-meson DAs to include all meson momentum resp. mass terms of harmonic twist-2 and twist-3 off-cone operators in order to get the meson mass corrections to the results obtained in Section 3.2. Vector meson mass corrections of order x^2 were already discussed by Ball and Braun [15] and resummed meson mass corrections in the scalar case was given by Ball [16] (see also Ref. [11]).

The structure of the mass corrections of ρ-meson DAs of geometric twist are, from the group theoretical point of view, similar to the target mass corrections in deep inelastic scattering, which can be resummed using Nachtmann's method [107]. The main tool, in order to obtain the mass corrections, are the harmonic operators of definite geometric twist which have been determined in Section 3.2. Obviously, they are the harmonic extensions of the corresponding light-cone operators which I have already used in Section 3.2 for the classification of the corresponding meson light cone DAs with respect to geometric twist.

Let me now introduce the distribution functions for the harmonic operators of geometric twist sandwiched between the vacuum and the meson state, $\langle 0|O^{(\tau)}_\Gamma(x, -x)|\rho(P, \lambda)\rangle$. Thereby, I adopt again the definitions of Chernyak and Zhitnitsky [42] in the terminology of Ref. [14]. As usual, these matrix elements are related to the momentum P_α and polarization vector $e^{(\lambda)}_\alpha$ of the meson with helicity λ, $P^2 = m_\rho^2$, $(e^{(\lambda)}e^{(\lambda)}) = -1$, $(Pe^{(\lambda)}) = 0$.

First, we consider the chiral-even *vector operator*. Using the twist projections (4.4.40) we introduce the moments of the meson DAs, $\Phi^{(\tau)}_n$, of twist τ as reduced matrix elements according to

$$\langle 0|O^{(\tau)}_{\alpha n}(x)|\rho(P,\lambda)\rangle = f_\rho m_\rho \, \mathcal{P}^{(\tau)\beta}_{\alpha n}\left(e^{(\lambda)}_\beta (Px)^n \Phi^{(\tau)}_n\right), \qquad (3.3.37)$$

where f_ρ is again the vector meson decay constant. The corresponding meson distribution amplitudes $\hat{\Phi}^{(\tau)}(\xi)$ are given by inverting the moment integral,

$$\Phi^{(\tau)}_n = \int_{-1}^{1} \mathrm{d}\xi \, \xi^n \hat{\Phi}^{(\tau)}(\xi), \quad \xi = u - (1-u) = 2u - 1. \qquad (3.3.38)$$

These distribution amplitudes are dimensionless functions of ξ and describe the probability amplitudes to find the ρ-meson in a state with minimal number of constituents (quark and antiquark) which carry the momentum fractions u (quark) and $(1-u)$ (antiquark).

Taking into account expression (4.4.47) we obtain the local twist-2 matrix element in the x-space as follows:

$$\begin{aligned} &\langle 0|O^{\mathrm{tw2}}_{\alpha n}(x)|\rho(P,\lambda)\rangle = && (3.3.39)\\ &= \frac{1}{(n+1)^2} f_\rho m_\rho \, \Phi^{(2)}_n \Big\{ e^{(\lambda)}_\alpha h^2_n(P|x) + 2P_\alpha (e^{(\lambda)}x) h^3_{n-1}(P|x) - m_\rho^2 x_\alpha (e^{(\lambda)}x) h^3_{n-2}(P|x) \Big\}, \end{aligned}$$

which is analogous to Wandzura's expression, $\langle PS|O^{\mathrm{tw2}}_{5\alpha n}(q)|PS\rangle$, (Eq. (C.6) in [124]) for the target mass corrections in deep inelastic scattering in the q-space. After resummation we get the bilocal matrix element of geometric twist-2:

$$\begin{aligned} \langle 0|O^{\mathrm{tw2}}_{\alpha}(x,-x)|\rho(P,\lambda)\rangle &= f_\rho m_\rho \int_0^1 \mathrm{d}t \int_{-1}^{1} \mathrm{d}\xi \, \hat{\Phi}^{(2)}(\xi) \, (3 + \xi\partial_\xi) \, (2 + \xi\partial_\xi) \qquad (3.3.40)\\ &\times \left\{ e^{(\lambda)}_\alpha \mathcal{H}_2(P\xi|tx) + \left((4 + \xi\partial_\xi) \, (\mathrm{i}\xi t) P_\alpha - \frac{1}{2}(\mathrm{i}\xi t)^2 m_\rho^2 x_\alpha \right)(e^{(\lambda)}x)\mathcal{H}_3(P\xi|tx) \right\}. \end{aligned}$$

The trace subtraction of the twist-2 operator give rise to the so-called "kinematical" target-mass corrections. For the forward-scattering case, contributions of this type have been considered by Nachtmann. For the vector and tensor operators, we obtain mass corrections which are proportional to x_α [5] and terms proportional to x^2 which are resummed to Gegenbauer polynomials (local operators) or Bessel functions (nonlocal operators).

An analogous calculation using expression (4.4.49) gives the local twist-3 matrix element

$$\begin{aligned} \langle 0|O^{\mathrm{tw3}}_{\alpha n}(x)|\rho(P,\lambda)\rangle = \frac{2}{(n+1)} f_\rho m_\rho \, \Phi^{(3)}_n x^\beta \Big\{ & e^{(\lambda)}_{[\alpha} P_{\beta]} h^2_{n-1}(P|x) + \frac{n+2}{2(n+1)} m_\rho^2 x_{[\alpha} e^{(\lambda)}_{\beta]} h^2_{n-2}(P|x) \\ & + \frac{1}{n+1} m_\rho^2 x_{[\alpha} P_{\beta]} (e^{(\lambda)}x) h^3_{n-3}(P|x) \Big\}, \qquad (3.3.41) \end{aligned}$$

which is close to Wandzura's expression, $\langle PS|O^{\mathrm{tw3}}_{5\alpha n}(q)|PS\rangle$, (Eq. (C.9) in [124]). Additionally the resummed bilocal matrix element of twist-3 reads

$$\begin{aligned} \langle 0|O^{\mathrm{tw3}}_{\alpha}(x,-x)|\rho(P,\lambda)\rangle = f_\rho m_\rho \int_{-1}^{1} \mathrm{d}\xi \, \hat{\Phi}^{(3)}(\xi) \, (2 + \xi\partial_\xi) \, x^\beta \Big\{ & 2(\mathrm{i}\xi) e^{(\lambda)}_{[\alpha} P_{\beta]} \mathcal{H}_2(P\xi|x) \\ + m_\rho^2 \int_0^1 \mathrm{d}t \Big((\mathrm{i}\xi t)^2 x_{[\alpha} e^{(\lambda)}_{\beta]} \mathcal{H}_2(P\xi|tx) + (\mathrm{i}\xi t)^3 x_{[\alpha} P_{\beta]} (e^{(\lambda)}x) & \mathcal{H}_3(P\xi|tx) \Big) \Big\}. \qquad (3.3.42) \end{aligned}$$

---

[5] This kind of mass corrections survives on the light-cone (see trace terms in Section 3.2).

It is well-known that the twist-3 operator $O^{\rm tw3}_\alpha(x,-x)$ is related to other twist-3 operators containing total derivatives and operators of Shuryak-Vainshtein type by means of QCD equations of motion [9, 15]. Therefore, the matrix elements (3.3.41) and (3.3.42) include the contribution of the twist-3 operator containing total derivatives which is as large as those from twist-2 (see also [17]) and also give rise to so-called "dynamic" mass corrections. Dynamical mass-corrections have so far only been considered for the exclusive case, Ref. [15]. In addition the next higher twist contributions of the chiral-even vector operator are of twist-4 which also contain dynamical mass corrections. They act in the same direction as the mass corrections of twist-2 and twist-3 operators ($\sim x_\alpha$ and $\sim x^2$).

Now we consider the chiral-even *axial vector operator* $O_{5\alpha n}(x)$ with the same twist projectors (4.4.40) as for the chiral-even vector operator $O_{\alpha n}(x)$. We define the corresponding moments of the meson DAs $\Xi^{(\tau)}_n$ by

$$\langle 0|O^{(\tau)}_{5\alpha n}(x)|\rho(P,\lambda)\rangle = \frac{1}{2}\Big(f_\rho - f_\rho^{\rm T}\frac{m_u+m_d}{m_\rho}\Big) m_\rho\, \mathcal{P}^{(\tau)\beta}_{\alpha n}\Big(\epsilon_\beta{}^{\gamma\mu\nu} e^{(\lambda)}_\gamma P_\mu x_\nu (Px)^n \Xi^{(\tau)}_n\Big)\,, \tag{3.3.43}$$

where $f_\rho^{\rm T}$ denotes the tensor decay constant. First, we observe that the twist-2 contribution vanishes. The nontrivial local vacuum-to-meson matrix elements of this axial vector operator are of twist-3:

$$\langle 0|O^{\rm tw3}_{5\alpha n}(x)|\rho(P,\lambda)\rangle = \frac{1}{2}\Big(f_\rho - f_\rho^{\rm T}\frac{m_u+m_d}{m_\rho}\Big) m_\rho\, \epsilon_\alpha{}^{\beta\mu\nu} e^{(\lambda)}_\beta P_\mu x_\nu\, \Xi^{(3)}_n h^1_n(P|x), \tag{3.3.44}$$

and the bilocal matrix element of twist-3 reads

$$\langle 0|O^{\rm tw3}_{5\alpha}(x,-x)|\rho(P,\lambda)\rangle = \frac{1}{2}\Big(f_\rho - f_\rho^{\rm T}\frac{m_u+m_d}{m_\rho}\Big) m_\rho\, \epsilon_\alpha{}^{\beta\mu\nu} e^{(\lambda)}_\beta P_\mu x_\nu \int_{-1}^{1} {\rm d}\xi\, \hat\Xi^{(3)}(\xi)\,(1+\xi\partial_\xi) \times \mathcal{H}_1(P\xi|x). \tag{3.3.45}$$

By the way the twist-4 matrix element also vanishes and the next higher twist contribution would be of twist-5. Also here, the matrix elements (3.3.44) and (3.3.45) include the contribution of the twist-3 operator containing total derivatives.

The matrix element of the chiral-odd *scalar operator* (see Eq. (4.4.41)) is defined as

$$\langle 0|N^{\rm tw3}_n(x)|\rho(P,\lambda)\rangle = -{\rm i}\Big(f_\rho^{\rm T} - f_\rho\frac{m_u+m_d}{m_\rho}\Big) m_\rho^2 \mathcal{P}^{(3)}_n\Big(\big(e^{(\lambda)}x\big)(Px)^n \Upsilon^{(3)}_n\Big)\,, \tag{3.3.46}$$

where $\Upsilon^{(3)}_n$ is the moment of a spin-independent twist-3 distribution function. Using expression (4.4.51) the local matrix element is given as

$$\langle 0|N^{\rm tw3}_n(x)|\rho(P,\lambda)\rangle = -{\rm i}\Big(f_\rho^{\rm T} - f_\rho\frac{m_u+m_d}{m_\rho}\Big)\big(e^{(\lambda)}x\big) m_\rho^2\, \Upsilon^{(3)}_n h^1_n(P|x), \tag{3.3.47}$$

and for the bilocal matrix element of twist-3 we obtain

$$\langle 0|N^{\rm tw3}(x,-x)|\rho(P,\lambda)\rangle = -{\rm i}\Big(f_\rho^{\rm T} - f_\rho\frac{m_u+m_d}{m_\rho}\Big)\big(e^{(\lambda)}x\big) m_\rho^2 \int_{-1}^{1} {\rm d}\xi\, \hat\Upsilon^{(3)}(\xi)\,(1+\xi\partial_\xi)\,\mathcal{H}_1(P\xi|x). \tag{3.3.48}$$

The next higher twist contributions of the scalar chiral-odd operator are of order twist-5.

Now, we consider the matrix elements of the chiral-odd *skew tensor operators*. Using Eq. (4.4.42), the corresponding moments of the wave function may be introduced according to

$$\langle 0|M^{(\tau)}_{[\alpha\beta]n}(x)|\rho(P,\lambda)\rangle = \mathrm{i} f^{\mathrm{T}}_{\rho}\mathcal{P}^{(\tau)\mu\nu}_{[\alpha\beta]n}\left(\left(e^{(\lambda)}_{\mu}P_{\nu} - e^{(\lambda)}_{\nu}P_{\mu}\right)(Px)^{n}\Psi^{(\tau)}_{n}\right). \tag{3.3.49}$$

The local matrix element of the skew tensor operator of twist-2, performing the differentiations in the expression (4.4.53), is given by

$$\begin{aligned}\langle 0|M^{\mathrm{tw2}}_{[\alpha\beta]n}(x)|\rho(P,\lambda)\rangle &= \mathrm{i} f^{\mathrm{T}}_{\rho}\,\Psi^{(2)}_{n}\Big\{\frac{2}{n+1}\,e^{(\lambda)}_{[\alpha}P_{\beta]}h^{2}_{n}(P|x) + \frac{n+3}{(n+2)^{2}}\,m^{2}_{\rho}x_{[\alpha}e^{(\lambda)}_{\beta]}h^{2}_{n-1}(P|x)\\ &+ \frac{m^{2}_{\rho}}{(n+2)^{2}(n+1)}\Big\{4x_{[\alpha}e^{(\lambda)}_{\beta]}h^{3}_{n-1}(P|x) - \Big(2x_{[\alpha}e^{(\lambda)}_{\beta]}(Px) + 4x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)\Big)h^{3}_{n-2}(P|x)\\ &+ 12x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)h^{4}_{n-2}(P|x) - 6x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)(Px)h^{4}_{n-3}(P|x)\Big\}\Big\},\end{aligned} \tag{3.3.50}$$

and the resummed bilocal matrix element of twist-2 reads

$$\begin{aligned}\langle 0|M^{\mathrm{tw2}}_{[\alpha\beta]}(x,-x)|\rho(P,\lambda)\rangle &= \mathrm{i}\, f^{\mathrm{T}}_{\rho}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\Psi}^{(2)}(\xi)\,(3+\xi\partial_{\xi})\Big\{2\,(2+\xi\partial_{\xi})\,e^{(\lambda)}_{[\alpha}P_{\beta]}\mathcal{H}_{2}(P\xi|x)\\ &+ m^{2}_{\rho}\int_{0}^{1}\mathrm{d}t\,t\Big[(1+\xi\partial_{\xi})\,(\mathrm{i}\xi t)x_{[\alpha}e^{(\lambda)}_{\beta]}\mathcal{H}_{2}(P\xi|tx)\\ &+ \Big(2\,(4+\xi\partial_{\xi})\,(\mathrm{i}\xi t)x_{[\alpha}e^{(\lambda)}_{\beta]} - (\mathrm{i}\xi t)^{2}\big(x_{[\alpha}e^{(\lambda)}_{\beta]}(Px) + 2x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)\big)\Big)\mathcal{H}_{3}(P\xi|tx)\\ &+ \Big(2\,(5+\xi\partial_{\xi})\,(4+\xi\partial_{\xi})\,(\mathrm{i}\xi t)^{2}x_{[\alpha}P_{\beta]}(e^{(\lambda)}x) - (4+\xi\partial_{\xi})\,(\mathrm{i}\xi t)^{3}x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)(Px)\Big)\mathcal{H}_{4}(P\xi|tx)\Big]\Big\}.\end{aligned} \tag{3.3.51}$$

The local matrix element of the skew tensor operator of twist-3 is obtained from (4.4.55) as follows:

$$\langle 0|M^{\mathrm{tw3}}_{[\alpha\beta]n}(x)|\rho(P,\lambda)\rangle = -\frac{2}{(n+2)n}\,\mathrm{i} f^{\mathrm{T}}_{\rho}\,\Psi^{(3)}_{n}m^{2}_{\rho}\Big\{x_{[\alpha}e^{(\lambda)}_{\beta]}h^{2}_{n-1}(P|x) + 2x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)h^{3}_{n-2}(P|x)\Big\}, \tag{3.3.52}$$

and the corresponding bilocal matrix element reads

$$\begin{aligned}\langle 0|M^{\mathrm{tw3}}_{[\alpha\beta]}(x,-x)|\rho(P,\lambda)\rangle &= -2\mathrm{i} f^{\mathrm{T}}_{\rho}m^{2}_{\rho}\int_{0}^{1}\frac{\mathrm{d}t}{t}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\Psi}^{(3)}(\xi)\,(1+\xi\partial_{\xi})\\ &\times\Big\{(\mathrm{i}\xi t)x_{[\alpha}e^{(\lambda)}_{\beta]}\mathcal{H}_{2}(P\xi|tx) + (3+\xi\partial_{\xi})\,(\mathrm{i}\xi t)^{2}x_{[\alpha}P_{\beta]}(e^{(\lambda)}x)\mathcal{H}_{3}(P\xi|tx)\Big\}.\end{aligned} \tag{3.3.53}$$

The next higher twist contributions of the skew tensor operator would be of twist-4. Again, the matrix elements (3.3.47), (3.3.48), (3.3.52) and (3.3.53) include the contribution of the twist-3 operator containing total derivatives. Let us note, that the contribution of the additional twist-3 operator, Eq. (4.4.57), vanishes.

Finally, we remark that after projection onto the light-cone the matrix elements, which have been introduced in Section (3.2), are recovered. This can be easily checked by using the expansion of the Gegenbauer polynomials or the Poisson integral representation of Bessel functions. Let us point to the fact that the x^2-dependence is contained in the polynomials $h^{\nu}_{n}(P|x)$ and the functions $\mathcal{H}_\nu(P|x)$ in the case of the moments and the wave functions, respectively. Since the latter is given by

$$2\Gamma(\nu)\mathcal{H}_\nu(P|x) = 4^\nu \int_{-1}^{+1} \mathrm{d}t(1-t^2)^{\nu-1}\mathrm{e}^{\mathrm{i}(Px)\left(1+t\sqrt{1-m_\rho^2x^2/(Px)^2}\right)/2} \tag{3.3.54}$$

one would obtain a fairly simple expansion in powers of $m_\rho^2 x^2/(Px)^2$. Because the meson DAs of definite twist are independent from any coordinates or momenta, they coincide on-cone and off-cone. The whole information about the power corrections is contained in twist projection of the operators from which the matrix elements are taken.

Trace-subtractions of leading twist operators give rise to the so-called "kinematical" (or rather geometric) mass-corrections. They are formally analogous to Nachtmann corrections in inclusive processes. Operators of higher twist can be decomposed into operators containing total derivatives and quark-quark-gluon operators of Shuryak-Vainshtein type entering by the EOM. The contributions of operators with total derivatives is a specific feature in exclusive processes as well as in off-forward scattering, which make the structure of these corrections much more complex. Numerically, these corrections turned out to be relevant. We conclude that, at least for exclusive vector-meson DAs, a resummation of mass-corrections induced by trace-subtractions in the leading twist matrix element and the higher twist operators containing total derivatives are relevant for a good approximation.

Mass corrections of pion distributions

Amplitudes of processes involving pseudoscalar mesons (pions) can be expressed in terms of matrix elements of two-particle operators sandwiched between the vacuum and the meson state $\langle 0|\bar{u}(x)\Gamma U(x,-x)d(-x)|\pi(P)\rangle$. We are now in particular interested in pion-mass corrections. In contrast to the vector mesons, the matrix elements of the pion are just related to the pion momentum P with $P^2 = m_\pi^2$ where m_π denotes the pion mass. Therefore, the DAs of pions are given by totally symmetric harmonic operators. Because no asymmetric Young tableaux give a contribution we can carry out the classification of the DAs by means of the totally symmetric scalar two particle operators of any twist. Thus we are able to make a classification of pion DAs up to infinite twist.

We start with the two particle DAs of minimal twist. The light-cone matrix-element of the pseudo scalar operator of twist-2 reads [42, 32]

$$\langle 0|O_5^{\mathrm{tw2}}(\tilde{x},-\tilde{x})|\pi(P)\rangle = \mathrm{i}f_\pi(P\tilde{x})\int_{-1}^{1}\mathrm{d}\xi\,\hat{\phi}_\pi^{(2)}(\xi)\,\mathrm{e}^{\mathrm{i}\xi P\tilde{x}}, \tag{3.3.55}$$

where $\hat{\phi}_\pi^{(2)}(\xi)$ is the twist-2 DA.

Using Eqs. (4.4.18) and (4.4.19), we are able to give the following (infinite) twist decomposition of the corresponding operator and of the DAs as follows

$$\begin{aligned}\langle 0|O_5(x,-x)|\pi(P)\rangle &= \langle 0|O_5^{\mathrm{tw2}}(x,-x)|\pi(P)\rangle \\ &\quad + \sum_{j=1}^{\infty}\frac{x^{2j}}{4^j j!(j-1)!}\int_0^1 \mathrm{d}t\,t\,(1-t)^{j-1}\langle 0|O_5^{\mathrm{tw}(2+2j)}(tx,-tx)|\pi(P)\rangle,\end{aligned} \tag{3.3.56}$$

with

$$\langle 0|O_5^{\mathrm{tw}(2+2j)}(tx,-tx)|\pi(P)\rangle = \mathrm{i}f_\pi \int_{-1}^{1}\mathrm{d}\xi\, \hat{\phi}_\pi^{(2+2j)}(\xi)\left(-m_\pi^2\xi^2\right)^j \Bigg\{ \left(t(Px)(2+\xi\partial_\xi) - \frac{1}{2}\,\mathrm{i}\xi t^2 m_\pi^2 x^2\right) \\ \times (3+\xi\partial_\xi)\,\mathcal{H}_2(P\xi|tx) - 2j\frac{\mathrm{i}}{\xi}\,(1+\xi\partial_\xi)\,\mathcal{H}_1(P\xi|tx)\Bigg\}. \quad (3.3.57)$$

Here $\hat{\phi}_\pi^{(2+2j)}(\xi)$ is a pion DA of twist $\tau = 2+2j$. The harmonic twist-2 operator is obtained from Eq. (3.3.57) by setting $t=1$ and $j=0$ and gives the pure kinematical target-mass corrections.

The light-cone matrix-element of the pseudo scalar operator of twist-3 is given by [32]

$$\langle 0|N_5^{\mathrm{tw3}}(\tilde{x},-\tilde{x})|\pi(P)\rangle = \frac{f_\pi m_\pi^2}{m_u+m_d}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\phi}_p^{(3)}(\xi)\,\mathrm{e}^{\mathrm{i}\xi P\tilde{x}}. \quad (3.3.58)$$

The (infinite) twist decomposition of the bilocal scalar operator $N_5(x,-x) = \bar{u}(x)\mathrm{i}\gamma_5 d(-x)$ into harmonic operators $N^{\mathrm{tw}(3+2j)}(x,-x)$ of twist $\tau = 3+2j$, $j = 0,1,2,\dots,$ is given by Eqs. (4.4.7) and (4.4.8) and reads

$$\langle 0|N_5(x,-x)|\pi(P)\rangle = \langle 0|N_5^{\mathrm{tw3}}(x,-x)|\pi(P)\rangle \\ + \sum_{j=1}^{\infty}\frac{x^{2j}}{4^j j!(j-1)!}\int_0^1 \mathrm{d}t\,t\,(1-t)^{j-1}\langle 0|N_5^{\mathrm{tw}(3+2j)}(tx,-tx)|\pi(P)\rangle, \quad (3.3.59)$$

with

$$\langle 0|N_5^{\mathrm{tw}(3+2j)}(tx,-tx)|\pi(P)\rangle = \frac{f_\pi m_\pi^2}{m_u+m_d}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\phi}_p^{(3+2j)}(\xi)\left(-m_\pi^2\xi^2\right)^j(1+\xi\partial_\xi)\,\mathcal{H}_1(P\xi|tx). \quad (3.3.60)$$

Eq. (3.3.60) is the generalization of Ball's [15] pion mass corrections of leading twist for any twist. These higher twist operators give also rise to dynamical target-mass corrections.

Another twist-3 light-cone matrix element is given as [32]

$$\langle 0|M_{5[\alpha\beta]}^{\mathrm{tw3}}(\tilde{x},-\tilde{x})|\pi(P)\rangle = P_{[\alpha}\tilde{x}_{\beta]}\frac{\mathrm{i}f_\pi m_\pi^2}{3(m_u-m_d)}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\phi}_\sigma^{(3)}(\xi)\,\mathrm{e}^{\mathrm{i}\xi P\tilde{x}}. \quad (3.3.61)$$

Using the parametrization of Eq. (3.3.61) together with Eqs. (4.4.21) and (4.4.22), we finally get the infinite twist decomposition:

$$\langle 0|M_5(x,-x)|\pi(P)\rangle = \langle 0|M_5^{\mathrm{tw3}}(x,-x)|\pi(P)\rangle \quad (3.3.62) \\ + \sum_{j=1}^{\infty}\frac{x^{2j}}{4^j j!(j-1)!}\int_0^1 \mathrm{d}t\,t\,(1-t)^{j-1}\,\langle 0|M_5^{\mathrm{tw}(3+2j)}(tx,-tx)|\pi(P)\rangle,$$

with

$$\langle 0|M_5^{\mathrm{tw}(3+2j)}(tx,-tx)|\pi(P)\rangle = \frac{f_\pi m_\pi^2}{6(m_u-m_d)}\int_{-1}^{1}\mathrm{d}\xi\,\hat{\phi}_\sigma^{(3+2j)}(\xi)\left(-m_\pi^2\xi^2\right)^j\,t\xi\left((Px)^2 - m_\pi^2x^2\right) \\ \times(2+\xi\partial_\xi)\,(3+\xi\partial_\xi)\,\mathcal{H}_2(P\xi|tx). \quad (3.3.63)$$

Again these higher twist operators contain also dynamical target-mass corrections.

Let me, additionally, note that the formal structure of Eqs. (3.3.57), (3.3.60) and (3.3.63) is, up to the factor $(-m_\pi^2\xi^2)^j$, the same for any twist.

3.3.3 Conclusions

In this Section, we introduced a general procedure of parametrizing non-forward matrix elements of off-cone QCD operators $\mathcal{O}_\Gamma(\kappa x, -\kappa x)$ in terms of two-variable distribution amplitudes $f_a^{(\tau)}(\mathbb{Z})$ of well-defined geometric twist τ, namely, single variable hadron distributions, double distributions etc..

The procedure relies on the unique twist decomposition of non-local operators off the light-cone leading to an infinite series of operators with growing twist. This decomposition is completely of group theoretical origin and is equivalent to the decomposition of the local operators into irreducible tensor representations of the Lorentz group.

Using these results we determined the off-cone power corrections to various double distributions and the vector as well as scalar meson distribution amplitudes being the inputs of the corresponding scattering amplitudes and hadronic form factors, respectively. These power corrections, in the case of local operators, are expressed in terms of Gegenbauer polynomials being multiplied with the moments of the distribution amplitudes and, in the case of non-local operators, in terms of Bessel functions now multiplied with the distribution amplitudes directly. Accordingly, the off-cone expressions are obtained from the on-cone ones by harmonic extension.

Concerning the computation of the scattering amplitudes of physical relevance some Fourier transformation has to be carried out whose result mainly depends on the (singular) coefficient functions as well as on the Gegenbauer polynomials or Bessel functions (see Appendix D). The easiest way to calculate the power corrections of scattering amplitudes is to use the local operators in terms of Gegenbauer polynomials in the q-space. The currently most interesting example is the (deeply) virtual Compton scattering.

Chapter 4

The twist decomposition of nonlocal tensor operators

In this Chapter we give the explicite twist decomposition of (non)local operators on the light cone as well as off-cone. In Section 4.1 we discuss the basics of irreducible tensors with respect to the Lorentz and rotation group in $2h$-dimensional spacetime ($2h \geq 3$) in the framework of polynomial technique. We are using the harmonic irreducible tensor polynomials with $h = 2$ in order to rewrite harmonic operators in terms of Gegenbauer polynomials and Bessel functions in Section 4.4. We have used these harmonic operators in Section 3.3 to calculate the power corrections of distribution amplitudes with definite geometric twist. In Section 4.2 we decompose a general bilocal tensor operator of second rank. In this connection we derive the twist decomposition of the light-cone operators $O_\alpha(\kappa_1\tilde{x}, \kappa_2\tilde{x})$ and $M_{[\alpha\beta]}(\kappa_1\tilde{x}, \kappa_2\tilde{x})$ which we used in Section 3.1 and 3.2. In Section 4.3 we discuss the twist decomposition of trilocal and multilocal operators on the light-cone.

4.1 Irreducible tensor polynomials and harmonic extension in $2h$-dimensional spacetime – polynomial technique

The objective of the present Section is to use the polynomial technique for symmetric tensor and some asymmetric tensor representations of the orthogonal group as well as the Lorentz group in $2h$-dimensions ($2h \geq 3$). By means of the polynomial technique we can use a $2h$-vector as a device for writing tensors with special symmetries in an analytic form.

In order to construct irreducible vector and tensor polynomials corresponding to irreducible tensors with special Young symmetry, we generalize the homogeneous polynomial technique which is well-known for the scalar case in constructing irreducible symmetric tensor representations of the orthogonal group $SO(2h, \mathbb{C})$.

(i) Let us now consider a symmetric traceless tensor $\overset{\circ}{T}_{\mu_1\ldots\mu_n}$ corresponding to the Young tableau $\boxed{\quad n \quad}$. Such tensors are, for the group $SO(2h, \mathbb{C})$, in one-to-one correspondence with harmonic polynomials of the complex $2h$-vector ζ:

$$\overset{\circ}{T}_n(\zeta) \equiv \zeta^{\mu_1} \cdots \zeta^{\mu_n}\, \overset{\circ}{T}_{\mu_1\ldots\mu_n}, \qquad \zeta \in \mathbb{C}^{2h} \qquad \Delta\overset{\circ}{T}_n(\zeta) = 0, \tag{4.1.1}$$

or, equivalently, with their restriction $T_n(z)$ on the complex light cone $K_{2h}(\mathbb{C})$:

$$T_n(z) \equiv z^{\mu_1} \cdots z^{\mu_n}\, T_{\mu_1\ldots\mu_n} = \lim_{\zeta \to z, z^2 = 0} \overset{\circ}{T}_n(\zeta). \tag{4.1.2}$$

Then, the space of (complex) harmonic polynomials $\overset{\circ}{T}_n(\zeta)$ and the space of homogeneous polynomials $T_n(z)$ on the complex light cone provide convenient realizations of the carrier space of irreducible finite dimensional symmetric tensor representations of the group $SO(2h,\mathbb{C})$ (see Ref. [19]).

For symmetric tensor representations of the group $SO(1,2h-1)$ the elements of the carrier space are harmonic (more strictly: $\Box$-harmonic) polynomials of degree n, $\overset{\circ}{T}_n(x)$, with respect to the real $2h$-vector x, iff

$$\Box\overset{\circ}{T}_n(x)=0, \qquad x\in M^{2h} \tag{4.1.3}$$

or, alternatively, their restrictions $T_n(\tilde{x})$ on the real light cone $K_{2h}(\mathbb{R})$. The tensor representations $\overset{\circ}{T}_n(\zeta)$ and $T_n(z)$ of $SO(2h,\mathbb{C})$ are equivalent to the analytic continuation of the representations $\overset{\circ}{T}_n(x)$ and $T_n(\tilde{x})$ of $SO(1,2h-1)$, respectively. In the following we consider finite dimensional representations of the group $SO(1,2h-1)$.

Let us now regard the equivalence between the representation $\overset{\circ}{T}_n(x)$ and $T_n(\tilde{x})$. For this we note the decomposition of any homogeneous polynomial in the form

$$T_n(x)=\overset{\circ}{T}_n(x)+x^2T_{n-2}(x), \quad \overset{\circ}{T}_n(x)\in\mathcal{H}^{2h,n}, \tag{4.1.4}$$

where $T_n(x)\in\mathcal{R}^{2h,n}$ is a homogeneous polynomial of degree n in x. On the light cone, the value of any polynomial coincides with the value on the cone of a uniquely defined harmonic polynomial. This establishes an isomorphism between the coset space $\mathcal{R}^{2h,n}/x^2\mathcal{R}^{2h,n-2}$ and the space $\mathcal{H}^{2h,n}$.

By means of the extension $T_n(x)$ for any polynomial $T_n(\tilde{x})$ on the light cone, the harmonic extension is defined by

$$\Box\overset{\circ}{T}_n(x)=0, \quad \overset{\circ}{T}_n(x)|_{x^2=0}\equiv T_n(x)|_{x=\tilde{x}}. \tag{4.1.5}$$

Therefore, the harmonic extension establishes an one-to-one relation between homogeneous polynomials on the light-cone and corresponding harmonic polynomials off-cone which provide a convenient realization for symmetric tensor representations of the Lorentz group and the same is valid for tensor polynomials carrying more complicated tensor representations of the Lorentz group. It is important to note that the unique harmonic extension must not destroy the type of Lorentz representation.

The solutions of

$$\Box\overset{\circ}{T}_n(x)=0, \tag{4.1.6}$$

are *(scalar) harmonic polynomials* of order n corresponding to symmetric traceless tensors of rank n ($\overset{\circ}{T}_n(x)\equiv T_n^{(n)}(x)$). They are given by (see, e.g., [121], Chapter IX) and [19])

$$\overset{\circ}{T}_n(x)=\mathcal{P}_n^{(n)}(x)T_n(x), \tag{4.1.7}$$

with the harmonic projection operator

$$\mathcal{P}_n^{(n)}(x)\equiv H_n^{(2h)}(x^2|\Box)=\sum_{k=0}^{[\frac{n}{2}]}\frac{(-1)^k(h+n-k-2)!}{4^kk!(h+n-2)!}\,x^{2k}\,\Box^k\,. \tag{4.1.8}$$

Therefore, the unique harmonic extension $\overset{\circ}{T}_n(x)$ of $T_n(\tilde{x})$ is uniquely given by (4.1.7) and (4.1.8). Additionally, the restriction of Eq. (4.1.8) to the light cone is $\mathcal{P}^{(n)}_n(\tilde{x}) = 1$. The condition of tracelessness (4.1.6) is trivially satisfied on the light cone by $\tilde{x}^2 = 0$. For $T_n(\tilde{x}) = (a\tilde{x})^n$ the harmonic extension can be rewritten as

$$\begin{aligned}\overset{\circ}{T}_n(x) &= \frac{2^n n!(2h+n-3)!}{(2h+2n-3)!}\left(\sqrt{a^2x^2}\right)^n P_n^{\left(h-\frac{3}{2},\,h-\frac{3}{2}\right)}\left(\frac{ax}{\sqrt{a^2x^2}}\right)\\ &= \frac{n!}{(h-1)_n}\left(\frac{1}{2}\sqrt{a^2x^2}\right)^n C_n^{h-1}\left(\frac{ax}{\sqrt{a^2x^2}}\right), \qquad (4.1.9)\end{aligned}$$

with $(h-1)_n = \Gamma(h-1+n)/\Gamma(h-1)$. Here $P_n^{\left(h-\frac{3}{2},\,h-\frac{3}{2}\right)}$ are the Jacobi polynomials and C_n^{h-1} are the Gegenbauer polynomials of degree n. Note that in three dimensions $C_n^{1/2}$ coincide with the Legendre polynomials.

A vector polynomial of degree $n-1$ in x for the Young tableau $\begin{array}{|c|c|}\hline \alpha & n-1 \\ \hline\end{array}$ is recovered by straightforward differentiation as

$$\begin{aligned}T^{(n)}_{\alpha n-1}(x) &= \frac{1}{n}\,\partial_\alpha \overset{\circ}{T}_n(x),\\ &= \frac{1}{n}\sum_{k=0}^{\left[\frac{n-1}{2}\right]}\frac{(-1)^k(h+n-k-3)!}{4^k k!(h+n-2)!}\,x^{2k}\left\{(h+n-k-2)\partial_\alpha - \frac{1}{2}\,x_\alpha\Box\right\}\Box^k T_n(x). \qquad (4.1.10)\end{aligned}$$

The conditions of tracelessness are now given by

$$\partial^\alpha T^{(n)}_{\alpha n-1}(x) = 0, \quad \Box T^{(n)}_{\alpha n-1}(x) = 0. \qquad (4.1.11)$$

On the light-cone, the vector polynomial of degree $n-1$ in $\tilde{x}$ is obtained from $k=0$ in Eq. (4.1.10) as follows

$$T^{(n)}_{\alpha n-1}(\tilde{x}) = \frac{1}{n(h+n-2)}\,\mathrm{d}_\alpha T_n(\tilde{x}), \qquad (4.1.12)$$

where the inner derivative d_α, as a lowering operator on the light cone by construction, maps the scalar polynomial of degree n to a vector polynomial of degree $n-1$. The condition of tracelessness on the light-cone can be formulated by the help of the inner derivative:

$$\mathrm{d}^\alpha T^{(n)}_{\alpha n-1}(\tilde{x}) = 0, \qquad (4.1.13)$$

and is easily proved by the property $\mathrm{d}^2 = 0$. The harmonic extension of (4.1.12) is given by

$$T^{(n)}_{\alpha n-1}(x) = \frac{1}{n(h+n-2)}\,\mathcal{P}^{(n)}_{\alpha n-1}(x) T_n(x), \qquad (4.1.14)$$

where the harmonic projection operator of vector polynomials of type $[n] = (n)$ is defined by

$$\begin{aligned}\mathcal{P}^{(n)}_{\alpha n-1}(x) &:= (h+n-2)\partial_\alpha \mathcal{P}^{(n)}_n(x)\\ &= \sum_{k=0}^{\left[\frac{n-1}{2}\right]}\frac{(-1)^k(h+n-k-3)!}{4^k k!(h+n-3)!}\,x^{2k}\mathcal{D}_\alpha(k)\,\Box^k \qquad (4.1.15)\end{aligned}$$

and where the differential operator,

$$\mathcal{D}_\alpha(k) := (h-1+k+x\partial)\partial_\alpha - \frac{1}{2}\,x_\alpha\Box, \tag{4.1.16}$$

can be understood as a formal off-cone modification of the interior differentiation d_α which gives in the limit $k=0$ and $x \to \tilde{x}$ the interior differential operator ($\mathcal{D}_\alpha(0) = \mathrm{d}_\alpha$ and $\mathcal{P}^{(n)}_{\alpha n-1}(\tilde{x}) = \mathrm{d}_\alpha$). The *prescription for the harmonic extension* is: Replace the interior derivative d_α by the differential operator $\mathcal{D}_\alpha(k)$ which has to be sandwiched in the scalar harmonic projection operator of the corresponding polynomial degree between x^{2k} and $\Box^k$ so that in the limit $k=0$ the projector of the Lorentz spin on the light-cone is recovered.

The generalization of (4.1.12) is

$$T^{(n)}_{\mu_1\cdots\mu_l n-l}(\tilde{x}) = \frac{(n-l)!}{n!}\frac{(h+n-2-l)!}{(h+n-2)!}\,\mathrm{d}_{\mu_1}\ldots\mathrm{d}_{\mu_l}\,T_n(\tilde{x}), \tag{4.1.17}$$

and its harmonic extension is given by

$$T^{(n)}_{\mu_1\cdots\mu_l n-l}(x) = \frac{(n-l)!}{n!}\frac{(h+n-2-l)!}{(h+n-2)!}\,\mathcal{P}^{(n)}_{\mu_1\cdots\mu_l n-l}(x)T_n(x), \tag{4.1.18}$$

where

$$\mathcal{P}^{(n)}_{\mu_1\cdots\mu_l n-l}(x) = \sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k(h+n-k-2-l)!}{4^k k!(h+n-2-l)!}\,x^{2k}\mathcal{D}_{\mu_1}(k)\ldots\mathcal{D}_{\mu_l}(k)\,\Box^k. \tag{4.1.19}$$

(ii) Now we consider an irreducible tensor $T^{(n,1)}_{[\alpha\beta]\mu_1\cdots\mu_{n-1}}$ having the symmetry type of the Young tableau $\begin{array}{|c|c|}\hline \beta & n-1 \\ \hline \alpha \\ \cline{1-1}\end{array}$.

In order to obtain an irreducible tensor, it has to be:
(a) traceless in $\mu_1\cdots\mu_{n-1}$;
(b) traceless in one antisymmetric and one symmetric index.
Using the polynomial technique we replace $T^{(n,1)}_{[\alpha\beta]\mu_1\cdots\mu_{n-1}}$ by a homogeneous polynomial of degree $n-1$ in x. Then we can build the tensor polynomial carrying the required symmetry by means of

$$T^{(n,1)}_{[\alpha\beta]n-1}(x) = \frac{2}{n+1}\,\partial_{[\beta}\overset{\circ}{T}_{\alpha]n}(x), \tag{4.1.20}$$

where $\overset{\circ}{T}_{\mu n}(x)$ has to be harmonic and traceless. Let me note that the condition of the correct symmetry type can be formulated as

$$\epsilon^{\alpha\beta\nu\mu_1\ldots\mu_{2h-3}}\partial_\nu T^{(n,1)}_{[\alpha\beta]n-1}(x) = 0. \tag{4.1.21}$$

The conditions of tracelessness for the irreducible tensor polynomial are given by

$$\Box T^{(n,1)}_{[\alpha\beta]n-1}(x) = 0, \quad \partial^\alpha T^{(n,1)}_{[\alpha\beta]n-1}(x) = 0, \tag{4.1.22}$$

and can be reduced to the conditions of tracelessness for $\overset{\circ}{T}_{\mu n}(x)$:

$$\Box\overset{\circ}{T}_{\alpha n}(x) = 0, \quad \partial^\alpha\overset{\circ}{T}_{\alpha n}(x) = 0, \tag{4.1.23}$$

resulting from (a) and and (b), respectively. We make an Ansatz with the special structure of traces of this tensor as follows:

$$\overset{\circ}{T}_{\mu n}(x) = \left\{ \delta_\mu^\nu - a_n x_\mu \partial^\nu - b_n x^2 \partial_\mu \partial^\nu \right\} \mathcal{P}_n^{(n)}(x) T_{\nu n}(x). \tag{4.1.24}$$

From the conditions of tracelessness (4.1.23) we obtain two equations for the coefficients $a_n,\ b_n$:

$$\begin{aligned} 1 &= (2h+n-1)a_n + 2(n-1)b_n, \\ 0 &= a_n + 2(h+n-2)b_n. \end{aligned}$$

The unique solution is:

$$a_n = \frac{h+n-2}{(h+n-1)(2h+n-3)}, \qquad b_n = -\frac{1}{2(h+n-1)(2h+n-3)}.$$

Then the harmonic and traceless vector polynomial is (not irreducible):

$$\begin{aligned} \overset{\circ}{T}_{\mu n}(x) = \Big\{ \delta_\mu^\nu &- \frac{h+n-2}{(h+n-1)(2h+n-3)}\, x_\mu \partial^\nu \\ &+ \frac{1}{2(h+n-1)(2h+n-3)}\, x^2 \partial_\mu \partial^\nu \Big\} \mathcal{P}_n^{(n)}(x) T_{\nu n}(x). \end{aligned} \tag{4.1.25}$$

Finally, using (4.1.20) and (4.1.25) we obtain for the irreducible representation of type $[n+1] = (n,1)$ as tensor polynomial of degree $n-1$:

$$\begin{aligned} T^{(n,1)}_{[\alpha\beta]n-1}(x) &= \frac{2}{n+1} \left\{ \delta^\mu_{[\alpha} \partial_{\beta]} - \frac{1}{2h+n-3}\, x_{[\alpha} \partial_{\beta]} \partial^\mu \right\} \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (h+n-k-2)!}{4^k k! (h+n-2)!}\, x^{2k} \Box^k T_{\mu n}(x), \\ &= \frac{2}{n+1} \sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k (h+n-k-3)!}{4^k k! (h+n-2)!}\, x^{2k} \Big\{ \delta^\mu_{[\alpha} \Big((h+n-k-2)\partial_{\beta]} - \frac{1}{2}\, x_{\beta]} \Box \Big) \\ &\qquad - \frac{1}{2h+n-3}\, x_{[\alpha} \partial_{\beta]} \Big((h+n-k-2)\partial^\mu - \frac{1}{2}\, x^\mu \Box \Big) \Big\} \Box^k T_{\mu n}(x), \end{aligned} \tag{4.1.26}$$

and the irreducible representation of type $(n,1)$ as tensor polynomial of degree $n-2$ is obtained from Eq. (4.1.26) by means of partial differentiation with respect to the index γ

$$\begin{aligned} T^{(n,1)}_{[\alpha\beta]\gamma n-2}(x) &= \frac{1}{n-1}\, \partial_\gamma T^{(n,1)}_{[\alpha\beta]n-1}(x), \\ &= \frac{2}{(n+1)(n-1)} \left\{ \delta^\mu_{[\alpha} \partial_{\beta]} \partial_\gamma - \frac{1}{2h+n-3} \left(g_{\gamma[\alpha} \partial_{\beta]} + x_{[\alpha} \partial_{\beta]} \partial_\gamma \right) \partial^\mu \right\} \\ &\qquad \times \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (h+n-k-2)!}{4^k k! (h+n-2)!}\, x^{2k} \Box^k T_{\mu n}(x), \\ &= \frac{2}{(n+1)(n-1)} \sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k (h+n-k-4)!}{4^k k! (h+n-2)!}\, x^{2k} \Big\{ \delta^\mu_{[\alpha} \Big((h+n-k-3)\partial_{\beta]} - \frac{1}{2}\, x_{\beta]} \Box \Big) \\ &\qquad \times \Big((h+n-k-2)\partial_\gamma - \frac{1}{2}\, x_\gamma \Box \Big) \\ &- \frac{1}{2h+n-3} \Big\{ g_{\gamma[\alpha} \Big((h+n-k-3)\partial_{\beta]} - \frac{1}{2}\, x_{\beta]} \Box \Big) + x_{[\alpha} \partial_{\beta]} \Big((h+n-k-3)\partial_\gamma - \frac{1}{2}\, x_\gamma \Box \Big) \Big\} \\ &\qquad \times \Big((h+n-k-2)\partial^\mu - \frac{1}{2}\, x^\mu \Box \Big) \Big\} \Box^k T_{\mu n}(x), \end{aligned} \tag{4.1.27}$$

The irreducible tensor polynomial of degree $n-1$ on the light-cone is obtained from Eq. (4.1.26) for $k=0$ as

$$T^{(n,1)}_{[\alpha\beta]n-1}(\tilde{x}) = \frac{2}{(n+1)(h+n-2)}\left\{\delta^{\mu}_{[\alpha}\mathrm{d}_{\beta]} + \frac{1}{2h+n-3}\,X_{[\alpha\beta]}\mathrm{d}^{\mu}\right\}T_{\mu n}(\tilde{x}), \tag{4.1.28}$$

where $X_{[\alpha\beta]} = -\tilde{x}_{[\alpha}\tilde{\partial}_{\beta]}$. Now the condition of tracelessness is imposed as

$$\mathrm{d}^{\alpha}T^{(n,1)}_{[\alpha\beta]n-1}(\tilde{x}) = 0, \tag{4.1.29}$$

and can be easily checked by means of the algebraic properties of the interior derivative on the cone. Additionally, the condition of the correct symmetry type is given on the cone by [49]

$$\epsilon^{\alpha\beta\nu\mu_1\ldots\mu_{2h-3}}\mathrm{d}_{\nu}T^{(n,1)}_{[\alpha\beta]n-1}(\tilde{x}) = 0. \tag{4.1.30}$$

The harmonic extension of $T^{(n,1)}_{[\alpha\beta]n-1}(\tilde{x})$ is defined by

$$T^{(n,1)}_{[\alpha\beta]n-1}(x) = \frac{2}{(n+1)(h+n-2)}\,\mathcal{P}^{(n,1);\,\mu}_{[\alpha\beta]n-1}(x)T_{\mu n}(x), \tag{4.1.31}$$

with

$$\mathcal{P}^{(n,1);\,\mu}_{[\alpha\beta]n-1}(x) = \sum_{k=0}^{[\frac{n-1}{2}]}\frac{(-1)^k(h+n-k-3)!}{4^k k!(h+n-3)!}\,x^{2k}\left\{\delta^{\mu}_{[\alpha}\mathcal{D}_{\beta]}(k) + \frac{1}{2h+n-3}\,X_{[\alpha\beta]}\mathcal{D}^{\mu}(k)\right\}\Box^k, \tag{4.1.32}$$

where we used $X_{[\alpha\beta]} = -x_{[\alpha}\partial_{\beta]}$ in the off-cone expression. Eventually, the restriction of the irreducible representation (4.1.27) to the light cone is given by

$$\begin{aligned} T^{(n,1)}_{[\alpha\beta]\gamma n-2}(\tilde{x}) &= \frac{1}{(n-1)(h+n-3)}\,\mathrm{d}_{\gamma}T^{(n,1)}_{[\alpha\beta]n-1}(\tilde{x}) \\ &= \frac{2}{(n+1)(n-1)(h+n-2)(h+n-3)} \\ &\qquad\times\left\{\delta^{\mu}_{[\alpha}\mathrm{d}_{\beta]}\mathrm{d}_{\gamma} - \frac{1}{2h+n-3}\left(g_{\gamma[\alpha}\mathrm{d}_{\beta]} - X_{[\alpha\beta]}\mathrm{d}_{\gamma}\right)\mathrm{d}^{\mu}\right\}T_{\mu n}(\tilde{x}), \end{aligned} \tag{4.1.33}$$

It satisfies the conditions of tracelessness

$$\mathrm{d}^{\alpha}T^{(n,1)}_{[\alpha\beta]\gamma n-2}(\tilde{x}) = 0, \quad \mathrm{d}^{\gamma}T^{(n,1)}_{[\alpha\beta]\gamma n-2}(\tilde{x}) = 0. \tag{4.1.34}$$

It is important to note that the formulas (4.1.26) and (4.1.27) are the unique harmonic extensions (4.1.28) and (4.1.33), respectively. A harmonic projector $\mathcal{P}^{(n,1);\,\mu}_{[\alpha\beta]\gamma n-2}(x)$ can be proven in the same way.

(iii) Now we regard an irreducible tensor $T^{(n,1,1)}_{[\alpha\beta\gamma]\mu_1\cdots\mu_{n-1}}$ having the symmetry type of the Young tableau $\begin{array}{|c|c|}\hline \gamma & n-1 \\ \hline \beta \\ \cline{1-1} \alpha \\ \cline{1-1}\end{array}$. We are using the convention $T_{[\alpha\beta\gamma]} = \frac{1}{6}(T_{\alpha\beta\gamma} - T_{\alpha\gamma\beta} + T_{\beta\gamma\alpha} - T_{\beta\alpha\gamma} + T_{\gamma\alpha\beta} - T_{\gamma\beta\alpha})$.

In order to obtain an irreducible tensor, it has to be:

(a) traceless in $\mu_1\cdots\mu_{n-1}$;

(b) traceless in one antisymmetric and one symmetric index.

Again, by means of the polynomial technique, the tensor $T^{(n,1,1)}_{[\alpha\beta\gamma]\mu_1\cdots\mu_{n-1}}$ can be replaced by a homogeneous polynomial of degree $n-1$ in x. Then we obtain the tensor polynomial carrying the required symmetry as follows

$$\begin{aligned} T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x) &= \frac{3}{n+2}\,\delta^{\mu}_{[\alpha}\delta^{\nu}_{\beta}\partial_{\gamma]}\,\overset{\circ}{T}_{\mu\nu n}(x), \\ &= \frac{1}{n+2}\left(\delta^{\mu}_{\alpha}\delta^{\nu}_{\beta}\partial_{\gamma} + \delta^{\mu}_{\beta}\delta^{\nu}_{\gamma}\partial_{\alpha} + \delta^{\mu}_{\gamma}\delta^{\nu}_{\alpha}\partial_{\beta}\right)\overset{\circ}{T}_{[\mu\nu]n}(x), \end{aligned} \tag{4.1.35}$$

where $\overset{\circ}{T}_{[\mu\nu]n}(x)$ has to be traceless. The (symmetry) condition of the Young tableau type is given, by means of the polynomial technique, by

$$\epsilon^{\alpha\beta\gamma\nu\mu_1\ldots\mu_{2h-4}}\partial_\nu T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x) = 0. \tag{4.1.36}$$

The conditions of tracelessness for the irreducible tensor polynomial are given by

$$\Box T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x) = 0, \quad \partial^\alpha T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x) = 0, \tag{4.1.37}$$

and give the following conditions of tracelessness for $\overset{\circ}{T}_{[\alpha\beta]n}(x)$:

$$\Box \overset{\circ}{T}_{[\alpha\beta]n}(x) = 0, \quad \partial^\alpha \overset{\circ}{T}_{[\alpha\beta]n}(x) = 0. \tag{4.1.38}$$

Using the same procedure as before we may construct the harmonic skew tensor polynomial $\overset{\circ}{T}_{[\mu\nu]n}(x)$. The solutions of Eqs. (4.1.38) are the *antisymmetric harmonic tensor polynomials* $\overset{\circ}{T}_{[\alpha\beta]n}(x)$ of order n. The most general Ansatz for it has the following structure:

$$\overset{\circ}{T}_{[\alpha\beta]n} = \left\{\delta^{\mu}_{\alpha}\delta^{\nu}_{\beta} - a_n x_{[\alpha}\delta^{\nu}_{\beta]}\partial^{\mu} - b_n x^2 \partial^{\mu}\partial_{[\alpha}\delta^{\nu}_{\beta]} - c_n x^{\mu} x_{[\alpha}\partial_{\beta]}\partial^{\nu}\right\}\mathcal{P}^{(n)}(x) T_{[\mu\nu]n}(x). \tag{4.1.39}$$

The corresponding system of linear equations reads

$$\begin{aligned} 1 &= (2h+n-2)a_n/2 + (n-1)b_n \\ 0 &= 2b_n + (2h+n-2)c_n \\ 0 &= a_n + c_n + 2(h+n-2)b_n\,. \end{aligned}$$

It is solved by the following values:

$$(a_n + c_n,\ b_n,\ c_n) = \frac{1}{(h+n-1)(2h+n-4)}\left(2(h+n-2),\ -1,\ \frac{2}{2h+n-2}\right).$$

The harmonic and traceless skew tensor polynomial is given by

$$\begin{aligned} \overset{\circ}{T}_{[\alpha\beta]n}(x) = \Big\{\delta^{\mu}_{[\alpha}\delta^{\nu}_{\beta]} &+ \frac{2}{(h+n-1)(2h+n-4)}\left((h+n-2)x_{[\alpha}\delta^{\mu}_{\beta]}\partial^{\nu} - \frac{1}{2}\,x^2\partial_{[\alpha}\delta^{\mu}_{\beta]}\partial^{\nu}\right) \\ &- \frac{2}{(h+n-1)(2h+n-4)(2h+n-2)}\,x_{[\alpha}\partial_{\beta]}x^{\mu}\partial^{\nu}\Big\}\mathcal{P}^{(n)}_{n}\left(x^2|\Box\right) T_{[\mu\nu]n}(x). \end{aligned} \tag{4.1.40}$$

Finally, we obtain for the irreducible tensor polynomial of degree $n-1$ of Young tableau $[n+2] = (n,1,1)$:

$$T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x) = \frac{3}{n+2}\Big\{\delta^\mu_{[\alpha}\delta^\nu_\beta\partial_{\gamma]} - \frac{2}{2h+n-4}\, x_{[\alpha}\partial_\beta\delta^\mu_{\gamma]}\partial^\nu\Big\} \times \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k(h+n-k-2)!}{4^k k!(h+n-2)!}\, x^{2k}\Box^k T_{[\mu\nu]n}(x), \tag{4.1.41}$$

$$= \frac{3}{n+2}\sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k(h+n-k-3)!}{4^k k!(h+n-2)!}\, x^{2k}\Big\{\delta^\mu_{[\alpha}\delta^\nu_\beta\Big((h+n-k-2)\partial_{\gamma]} - \frac{1}{2}\, x_{\gamma]}\Box\Big) - \frac{2}{2h+n-4}\, x_{[\alpha}\partial_\beta\delta^\mu_{\gamma]}\Big((h+n-k-2)\partial^\nu - \frac{1}{2}\, x^\nu\Box\Big)\Big\}\Box^k T_{[\mu\nu]n}(x),$$

and for its restriction to the light cone:

$$T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(\tilde{x}) = \frac{3}{(n+2)(h-n-2)}\Big\{\delta^\mu_{[\alpha}\delta^\nu_\beta \mathrm{d}_{\gamma]} + \frac{2}{2h+n-4}\, X_{[\alpha\beta}\delta^\mu_{\gamma]}\mathrm{d}^\nu\Big\}T_{[\mu\nu]n}(\tilde{x}). \tag{4.1.42}$$

Eq. (4.1.41) is the unique harmonic extension of (4.1.42). On the light cone, the condition of tracelessness is given by

$$\mathrm{d}^\alpha T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(\tilde{x}) = 0, \tag{4.1.43}$$

and the condition of the symmetry type reads

$$\epsilon^{\alpha\beta\gamma\nu\mu_1\ldots\mu_{2h-4}}\mathrm{d}_\nu T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(\tilde{x}) = 0. \tag{4.1.44}$$

The harmonic extension of $T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(\tilde{x})$ is defined by

$$T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x) = \frac{3}{(n+2)(h+n-2)}\,\mathcal{P}^{(n,1,1);\,\mu\nu}_{[\alpha\beta\gamma]n-1}(x)T_{[\mu\nu]n}(x), \tag{4.1.45}$$

with

$$\mathcal{P}^{(n,1,1);\,\mu\nu}_{[\alpha\beta\gamma]n-1}(x) = \sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k(h+n-k-3)!}{4^k k!(h+n-3)!}\, x^{2k}\Big\{\delta^\mu_{[\alpha}\delta^\nu_\beta\mathcal{D}_{\gamma]}(k) + \frac{2}{2h+n-4}\, X_{[\alpha\beta}\delta^\mu_{\gamma]}\mathcal{D}^\nu(k)\Big\}\Box^k. \tag{4.1.46}$$

By means of partial differentiation with respect to the index δ, one can also compute the tensor polynomial: $T^{(n,1,1)}_{[\alpha\beta\gamma]\delta n-2}(x) = \frac{1}{n-1}\,\partial_\delta T^{(n,1,1)}_{[\alpha\beta\gamma]n-1}(x)$.

(iv) Now we want to determine the reducible *symmetric harmonic tensor polynomials* $\overset{\circ}{T}_{(\alpha\beta)n}(x)$ which we need for the Young tableau $[n+2] = (n,2)$. We renounce to construct the proper irreducible tensor polynomial,

$$T^{(n,2)}_{[\alpha\rho][\beta\tau]n-2}(x) = \frac{8}{(n+1)n}\,\delta^\mu_{[\alpha}\partial_{\lambda]}\delta^\nu_{[\beta}\partial_{\gamma]}\delta^\lambda_{(\rho}\delta^\gamma_{\tau)}\overset{\circ}{T}_{(\mu\nu)n}(x), \tag{4.1.47}$$

which has the symmetry type of the Young tableau $\begin{array}{|c|c|c|}\hline \rho & \tau & n-2 \\ \hline \alpha & \beta \\ \cline{1-2}\end{array}$. In Section 4.2.4 we will consider the more easier tensor polynomial $T^{(n,2)}_{(\alpha\beta)n}(x) = x^\rho x^\tau \, T^{(n,2)}_{[\alpha\rho][\beta\tau]n-2}(x)$ (see Eq. (4.2.72)).
The conditions of tracelessness for the reducible tensor polynomial are:

$$\Box \overset{\circ}{T}_{(\alpha\beta)n}(x) = 0, \quad \partial^\alpha \overset{\circ}{T}_{(\alpha\beta)n}(x) = 0, \quad g^{\alpha\beta} \overset{\circ}{T}_{(\alpha\beta)n}(x) = 0. \tag{4.1.48}$$

Its general structure is as follows:

$$\begin{aligned} \overset{\circ}{T}_{(\alpha\beta)n}(x) = \Big\{ & \delta^\mu_\alpha \delta^\nu_\beta + a_n g_{\alpha\beta} x^{(\nu} \partial^{\mu)} + b_n x_{(\alpha} \delta^{(\nu}_{\beta)} \partial^{\mu)} + c_n x_{(\alpha} x^{(\nu} \partial^{\mu)} \partial_{\beta)} \\ & + d_n x^2 \delta^{(\nu}_{(\alpha} \partial_{\beta)} \partial^{\mu)} + e_n x^2 x^{(\nu} \partial^{\mu)} \partial_\alpha \partial_\beta + f_n x^2 x_{(\alpha} \partial_{\beta)} \partial^\mu \partial^\nu \\ & + g_n x_\alpha x_\beta \partial^\mu \partial^\nu + k_n x^2 g_{\alpha\beta} \partial^\mu \partial^\nu + l_n x^4 \partial_\alpha \partial_\beta \partial^\mu \partial^\nu \Big\} \breve{T}_{(\mu\nu)n}(x) \end{aligned} \tag{4.1.49}$$

with the partially traceless polynomials

$$\breve{T}_{(\mu\nu)n}(x) = \left(\delta^\rho_\mu \delta^\sigma_\nu - \frac{1}{2h} \, g_{\mu\nu} g^{\rho\sigma} \right) \mathcal{P}^{(n)}_n(x) T_{(\rho\sigma)n}(x). \tag{4.1.50}$$

It holds by construction

$$\Box \breve{T}_{(\mu\nu)n}(x) = 0, \qquad g^{\mu\nu} \breve{T}_{(\mu\nu)n}(x) = 0. \tag{4.1.51}$$

From Eqs. (4.1.48) the following system of linear equations for the nine unknown coefficients results:

$$\begin{aligned} 0 &= (n-1)c_n + b_n + 2h a_n \\ 0 &= d_n + g_n + (n-2) f_n + 2h k_n \\ 0 &= 2(h+n-2) d_n + c_n + b_n \\ 0 &= 2(h+n-2) e_n + c_n \\ 0 &= 2(h+n-2) f_n + c_n + 2 g_n \\ 0 &= 2(h+n-2) k_n + a_n + g_n \\ 0 &= 4(h+n-3) l_n + f_n + e_n \\ 0 &= (2h+n) b_n + 2(n-1) d_n + 2 a_n + 2 \\ 0 &= (2h+n) c_n + 2 d_n + 2 a_n + 4(n-2) e_n \\ 0 &= (2h+n) f_n + 2 k_n + d_n + 8(n-3) l_n + 2 e_n \\ 0 &= 2(2h+n-1) g_n + 4 k_n + c_n + 2(n-2) f_n + b_n. \end{aligned}$$

Its unique solution is

$$
\begin{aligned}
a_n &= \frac{h+n-2}{(h-1)(h+n)(2h+n-2)}, \\
b_n &= -2\,\frac{(h(h+n-1)-n+1)(h+n-2)}{(h-1)(h+n)(h+n-1)(2h+n-2)}, \\
c_n &= -2\,\frac{h+n-2}{(h-1)(h+n)(h+n-1)(2h+n-2)}, \\
d_n &= \frac{h(h+n-1)-n+2}{(h-1)(h+n)(h+n-1)(2h+n-2)}, \\
e_n &= \frac{1}{(h-1)(h+n)(h+n-1)(2h+n-2)}, \\
f_n &= -\frac{h(h+n-4)-2(n-3)}{(h-1)(h+n)(h+n-1)(2h+n-2)(2h+n-3)}, \\
g_n &= \frac{\big(h(h+n-2)-n+3\big)(h+n-2)}{(h-1)(h+n)(h+n-1)(2h+n-2)(2h+n-3)}, \\
k_n &= -\frac{h(h+n-3)-2n+3+\frac{1}{2}(h+n)(h+n-1)}{(h-1)(h+n)(h+n-1)(2h+n-2)(2h+n-3)}, \\
l_n &= \frac{h-3}{4(h-1)(h+n)(h+n-1)(2h+n-2)(2h+n-3)}.
\end{aligned}
$$

In the case $h = 2$ these coefficients simplify; they are given as follows:

$$
\begin{aligned}
a_n &= \frac{n}{(n+2)^2} & b_n &= -2\,\frac{n(n+3)}{(n+1)(n+2)^2}, \\
c_n &= -2\,\frac{n}{(n+1)(n+2)^2}, & d_n &= \frac{4+n}{(n+1)(n+2)^2}, \\
e_n &= \frac{1}{(n+1)(n+2)^2}, & f_n &= -2\,\frac{1}{(n+1)^2(n+2)^2}, \\
g_n &= \frac{n(n+3)}{(n+1)^2(n+2)^2}, & k_n &= -\frac{1}{2}\frac{n^2+3n+4}{(n+1)^2(n+2)^2}, \\
l_n &= -\frac{1}{4}\frac{1}{(n+1)^2(n+2)^2}\,.
\end{aligned}
$$

This finishes the determination of the harmonic tensor polynomials being necessary for the present study. In principle, the extension of the procedure to arbitrary tensors of higher order is obvious. However, the explicit computation, without any additional information about their general properties, will be quite complicated. Let us remark that, to the best of our knowledge, such quantities have not been considered in the mathematical literature.

4.2 Twist decomposition of a bilocal 2nd rank tensor operator

This Section is devoted to the twist decomposition of a general 2nd rank light-cone tensor operator. Since the twist decomposition is independent from the chirality we demonstrate it only for the chiral-even gluon operator (2.1.33). The twist decomposition of the chiral-odd gluon operator Eq. (2.1.34) is the same. In addition, we make the special choice $\kappa_1 = 0, \kappa_2 = \kappa$, thereby also simplifying the generalized covariant derivatives, Eq. (2.2.4), to the usual ones:

$$G_{\alpha\beta}(0,\kappa x) = F_{\alpha}{}^{\rho}(0)U(0,\kappa x)F_{\beta\rho}(\kappa x) = \sum_{n=0}^{\infty} \frac{\kappa^n}{n!} x^{\mu_1} \ldots x^{\mu_n} G_{\alpha\beta(\mu_1\ldots\mu_n)} \ , \tag{4.2.1}$$

$$\text{with} \qquad G_{\alpha\beta(\mu_1\ldots\mu_n)} = F_{\alpha}{}^{\rho}(y) D^y_{(\mu_1} \ldots D^y_{\mu_n)} F_{\beta\rho}(y)\big|_{y=0} \ ;$$

here the symbol $(\ldots)$ denotes symmetrization of the enclosed indices.

Physically, the forward matrix elements of the both gluon operators provide the gluon distribution functions [43, 103, 81].

The local tensor operators $G_{\alpha\beta(\mu_1\ldots\mu_n)}$ can be decomposed according to the Young patterns (i) to (iv) and possible higher ones (i.e., for $m_1 + m_2 = n+2, m_2 \geq 3$):

$$G_{\alpha\beta(\mu_1\ldots\mu_n)} = G^{(\mathrm{i})}_{\alpha\beta(\mu_1\ldots\mu_n)} + \alpha_{n+1} G^{(\mathrm{ii})}_{\alpha\beta(\mu_1\ldots\mu_n)} + \beta_n G^{(\mathrm{iii})}_{\alpha\beta(\mu_1\ldots\mu_n)} + \gamma_n G^{(\mathrm{iv})}_{\alpha\beta(\mu_1\ldots\mu_n)} + \ldots \ , \tag{4.2.2}$$

with the (nontrivial) normalization coefficients $f_{[m]}/m!$ of the Young operators given by $\alpha_{n+1} = 2(n+1)/(n+2)$ for $[m] = (n+1,1)$, $\beta_n = 3n/(n+2)$ for $[m] = (n,1,1)$ and $\gamma_n = 4(n-1)/(n+1)$ for $[m] = (n,2)$. The corresponding Clebsch-Gordan series in terms of representations (j_1, j_2) of the Lorentz group is

$$\begin{aligned}
&\left(\tfrac{1}{2},\tfrac{1}{2}\right) \otimes \left(\tfrac{1}{2},\tfrac{1}{2}\right) \otimes \left(\left(\tfrac{n}{2},\tfrac{n}{2}\right) \oplus \left(\tfrac{n-2}{2},\tfrac{n-2}{2}\right) \oplus \ldots\right) \\
&\quad = \left(\tfrac{n+2}{2},\tfrac{n+2}{2}\right) \oplus \left(\left(\tfrac{n+2}{2},\tfrac{n}{2}\right) \oplus \left(\tfrac{n}{2},\tfrac{n+2}{2}\right)\right) \oplus \left(\tfrac{n}{2},\tfrac{n}{2}\right) \\
&\qquad \oplus \left(\left(\tfrac{n+2}{2},\tfrac{n-2}{2}\right) \oplus \left(\tfrac{n-2}{2},\tfrac{n+2}{2}\right)\right) \\
&\qquad \oplus \left(\left(\tfrac{n}{2},\tfrac{n-2}{2}\right) \oplus \left(\tfrac{n-2}{2},\tfrac{n}{2}\right)\right) \oplus \left(\tfrac{n-2}{2},\tfrac{n-2}{2}\right) \oplus \ldots \ ;
\end{aligned} \tag{4.2.3}$$

the corresponding tensor spaces will be denoted by $\mathbf{T}(j_1, j_2)$. In the last line of Eq. (4.2.3) such representations are listed down which correspond to higher twist contributions contained in the trace terms of symmetry classes (i) – (iv). The canonical (or scale) dimension d of the local operator (4.2.2) is $n+4$, and the spin of the various contributions in (4.2.3) ranges from $n+2$ up to 1 or 0 if n is even or odd, respectively; therefore, the local operators with well-defined twist are irreducible tensors of the Lorentz group or, equivalently, of the group $SL(2,\mathbb{C}) \times \mathbb{R}_+$, having scale dimension d.

In general, according to the spin content and the rank of the corresponding local tensor operators, the nonlocal operator (4.2.1) for arbitrary x contains contributions of twist ranging from $\tau = 2$ until $\tau = \infty$. However, after projection onto the light–cone, $x \to \tilde{x}$, this infinite series terminates at least at $\tau = 6$ since higher order terms are proportional to x^2.

In order to be more explicit, as well as for later use, we introduce the (anti)symmetrization with respect to α and β and we define the following nonlocal operators:

$$G^{\pm}_{\alpha\beta}(0,\kappa x) \ := \ \tfrac{1}{2}\left(G_{\alpha\beta}(0,\kappa x) \pm G_{\beta\alpha}(0,\kappa x)\right) \ , \tag{4.2.4}$$

$$G^{\pm}_{\alpha}(0,\kappa x) \ := \ x^{\beta}\, G^{\pm}_{\alpha\beta}(0,\kappa x) \equiv G^{\pm}_{\alpha\bullet}(0,\kappa x) \ , \tag{4.2.5}$$

$$G(0,\kappa x) \ := \ x^{\alpha} x^{\beta}\, G^{+}_{\alpha\beta}(0,\kappa x) \ . \tag{4.2.6}$$

In fact, the twist decomposition of these bilocal light–ray operators reads:[1]

$$\begin{aligned}
G^{+}_{\alpha\beta}(0,\kappa\tilde{x}) =&\ G^{\rm tw2}_{(\alpha\beta)}(0,\kappa\tilde{x}) + G^{\rm tw3}_{(\alpha\beta)}(0,\kappa\tilde{x}) + G^{\rm tw4}_{(\alpha\beta)}(0,\kappa\tilde{x}) \\
&+ G^{\rm tw5}_{(\alpha\beta)}(0,\kappa\tilde{x}) + G^{\rm tw6}_{(\alpha\beta)}(0,\kappa\tilde{x})\,, \qquad (4.2.7)\\
G^{-}_{\alpha\beta}(0,\kappa\tilde{x}) =&\ G^{\rm tw3}_{[\alpha\beta]}(0,\kappa\tilde{x}) + G^{\rm tw4}_{[\alpha\beta]}(0,\kappa\tilde{x}) + G^{\rm tw5}_{[\alpha\beta]}(0,\kappa\tilde{x})\,, \qquad (4.2.8)\\
G^{+}_{\alpha}(0,\kappa\tilde{x}) =&\ G^{\rm tw2}_{(\alpha\bullet)}(0,\kappa\tilde{x}) + G^{\rm tw3}_{(\alpha\bullet)}(0,\kappa\tilde{x}) + G^{\rm tw4}_{(\alpha\bullet)}(0,\kappa\tilde{x})\,, \qquad (4.2.9)\\
G^{-}_{\alpha}(0,\kappa\tilde{x}) =&\ G^{\rm tw3}_{[\alpha\bullet]}(0,\kappa\tilde{x}) + G^{\rm tw4}_{[\alpha\bullet]}(0,\kappa\tilde{x})\,, \qquad (4.2.10)\\
G(0,\kappa\tilde{x}) =&\ G^{\rm tw2}(0,\kappa\tilde{x})\,. \qquad (4.2.11)
\end{aligned}$$

The various twist contributions individually decompose further according to the symmetry classes which may contribute. The explicit expressions for generic tensor operators are given in the following subsections, where it will be shown that Young patterns (i) and (ii) as well as (iv) contribute to the symmetric tensor operators and related vector and scalar operators, and Young patterns (ii) and (iii) contribute to the antisymmetric tensor operators and related vector operators.

Fortunately, in the case of forward scattering one has to consider the symmetry type (i) in the polarized and unpolarized case, and additionally, the symmetry type (ii) for the polarized case. But in order to consider non-forward matrix elements all symmetry types are involved.

4.2.1 Tensor operators of symmetry class (i)

Construction of nonlocal symmetric class-(i) operators of definite twist

Let us start with the simplest case of the totally symmetric traceless tensors, and their contractions with x, which have twist $\tau = 2$ and are contained in the tensor space $\mathbf{T}(\frac{n+2}{2}, \frac{n+2}{2})$. They have symmetry class (i) and are uniquely characterized by the following standard tableau (with normalizing factor 1):

$$\begin{array}{|c|c|c|c|c|}\hline \alpha & \beta & \mu_1 & \quad\cdots\quad & \mu_n \\ \hline \end{array} \qquad \stackrel{\wedge}{=} \qquad \mathop{\mathcal{S}}_{\alpha\beta\mu_1\ldots\mu_n} F_{\alpha}{}^{\rho}(0) D_{\mu_1}\ldots D_{\mu_n} F_{\beta\rho}(0) - \text{trace terms}\ ,$$

The irreducible local twist–2 operator reads

$$G^{\rm tw2(i)}_{(\alpha\beta\mu_1\ldots\mu_n)} = F^{\ \rho}_{(\alpha}(0) D_{\mu_1}\ldots D_{\mu_n} F_{\beta)\rho}(0) - \text{trace terms}\,.$$

Let us postpone the determination of the trace terms and make the resummation to the corresponding nonlocal operator in advance. This is obtained, at first, by contracting with $x^{\mu_1}\ldots x^{\mu_n}$ and rewriting in the following form:

$$G^{\rm tw2(i)}_{(\alpha\beta)n}(x) = x^{\mu_1}\ldots x^{\mu_n}\ G^{\rm tw2(i)}_{(\alpha\beta\mu_1\ldots\mu_n)} = \frac{1}{(n+2)(n+1)}\,\partial_\alpha\partial_\beta \overset{\circ}{G}_{n+2}(x), \qquad (4.2.12)$$

with

$$\begin{aligned}
\overset{\circ}{G}_{n+2}(x) &= G_{n+2}(x) - \text{ trace terms}\ ,\\
G_{n+2}(x) &\equiv x^{\mu} x^{\nu} F_{\mu}{}^{\rho}(0)(xD)^{n} F_{\nu\rho}(0)\ .
\end{aligned}$$

---

[1] We use the notation $A_{(\alpha\beta)} \equiv \frac{1}{2}(A_{\alpha\beta} + A_{\beta\alpha})$ and $A_{[\alpha\beta]} \equiv \frac{1}{2}(A_{\alpha\beta} - A_{\beta\alpha})$.

Here, $\overset{\circ}{G}_{n+2}(x)$ is a harmonic polynomial of order $n+2$, cf. Eq. (4.1.7):

$$\overset{\circ}{G}_{n+2}(x) = H^{(4)}_{n+2}\big(x^2|\Box\big)\, G_{n+2}(x) \equiv \sum_{k=0}^{[\frac{n+2}{2}]} \frac{(n+2-k)!}{(n+2)!k!}\left(\frac{-x^2}{4}\right)^k \Box^k G_{n+2}(x)\,.$$

Then, using $((n+2)(n+1))^{-1} = \int_0^1 \mathrm{d}\lambda (1-\lambda)\lambda^n$ and the integral representation of Euler's beta function we obtain the nonlocal *twist–2 tensor operator*:[2]

$$G^{\rm tw2}_{(\alpha\beta)}(0,\kappa x) = \sum_{n=0}^{\infty} \frac{\kappa^n}{n!}\, G^{\rm tw2(i)}_{(\alpha\beta)n}(x) = \partial_\alpha \partial_\beta \int_0^1 \mathrm{d}\lambda (1-\lambda)\, \overset{\circ}{G}(0,\kappa\lambda x), \tag{4.2.13}$$

with

$$\overset{\circ}{G}(0,\kappa x) = G(0,\kappa x) + \sum_{k=1}^{\infty} \int_0^1 \mathrm{d}t\, t \left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{k!(k-1)!} \left(\frac{1-t}{t}\right)^{k-1} G(0,\kappa t x). \tag{4.2.14}$$

The operator $G^{\rm tw2}_{(\alpha\beta)}(0,\kappa x)$ satisfies the conditions of a harmonic tensor function:

$$g^{\alpha\beta} G^{\rm tw2}_{(\alpha\beta)}(0,\kappa x) = 0, \quad \Box G^{\rm tw2}_{(\alpha\beta)}(0,\kappa x) = 0, \quad \partial^\alpha G^{\rm tw2}_{(\alpha\beta)}(0,\kappa x) = 0\,. \tag{4.2.15}$$

Reduction to vector and scalar operators

Now, we specify to the nonlocal vector and scalar operators. Multiplying Eq. (4.2.13) by x^β and observing the equality $(x\partial_x)\overset{\circ}{G}(0,\kappa\lambda x) = (\lambda\partial_\lambda + 2)\overset{\circ}{G}(0,\kappa\lambda x)$ gives the *twist–2 vector operator*,

$$G^{\rm tw2}_{(\alpha\bullet)}(0,\kappa x) = \partial_\alpha \int_0^1 \mathrm{d}\lambda\, \lambda\, \overset{\circ}{G}(0,\kappa\lambda x), \tag{4.2.16}$$

which satisfies the conditions

$$\Box\; G^{\rm tw2}_{(\alpha\bullet)}(0,\kappa x) = 0, \quad \partial^\alpha\; G^{\rm tw2}_{(\alpha\bullet)}(0,\kappa x) = 0\,. \tag{4.2.17}$$

Multiplying by $x^\alpha x^\beta$ gives the *twist–2 scalar operator*,

$$G^{\rm tw2}(0,\kappa x) = \overset{\circ}{G}(0,\kappa x)\,, \tag{4.2.18}$$

which, by definition, satisfies the condition

$$\Box G^{\rm tw2}(0,\kappa x) = 0\,. \tag{4.2.19}$$

The expression (4.2.14) for the scalar operator has already been given by Balitsky and Braun [9].

Comparing Eqs. (4.2.13), (4.2.16) and (4.2.18) we may recognize that in the case of symmetry class (i) the tensor and vector operators are obtained from the scalar operator by very simple operations. Furthermore, we observe how in the case of the scalar operator the trace terms – being proportional to x^2 – are to be subtracted from $G(0,\kappa x)$ in order to make that operator traceless. In the case of vector and tensor operators such subtraction, because of the appearance of the derivatives, is more complicated.

---

[2] Here, and in the following, we omit the indication of the symmetry class of the nonlocal twist–2 operators because only totally symmetric tensors contribute. However, the trace terms being of higher twist must be classified according to their symmetry type.

Projection onto the light–cone

Let us now project onto the light–cone and, at the same time, also extend to the case of general values (κ_1, κ_2). Because of the derivatives ∂_α and ∂_β, appearing in Eq. (4.2.13), only the terms with $k = 1, 2$ in Eq. (4.2.14) contribute. The final expression for the *symmetric twist–2 light–cone tensor operator* is given by

$$G^{\rm tw2}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = \partial_\alpha\partial_\beta \int_0^1 {\rm d}\lambda(1-\lambda) G(\kappa_1\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{>{\rm (i)}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \tag{4.2.20}$$

$$\begin{aligned} G^{>{\rm (i)}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = &\int_0^1 {\rm d}\lambda \Big\{(1-\lambda+\lambda\ln\lambda)\left(\tfrac{1}{2} g_{\alpha\beta} + x_{(\alpha}\partial_{\beta)}\right)\Box \\ &+ \tfrac{1}{4}\big(2(1-\lambda) + (1+\lambda)\ln\lambda\big) x_\alpha x_\beta \Box^2\Big\} G(\kappa_1\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}. \end{aligned} \tag{4.2.21}$$

The operator $G^{>{\rm (i)}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x})$ contains the *higher twist contributions* which have to be subtracted from the first term, Eq. (4.2.20), in order to make the whole expression traceless. In order to disentangle the different terms of well-defined twist we observe that the tensor operator (4.2.21) consists of scalar and vector parts. Here, an operator being contained in the trace terms is called a scalar and vector part, if the expression multiplied by $g_{\alpha\beta}$, x_α or x_β is a scalar, like $\Box G(\kappa_1 x, \kappa_2 x)$ or $\partial^\mu G_\mu(\kappa_1 x, \kappa_2 x)$, and a vector, like $\partial_\beta G(\kappa_1 x, \kappa_2 x)$ or $\partial_\beta \partial^\mu G_\mu(\kappa_1 x, \kappa_2 x)$, respectively.[3] The scalar operators, $\Box G(\kappa_1 x, \kappa_2 x)|_{x=\tilde{x}}$ and $\Box^2 G(\kappa_1 x, \kappa_2 x)|_{x=\tilde{x}}$, occurring in Eq. (4.2.21) have already well-defined twist $\tau = 4$ and $\tau = 6$, respectively. To ensure that the vector operator $\partial_\beta \Box G(\kappa_1 x, \kappa_2 x)|_{x=\tilde{x}}$ also obtains well-defined twist, it is necessary to subtract its own trace terms which are of twist $\tau = 6$.

The higher twist operators contained in the trace terms of the twist–2 tensor operator are the following:

$$\begin{aligned} G^{>{\rm (i)}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = &\, G^{\rm tw4(i)a}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + G^{\rm tw4(i)b}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) \\ &+ G^{\rm tw6(i)a}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) + G^{\rm tw6(i)b}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \end{aligned} \tag{4.2.22}$$

with

$$\begin{aligned} G^{\rm tw4(i)a}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) &= \tfrac{1}{2}\, g_{\alpha\beta} \Box \int_0^1 {\rm d}\lambda (1-\lambda+\lambda\ln\lambda) G(\kappa_1\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\ G^{\rm tw4(i)b}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) &= x_{(\alpha}\partial_{\beta)} \Box \int_0^1 {\rm d}\lambda (1-\lambda+\lambda\ln\lambda) G(\kappa_1\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{\rm tw6(i)b}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}), \end{aligned}$$

and

$$\begin{aligned} G^{\rm tw6(i)a}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) &= \tfrac{1}{4}\, x_\alpha x_\beta \Box^2 \int_0^1 {\rm d}\lambda \big(2(1-\lambda) + (1+\lambda)\ln\lambda\big) G(\kappa_1\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\ G^{\rm tw6(i)b}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) &= \tfrac{1}{4}\, x_\alpha x_\beta \Box^2 \int_0^1 {\rm d}\lambda \Big(\tfrac{(1-\lambda)^2}{\lambda} - \tfrac{1-\lambda^2}{2\lambda} - \lambda\ln\lambda\Big) G(\kappa_1\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \end{aligned}$$

Obviously, by inspection of the spin content of the symmetry type (i), the local twist–4 and twist–6 operators carry the representation $\mathbf{T}(\frac{n}{2}, \frac{n}{2})$ and $\mathbf{T}(\frac{n-2}{2}, \frac{n-2}{2})$, respectively.

---

[3] Strictly speaking, x_α and x_β as well as $g_{\alpha\beta}$ which are necessary for the dimension and tensor structure of the whole expression do not belong to the higher twist operator itself.

Finally, completing the twist decomposition of the symmetric nonlocal tensor operator, we write down the symmetric twist–4 light–cone tensor operator contained in the expression (4.2.21)

$$\begin{aligned} G^{\mathrm{tw4(i)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= G^{\mathrm{tw4(i)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(i)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\ &= \int_0^1 \mathrm{d}\lambda \Big\{ (1-\lambda+\lambda\ln\lambda)\left(\tfrac{1}{2}\,g_{\alpha\beta} + x_{(\alpha}\partial_{\beta)}\right)\Box \\ &\qquad - \tfrac{1}{4}\Big(\tfrac{(1-\lambda)^2}{\lambda} - \tfrac{1-\lambda^2}{2\lambda} - \lambda\ln\lambda\Big) x_\alpha x_\beta \Box^2 \Big\}\, G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \end{aligned} \tag{4.2.23}$$

whereas the symmetric twist–6 light–cone operator contained in (4.2.21) reads

$$\begin{aligned} G^{\mathrm{tw6(i)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= G^{\mathrm{tw6(i)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(i)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\ &= \tfrac{1}{4}\, x_\alpha x_\beta \Box^2 \int_0^1 \mathrm{d}\lambda \Big(\tfrac{1-\lambda^2}{2\lambda} + \ln\lambda\Big)\, G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \end{aligned} \tag{4.2.24}$$

Vector and scalar light–ray operators

Contracting Eq. (4.2.20) with $\tilde{x}^\beta$, making use of formula $(x\partial_x)f(\lambda x) = \lambda\partial_\lambda f(\lambda x)$ and performing the partial integrations we obtain the final version of the *twist–2 light–cone vector operator*,

$$G^{\mathrm{tw2}}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\,\lambda \Big[\partial_\alpha + \tfrac{1}{2}(\ln\lambda) x_\alpha \Box\Big] G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \tag{4.2.25}$$

and the twist–4 light–cone vector operator,

$$G^{\mathrm{tw4(i)}}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = -\tfrac{1}{2}\, x_\alpha \Box \int_0^1 \mathrm{d}\lambda\, \lambda(\ln\lambda) G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \tag{4.2.26}$$

In order to obtain the *scalar twist–2 light–ray operator* we multiply (4.2.25) by $\tilde{x}^\alpha$. Then, the twist–4 part vanishes and the remaining twist–2 operator restores the scalar operator (compare Eq. (4.2.11)),

$$G^{\mathrm{tw2}}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G(\kappa_1\tilde{x},\kappa_2\tilde{x})\,. \tag{4.2.27}$$

Let us point to the fact that the trace of the original gluon tensor, $g^{\alpha\beta}G_{\alpha\beta}(\kappa_1\tilde{x},\kappa_2\tilde{x})$, is a twist–4 scalar operator. It is contained in $G^{\mathrm{tw4(i)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ as well as similar expressions occurring below, cf. also Eq. (4.2.85).

Condition of tracelessness on the cone

Finally, it should be remarked, that the conditions (4.2.15), (4.2.17) and (4.2.19), if translated into the corresponding ones containing derivatives with respect to $\tilde{x}$, no longer hold for the light–cone operators. This is clear because, by projecting onto the light–cone, part of the original structure of the operators has been lost. Nevertheless, the conditions of tracelessness of the light–cone operators may be formulated by using the *interior derivative* on the light–cone [19, 48, 67] which has been extensively used for the construction of local conformal operators [46, 49]. In four dimensions it is given by

$$\mathrm{d}_\alpha \equiv \big(1 + \tilde{x}\tilde{\partial}\big)\tilde{\partial}_\alpha - \tfrac{1}{2}\,\tilde{x}_\alpha\tilde{\partial}^2 \quad \text{with} \quad \tilde{\partial}_\alpha \equiv \frac{\partial}{\partial\tilde{x}^\alpha}\,,$$

and has the following properties (see Section 4.1)

$$\mathrm{d}^2 = 0\,, \quad [\mathrm{d}_\alpha, \mathrm{d}_\beta] = 0 \quad \text{and} \quad \mathrm{d}_\alpha \tilde{x}^2 = \tilde{x}^2\big(\mathrm{d}_\alpha + 2\tilde{\partial}_\alpha\big)\,.$$

Then, the conditions of tracelessness simplify, namely, they read:

$$g^{\alpha\beta} G^{\mathrm{tw2}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = 0\,, \qquad \mathrm{d}^\alpha G^{\mathrm{tw2}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = 0\,,$$

as well as

$$\mathrm{d}^\alpha G^{\mathrm{tw2}}_{(\alpha\bullet)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = 0\,.$$

Analogous conditions hold for the light–cone operators of definite twist in the case of symmetry classes (ii) – (iv) obtained below.

We remark that the interior derivative may be used in defining more directly totally symmetric local light–cone operators. For example it holds (see (4.1.12) and (4.1.17))

$$\begin{aligned} G^{\mathrm{tw2(i)}}_{(\alpha\beta)n}(\tilde{x}) &= \frac{1}{(n{+}2)^2(n{+}1)^2}\,\mathrm{d}_\alpha \mathrm{d}_\beta G_{n+2}(\tilde{x})\,, \\ G^{\mathrm{tw2(i)}}_{(\alpha\bullet)n}(\tilde{x}) &= \frac{1}{(n{+}2)^2}\,\mathrm{d}_\alpha G_{n+2}(\tilde{x})\,. \end{aligned}$$

However, in case of symmetry types (ii) – (iv) the corresponding expressions should be more complicated (see Section 4.1).

4.2.2 Tensor operators of symmetry class (ii)

Construction of nonlocal (anti)symmetric class-(ii) operators of definite twist: Young tableau A

Now we consider tensor operators, and their contractions with x, having symmetry class (ii) and whose local twist–3 parts are contained in $\mathbf{T}(\frac{n+2}{2}, \frac{n}{2}) \oplus \mathbf{T}(\frac{n}{2}, \frac{n+2}{2})$. Contrary to the totally symmetric case we have different possibilities to put the tensor indices into the corresponding Young pattern. Without presupposing any symmetry of indices α and β we should start with the following Young tableau:

$$\begin{array}{|c|c|c|c|c|}\hline \mu_1 & \mu_2 & \quad\cdots\quad & \mu_n & \beta \\ \hline \alpha \\ \cline{1-1} \end{array} \;\hat{=}\; \frac{2(n{+}1)}{n{+}2} \mathop{\mathcal{A}}_{\alpha\mu_1} \mathop{\mathcal{S}}_{\beta\mu_1\ldots\mu_n} F_\alpha{}^\rho(0) D_{\mu_1} \ldots D_{\mu_n} F_{\beta\rho}(0) - \text{trace terms},$$

with normalizing factor α_{n+1}. (The tableau with $\alpha \leftrightarrow \beta$ will be considered thereafter.) Denoting this symmetry behaviour by (iiA) we may write the local twist-3 tensor operator as follows:

$$\begin{aligned} G^{\mathrm{tw3(iiA)}}_{\alpha\beta\mu_1\ldots\mu_n} = {} & \frac{1}{n{+}2}\Big\{ F_\alpha{}^\rho(0) D_{(\mu_1} \ldots D_{\mu_n)} F_{\beta\rho}(0) + F_\alpha{}^\rho(0) D_\beta D_{(\mu_2} \ldots D_{\mu_n} F_{\mu_1)\rho}(0) \\ & + \sum_{l=2}^{n} F_\alpha{}^\rho(0) D_{(\mu_1} \ldots D_{\mu_{l-1}} D_{|\beta|} D_{\mu_{l+1}} \ldots D_{\mu_n} F_{\mu_l)\rho}(0) - (\alpha \leftrightarrow \mu_1) \Big\} - \text{trace terms.} \end{aligned}$$

Proceeding in the same manner as in the last Subsection we multiply by $x^{\mu_1} \ldots x^{\mu_n}$ and obtain:

$$\begin{aligned} G^{\mathrm{tw3(iiA)}}_{\alpha\beta n}(x) &= \frac{2}{(n{+}2)n}\, x^\nu \delta^\mu_{[\alpha} \partial_{\nu]} \partial_\beta \overset{\circ}{G}_{\mu\bullet|n+1}(x) \qquad (4.2.28) \\ &= \frac{1}{(n{+}2)n}\Big(\delta^\mu_\alpha (x\partial) - x^\mu \partial_\alpha\Big) \partial_\beta \overset{\circ}{G}_{\mu\bullet|n+1}(x) \end{aligned}$$

with

$$\overset{\circ}{G}_{\mu\bullet|n+1}(x) \equiv x^\nu \overset{\circ}{G}_{\mu\nu|n}(x) = \Big\{\delta_\mu^\alpha - \tfrac{1}{(n+2)^2}\Big[x_\mu\partial^\alpha(x\partial) - \tfrac{1}{2}x^2\partial_\mu\partial^\alpha\Big]\Big\} H^{(4)}_{n+1}(x^2|\Box) G_{\alpha\bullet|n+1}(x)\,.$$

Here, $\overset{\circ}{G}_{\mu\bullet|n+1}(x)$ is the harmonic vector polynomial of order $n+1$ (see, Eq. (4.1.25), Section 4.1) and the symmetry behaviour of class (ii) is obtained through the differential operator in front of it. Now, using $((n+2)n)^{-1} = \int_0^1 \mathrm{d}\lambda(1-\lambda^2)\lambda^n/(2\lambda)$ and Euler's beta function we sum up to obtain the following nonlocal twist–3 tensor operator:

$$G^{\mathrm{tw3(iiA)}}_{\alpha\beta}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\, \tfrac{1-\lambda^2}{2\lambda}\Big(\delta^\mu_\alpha(x\partial) - x^\mu\partial_\alpha\Big)\partial_\beta \overset{\circ}{G}_{\mu\bullet}(0,\kappa\lambda x)\,, \tag{4.2.29}$$

with the nonlocal traceless (vector) operator

$$\begin{aligned}\overset{\circ}{G}_{\alpha\bullet}(0,\kappa x) &= G_{\alpha\bullet}(0,\kappa x) + \sum_{k=1}^\infty \int_0^1 \mathrm{d}t \left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{k!(k-1)!}\left(\frac{1-t}{t}\right)^{k-1} G_{\alpha\bullet}(0,\kappa t x) \\ &\quad - \big[x_\alpha\partial^\mu(x\partial) - \tfrac{1}{2}x^2\partial_\alpha\partial^\mu\big]\sum_{k=0}^\infty \int_0^1 \mathrm{d}\tau\tau \int_0^1 \mathrm{d}t\, t \left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{k!k!}\left(\frac{1-t}{t}\right)^k G_{\mu\bullet}(0,\kappa\tau t x).\end{aligned} \tag{4.2.30}$$

The following decomposition into symmetric and antisymmetric part is useful for the further calculations

$$G^{\mathrm{tw3(iiA)}}_{\alpha\beta}(0,\kappa x) = G^{\mathrm{tw3(iiA)}}_{[\alpha\beta]}(0,\kappa x) + G^{\mathrm{tw3(iiA)}}_{(\alpha\beta)}(0,\kappa x)\,, \tag{4.2.31}$$

with

$$G^{\mathrm{tw3(iiA)}}_{[\alpha\beta]}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\, \lambda \delta^\mu_{[\alpha}\partial_{\beta]} \overset{\circ}{G}_{\mu\bullet}(0,\kappa\lambda x)\,, \tag{4.2.32}$$

$$\begin{aligned}G^{\mathrm{tw3(iiA)}}_{(\alpha\beta)}(0,\kappa x) &= \int_0^1 \mathrm{d}\lambda\, \tfrac{1}{\lambda}\Big(\delta^\mu_{(\alpha}\partial_{\beta)} - \tfrac{1-\lambda^2}{2}\partial_\alpha\partial_\beta x^\mu\Big)\overset{\circ}{G}_{\mu\bullet}(0,\kappa\lambda x) \\ &= \int_0^1 \mathrm{d}\lambda\, \tfrac{1-\lambda^2}{2\lambda}\Big(\delta^\mu_{(\alpha}(x\partial) - x^\mu\partial_{(\alpha}\Big)\partial_{\beta)}\overset{\circ}{G}_{\mu\bullet}(0,\kappa\lambda x)\,.\end{aligned} \tag{4.2.33}$$

Again, these operators are harmonic tensor functions. The conditions of tracelessness for the tensor operator are

$$\begin{aligned} g^{\alpha\beta}G^{\mathrm{tw3(iiA)}}_{\alpha\beta}(0,\kappa x) &= 0\,, \qquad \Box G^{\mathrm{tw3(iiA)}}_{\alpha\beta}(0,\kappa x) = 0\,, \\ \partial^\alpha G^{\mathrm{tw3(iiA)}}_{\alpha\beta}(0,\kappa x) &= 0\,, \qquad \partial^\beta G^{\mathrm{tw3(iiA)}}_{\alpha\beta}(0,\kappa x) = 0\,. \end{aligned} \tag{4.2.34}$$

Reduction to vector operators

The corresponding twist–3 vector operator is obtained from Eq. (4.2.29) by multiplication by x^β:

$$G^{\mathrm{tw3(iiA)}}_{\alpha\bullet}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\, \lambda\Big(\delta^\mu_\alpha(x\partial) - x^\mu\partial_\alpha\Big)\overset{\circ}{G}_{\mu\bullet}(0,\kappa\lambda x). \tag{4.2.35}$$

Obviously, a corresponding scalar operator does not exist.

This vector operator fulfils the following conditions of tracelessness

$$\Box G^{\mathrm{tw3(iiA)}}_{\alpha\bullet}(0,\kappa x) = 0\,, \qquad \partial^\alpha G^{\mathrm{tw3(iiA)}}_{\alpha\bullet}(0,\kappa x) = 0\,. \tag{4.2.36}$$

Projection onto the light–cone

(a) The calculation of the *antisymmetric* tensor operator $G^{\rm tw3(iiA)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ on the light–cone is similar to that of $M^{\rm tw2}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ in [96, 63]; for the details we refer to it. The resulting expression is:

$$
\begin{aligned}
G^{\rm tw3(iiA)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \int_0^1 {\rm d}\lambda\, \lambda \delta^{\mu}_{[\alpha}\partial_{\beta]} G_{\mu\bullet}(\kappa_1\lambda\tilde{x},\kappa_2\lambda\tilde{x})\big|_{x=\tilde{x}} - G^{>{\rm (iiA)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}), && (4.2.37)\\
G^{>{\rm (iiA)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}\int_0^1 {\rm d}\lambda\,(1-\lambda)\Big\{\big(2x_{[\alpha}\partial_{\beta]}\partial^{\mu} - x_{[\alpha}\delta^{\mu}_{\beta]}\Box\big) G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\
&\quad -(1-\lambda+\lambda\ln\lambda)x_{[\alpha}\partial_{\beta]}\Box G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\Big\}\,. && (4.2.38)
\end{aligned}
$$

$G^{>{\rm (iiA)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ contains twist–4 and twist–5 contributions, but twist–6 contributions do not appear due to $x_{[\alpha}x_{\beta]} = 0$. The higher twist operator (4.2.38) contains two vector operators multiplied by x_α and x_β, respectively. For their twist decomposition one has to take into account Young pattern (i) as well as (ii). The procedure is analogous to the decomposition of the vector operator $O_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})$ made in [96, 63]. After a straightforward calculation we obtain

$$
G^{>{\rm (iiA)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw4(iiA)a}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4(iiA)b}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw5(iiA)a}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \qquad (4.2.39)
$$

with

$$
\begin{aligned}
G^{\rm tw4(iiA)a}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}\, x_{[\alpha}\partial_{\beta]}\partial^{\mu}\int_0^1 {\rm d}\lambda\, \frac{1-\lambda^2}{\lambda} G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\rm tw4(iiA)b}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\frac{1}{2}\, x_{[\alpha}\partial_{\beta]}\Box\int_0^1 {\rm d}\lambda \Big(\frac{1-\lambda^2}{2\lambda} + \lambda\ln\lambda\Big) G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\rm tw5(iiA)a}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -x_{[\alpha}\big(\delta^{\mu}_{\beta]}(x\partial) - x^{\mu}\partial_{\beta]}\big)\Box\int_0^1 {\rm d}\lambda\, \frac{(1-\lambda)^2}{4\lambda} G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,.
\end{aligned}
$$

(b) Now, we determine the *symmetric* tensor operator $G^{\rm tw3(iiA)}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ on the light–cone. Putting Eq. (4.2.30) into (4.2.33), after some lengthy but straightforward calculation (taking into account only the relevant terms of the k-summation and performing some partial integrations), we get the following result:

$$
G^{\rm tw3(iiA)}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 {\rm d}\lambda\, \frac{1-\lambda^2}{2\lambda}\Big(\delta^{\mu}_{(\alpha}(x\partial) - x^{\mu}\partial_{(\alpha}\Big)\partial_{\beta)} G_{\mu\bullet}(\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{>{\rm (iiA)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \qquad (4.2.40)
$$

with the higher twist contributions of the trace terms

$$
\begin{aligned}
G^{>{\rm (iiA)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 {\rm d}\lambda \Big\{ &\Big[\frac{1-\lambda^2}{2\lambda} g_{\alpha\beta}\partial^{\mu} + \frac{1-\lambda}{2\lambda}\delta^{\mu}_{(\alpha}x_{\beta)}\Box + (1-\lambda)x_{(\alpha}\partial_{\beta)}\partial^{\mu} \\
&- \frac{(1-\lambda)^2}{4\lambda} x_\alpha x_\beta \partial^{\mu}\Box\Big] G_{\mu\bullet}(\kappa_1\lambda\tilde{x},\kappa_2\lambda\tilde{x}) \\
&- \Big[\frac{1}{2}\Big(\frac{1-\lambda^2}{2\lambda} + \lambda\ln\lambda\Big) g_{\alpha\beta}\Box + \Big(\frac{1}{2}(1-\lambda) + \frac{1-\lambda^2}{4\lambda} + \lambda\ln\lambda\Big) x_{(\alpha}\partial_{\beta)}\Box \\
&+ \frac{1}{4}\Big(\frac{1-\lambda^2}{2\lambda} - \frac{(1-\lambda)^2}{\lambda} + \lambda\ln\lambda\Big) x_\alpha x_\beta \Box^2\Big] G(\kappa_1\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}. \qquad (4.2.41)
\end{aligned}
$$

It is obvious that Eq. (4.2.41) contains scalar and vector operators. Again, using Young patterns (i) and (ii) and subtracting the trace terms, we can decompose the vector part appearing in Eq. (4.2.41) into twist–4, twist–5 and twist–6 operators. This procedure is analogous to the twist decomposition of the (vector) quark operator $O_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})$ in [96, 63]I. From the twist–4 and twist–5 vector part we have to subtract their trace terms being of twist–6 and add it to the other twist–6 scalar operator. Let us recall that the scalar twist–4 and twist–6 operators are already traceless on the cone. In that manner we get the following decomposition:

$$
\begin{aligned}
G^{>\mathrm{(iiA)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =\,& G^{\mathrm{tw4(iiA)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw4(iiA)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw4(iiA)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
&+G^{\mathrm{tw4(iiA)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw4(iiA)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw4(iiA)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
&+G^{\mathrm{tw5(iiA)d}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw6(iiA)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw6(iiA)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
&+G^{\mathrm{tw6(iiA)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw6(iiA)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw6(iiA)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
&+G^{\mathrm{tw6(iiA)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw6(iiA)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\mathrm{tw6(iiA)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}),
\end{aligned}
\tag{4.2.42}
$$

with

$$
\begin{aligned}
G^{\mathrm{tw4(iiA)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\tfrac{1}{2}g_{\alpha\beta}\Box\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda}+\lambda\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw4(iiA)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\tfrac{1}{2}x_{(\alpha}\partial_{\beta)}\Box\int_0^1 \mathrm{d}\lambda\Big((1-\lambda)+\frac{1-\lambda^2}{2\lambda}+2\lambda\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\\
&-G^{\mathrm{tw6(iiA)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,\\
G^{\mathrm{tw4(iiA)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\tfrac{1}{2}x_{(\alpha}\partial_{\beta)}\Box\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\\
&-G^{\mathrm{tw6(iiA)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw6(iiA)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&\,\tfrac{1}{4}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{(1-\lambda)^2}{\lambda}-\frac{1-\lambda^2}{2\lambda}-\lambda\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iiA)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\tfrac{1}{4}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{(1-\lambda)^2}{2\lambda}-\frac{3}{2}\frac{1-\lambda^2}{2\lambda}-\lambda\ln\lambda-\frac{\ln\lambda}{2\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iiA)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&\,\tfrac{1}{4}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}+\frac{\ln^2\lambda}{2\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iiA)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\tfrac{1}{2}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}+\frac{\ln^2\lambda}{2\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the scalar operator $G(\kappa_1\tilde{x},\kappa_2\tilde{x})$, as well as

$$
\begin{aligned}
G^{\mathrm{tw4(iiA)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&\,\tfrac{1}{2}g_{\alpha\beta}\partial^\mu\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda}G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}},\\
G^{\mathrm{tw4(iiA)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&\,x_{(\alpha}\partial_{\beta)}\partial^\mu\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}\Big)G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\\
&-G^{\mathrm{tw6(iiA)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,\\
G^{\mathrm{tw4(iiA)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&\,x_{(\alpha}\partial_{\beta)}\partial^\mu\int_0^1 \mathrm{d}\lambda(1-\lambda)G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}-G^{\mathrm{tw6(iiA)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw5(iiA)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = &-\frac{1}{2}x_{(\alpha}\big(\delta^{\mu}_{\beta)}(x\partial)-x^{\mu}\partial_{\beta)}\big)\Box\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}\Big)G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
&- G^{\mathrm{tw6(iiA)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) - G^{\mathrm{tw6(iiA)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw6(iiA)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\frac{1}{4}x_\alpha x_\beta\Box\partial^\mu\int_0^1 \mathrm{d}\lambda\frac{(1-\lambda)^2}{\lambda}G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iiA)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\frac{1}{2}x_\alpha x_\beta\Box\partial^\mu\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}+\frac{\ln^2\lambda}{2\lambda}\Big)G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iiA)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}x_\alpha x_\beta\Box\partial^\mu\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}+\frac{\ln\lambda}{\lambda}+\frac{\ln^2\lambda}{\lambda}\Big)G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iiA)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{4}x_\alpha x_\beta\Box\partial^\mu\int_0^1 \mathrm{d}\lambda\,\frac{(1-\lambda)^2}{\lambda}G_{\mu\bullet}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the vector operator $G_{\mu\bullet}(\kappa_1\tilde{x},\kappa_2\tilde{x})$.

Contributions of Young tableau B

Let us now consider the other Young tableau which is obtained from the former one by exchanging α and β:

$$
\begin{array}{|c|c|c|c|c|}\hline \mu_1 & \mu_2 & \quad\cdots\quad & \mu_n & \alpha \\ \hline \beta \\ \cline{1-1}\end{array}
\;\hat{=}\; \frac{2(n+1)}{n+2}\mathop{\mathcal{A}}_{\beta\mu_1}\mathop{\mathcal{S}}_{\alpha\mu_1\ldots\mu_n} F_{\alpha}{}^{\rho}(0)D_{\mu_1}\ldots D_{\mu_n}F_{\beta\rho}(0) - \text{trace terms.}
$$

This symmetry behaviour will be denoted by (iiB). The corresponding nonlocal twist–3 operator which we obtain after analogous calculations is (with self-explaining denotations):

$$
G^{\mathrm{tw3(iiB)}}_{\alpha\beta}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\,\frac{1-\lambda^2}{2\lambda}\Big(\delta^{\mu}_{\beta}(x\partial)-x^{\mu}\partial_{\beta}\Big)\partial_\alpha\overset{\circ}{G}_{\bullet\mu}(0,\kappa\lambda x)\,. \tag{4.2.43}
$$

Its decomposition into symmetric and antisymmetric part yields

$$
G^{\mathrm{tw3(iiB)}}_{\alpha\beta}(0,\kappa x) = G^{\mathrm{tw3(iiB)}}_{[\alpha\beta]}(0,\kappa x) + G^{\mathrm{tw3(iiB)}}_{(\alpha\beta)}(0,\kappa x)\,, \tag{4.2.44}
$$

with

$$
G^{\mathrm{tw3(iiB)}}_{[\alpha\beta]}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\,\lambda\delta^{\mu}_{[\beta}\partial_{\alpha]}\overset{\circ}{G}_{\bullet\mu}(0,\kappa\lambda x)\,, \tag{4.2.45}
$$

$$
G^{\mathrm{tw3(iiB)}}_{(\alpha\beta)}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\,\frac{1-\lambda^2}{2\lambda}\Big(\delta^{\mu}_{(\alpha}(x\partial)-x^{\mu}\partial_{(\alpha}\Big)\partial_{\beta)}\overset{\circ}{G}_{\bullet\mu}(0,\kappa\lambda x)\,. \tag{4.2.46}
$$

The projection onto the light–cone and the calculation of the higher twist operators contained in the trace terms is completely analogous to case (A) and should be omitted here.

Construction of the complete twist–3 tensor operators on the light–cone

In order to obtain the complete twist–3 operator it is necessary to add both twist–3 operators (4.2.31) and (4.2.44) resulting from the Young patterns (iiA) and (iiB). Since no further Young pattern contributes to twist–3 operators we omit the index (ii). After projection onto the light–cone the final result is

$$G^{\rm tw3}_{\alpha\beta}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw3}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw3}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \tag{4.2.47}$$

with the *antisymmetric twist–3 light–cone tensor operator*[4]

$$G^{\rm tw3}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =2\int_0^1 {\rm d}\lambda\,\lambda\partial_{[\beta}G^{-}_{\alpha]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{\rm tw4(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) - G^{\rm tw5(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}), \tag{4.2.48}$$

where

$$G^{\rm tw4(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = x_{[\alpha}\partial_{\beta]}\partial^{\mu}\int_0^1 {\rm d}\lambda\,\frac{1-\lambda^2}{\lambda}\;G^{-}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \tag{4.2.49}$$

$$G^{\rm tw5(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = -x_{[\alpha}\big(\delta^{\mu}_{\beta]}(x\partial) - x^{\mu}\partial_{\beta]}\big)\Box\int_0^1 {\rm d}\lambda\;\frac{(1-\lambda)^2}{2\lambda}G^{-}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \tag{4.2.50}$$

The local twist–4 operators are contained in the tensor space $\mathbf{T}(\frac{n}{2},\frac{n}{2})$ and the local twist–5 operators are contained in the tensor space $\mathbf{T}(\frac{n}{2},\frac{n-2}{2})\oplus\mathbf{T}(\frac{n-2}{2},\frac{n}{2})$.

In turn, let us remark that the bilocal light–ray operator $G^{\rm tw3}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ is the same as the antisymmetric tensor operator which obtains from the Young tableau $\begin{array}{|c|c|c|c|c|}\hline \beta & \mu_1 & \mu_2 & \cdots & \mu_n \\ \hline \alpha \\ \cline{1-1}\end{array}$. That tableau has been used for the computation of $M^{\rm tw2}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ in Ref. [96]. Furthermore, let us note that the local operators contained in $M^{\rm tw2}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ and in $G^{\rm tw3}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ are in complete agreement with the local operators on the light–cone determined by Dobrev and Ganchev [49] by means of the interior derivative.

From the above point of view our result for the antisymmetric part of twist–3 operator (4.2.47) could be obtained much easier. However, the *symmetric twist–3 light–cone tensor operator* is much more involved. It is given by[5]

$$\begin{aligned} G^{\rm tw3}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = &\int_0^1 {\rm d}\lambda\,\frac{1-\lambda^2}{2\lambda}\Big(\delta^{\mu}_{(\alpha}(x\partial) - x^{\mu}\partial_{(\alpha}\Big)\partial_{\beta)}G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ &- G^{>{\rm (ii)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \end{aligned} \tag{4.2.51}$$

where $G^{>{\rm (ii)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ includes all the trace terms having higher twist; these operators are given by

$$\begin{aligned} G^{\rm tw4(ii)a}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&- g_{\alpha\beta}\Box\int_0^1 {\rm d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \lambda\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\ G^{\rm tw4(ii)b}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&- x_{(\alpha}\partial_{\beta)}\Box\int_0^1 {\rm d}\lambda\Big((1-\lambda)+\frac{1-\lambda^2}{2\lambda}+2\lambda\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ &- G^{\rm tw6(ii)b}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \\ G^{\rm tw4(ii)d1}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =&- x_{(\alpha}\partial_{\beta)}\Box\int_0^1 {\rm d}\lambda\Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ &- G^{\rm tw6(ii)d1}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \end{aligned}$$

---

[4] Here, we used the abbreviation $G^{-}_{\mu}\equiv x^{\nu}G_{[\mu\nu]}$ instead of $G_{[\mu\bullet]}$ in order not to come into conflict with the antisymmetrization of the indices α and β.

[5] Similarly, we use the notation G^{+}_{μ} instead of $G_{(\mu\bullet)}$.

and

$$
\begin{aligned}
G^{\mathrm{tw6(ii)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& \frac{1}{2}x_\alpha x_\beta \Box^2 \int_0^1 \mathrm{d}\lambda \Big(\frac{(1-\lambda)^2}{\lambda} - \frac{1-\lambda^2}{2\lambda} - \lambda\ln\lambda\Big) G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(ii)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\frac{1}{2}x_\alpha x_\beta \Box^2 \int_0^1 \mathrm{d}\lambda \Big(\frac{(1-\lambda)^2}{2\lambda} - \frac{3}{2}\frac{1-\lambda^2}{2\lambda}\\
&\qquad\qquad - \lambda\ln\lambda - \frac{\ln\lambda}{2\lambda}\Big) G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(ii)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& \frac{1}{2}x_\alpha x_\beta \Box^2 \int_0^1 \mathrm{d}\lambda \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda} + \frac{\ln^2\lambda}{2\lambda}\Big) G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(ii)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -x_\alpha x_\beta \Box^2 \int_0^1 \mathrm{d}\lambda \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda} + \frac{\ln^2\lambda}{2\lambda}\Big) G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the scalar operator $G(\kappa_1 x,\kappa_2 x)$, as well as

$$
\begin{aligned}
G^{\mathrm{tw4(ii)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& g_{\alpha\beta}\partial^\mu \int_0^1 \mathrm{d}\lambda \frac{1-\lambda^2}{\lambda} G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw4(ii)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& 2x_{(\alpha}\partial_{\beta)}\partial^\mu \int_0^1 \mathrm{d}\lambda \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\\
&- G^{\mathrm{tw6(ii)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,\\
G^{\mathrm{tw4(ii)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& 2x_{(\alpha}\partial_{\beta)}\partial^\mu \int_0^1 \mathrm{d}\lambda (1-\lambda) G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\\
&- G^{\mathrm{tw6(ii)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw5(ii)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -x_{(\alpha}\big(\delta^\mu_{\beta)}(x\partial) - x^\mu\partial_{\beta)}\big)\Box\\
&\times \int_0^1 \mathrm{d}\lambda \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
&- G^{\mathrm{tw6(ii)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) - G^{\mathrm{tw6(ii)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw6(ii)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -\frac{1}{2}x_\alpha x_\beta \Box\partial^\mu \int_0^1 \mathrm{d}\lambda\, \frac{(1-\lambda)^2}{\lambda} G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(ii)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& -x_\alpha x_\beta \Box\partial^\mu \int_0^1 \mathrm{d}\lambda \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda} + \frac{\ln^2\lambda}{2\lambda}\Big) G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(ii)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& x_\alpha x_\beta \Box\partial^\mu \int_0^1 \mathrm{d}\lambda \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda} + \frac{\ln^2\lambda}{\lambda}\Big) G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(ii)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& \frac{1}{2}x_\alpha x_\beta \Box\partial^\mu \int_0^1 \mathrm{d}\lambda \frac{(1-\lambda)^2}{\lambda} G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the (symmetric) vector operator $G^+_\mu(\kappa_1 x,\kappa_2 x)$.

Determination of complete light–cone vector operators

Now we are able to determine the twist–3 vector operator resulting from the tensor operator (4.2.47) by multiplying by x^β. Because neither symmetry type (iii) nor (iv) may contribute to the vector operator this will be the final expression. The *twist–3 light–cone vector operator* consists of two parts, one originating from the symmetric and the other from the antisymmetric tensor operator,

$$G^{\rm tw3}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw3}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw3}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \tag{4.2.52}$$

where

$$G^{\rm tw3}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 {\rm d}\lambda\lambda\Big[\delta^{\mu}_{\alpha}(x\partial) - x^{\mu}\partial_{\alpha} - x_{\alpha}\big(\partial^{\mu} + \ln\lambda\,\square x^{\mu}\big)\Big] G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \tag{4.2.53}$$

$$G^{\rm tw3}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = \int_0^1 {\rm d}\lambda\,\lambda\Big[\delta^{\mu}_{\alpha}(x\partial) - x^{\mu}\partial_{\alpha} - x_{\alpha}\partial^{\mu}\Big] G^{-}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \tag{4.2.54}$$

and the twist–4 vector operator which is contained in the trace terms of that twist–3 vector operator is given by

$$G^{\rm tw4(ii)}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw4(ii)}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \tag{4.2.55}$$

where

$$G^{\rm tw4(ii)}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = x_{\alpha}\int_0^1 {\rm d}\lambda\,\lambda\big[\partial^{\mu} + (\ln\lambda)\square x^{\mu}\big] G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \tag{4.2.56}$$

$$G^{\rm tw4}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = x_{\alpha}\int_0^1 {\rm d}\lambda\,\lambda\,\partial^{\mu} G^{-}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \tag{4.2.57}$$

The antisymmetric part of the twist–4 vector operator is already complete. The complete *twist–4 light–cone vector operator* is given by

$$\begin{aligned} G^{\rm tw4}_{\alpha}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= G^{\rm tw4}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\ &= x_{\alpha}\int_0^1 {\rm d}\lambda\,\lambda\big[\partial^{\mu} + \frac{\ln\lambda}{2}\square x^{\mu}\big] G_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \end{aligned} \tag{4.2.58}$$

where the symmetric twist–4 vector operator obtains by adding together expressions (4.2.26) and (4.2.55),

$$\begin{aligned} G^{\rm tw4}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= G^{\rm tw4(i)}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4(ii)}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\ &= x_{\alpha}\int_0^1 {\rm d}\lambda\,\lambda\big[\partial^{\mu} + \frac{\ln\lambda}{2}\square x^{\mu}\big] G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \end{aligned} \tag{4.2.59}$$

Furthermore, putting together expressions (4.2.25), (4.2.53), (4.2.59) and (4.2.54), (4.2.57) we obtain for the complete decomposition of the *light–cone vector operator* the following results anticipated by Eqs. (4.2.9) and (4.2.10), respectively:

$$G_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw2}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw3}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4}_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \tag{4.2.60}$$

$$G_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw3}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4(ii)}_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \tag{4.2.61}$$

4.2.3 Tensor operators of symmetry class (iii)

Construction of nonlocal antisymmetric class-(iii) operators of definite twist

Now we consider the twist–4 and twist–5 contributions originating from the (unique) Young tableau related to the symmetry class (iii) (with the normalizing factor β_n):

$$\begin{array}{|c|c|c|c|c|}\hline \mu_1 & \mu_2 & \mu_3 & \quad\cdots\quad & \mu_n \\ \hline \beta \\ \cline{1-1} \alpha \\ \cline{1-1} \end{array} \;\hat{=}\; \frac{3n}{n+2} \mathop{\mathcal{A}}_{\alpha\beta\mu_1} \mathop{\mathcal{S}}_{\mu_1\ldots\mu_n} F_\alpha{}^\rho(0) D_{\mu_1} \ldots D_{\mu_n} F_{\beta\rho}(0) - \text{trace terms}\,.$$

The corresponding traceless local operator having twist $\tau = 4$ and being contained in the tensor space $\mathbf{T}\big(\frac{n}{2}, \frac{n}{2}\big)$ is given by:

$$\begin{aligned} G^{\mathrm{tw4(iii)}}_{\alpha\beta\mu_1\ldots\mu_n} \;=\; \frac{n}{n+2}\Big(& F_{[\alpha|}{}^\rho(0) D_{(\mu_1} \ldots D_{\mu_n)} F_{|\beta]\rho}(0) - F_{[\alpha|}{}^\rho(0) D_{(\beta} D_{\mu_2} \ldots D_{\mu_n)} F_{|\mu_1]\rho}(0) \\ & - F_{[\mu_1|}{}^\rho(0) D_{(\alpha} D_{\mu_2} \ldots D_{\mu_n)} F_{|\beta]\rho}(0)\Big) - \text{trace terms.} \end{aligned}$$

Now, contracting this expression with $x^{\mu_1} \ldots x^{\mu_n}$ we obtain:

$$\begin{aligned} G^{\mathrm{tw4(iii)}}_{[\alpha\beta]n}(x) &= x^{\mu_1} \ldots x^{\mu_n} G^{\mathrm{tw4(iii)}}_{[\alpha\beta]\mu_1\ldots\mu_n} \\ &= \frac{3}{n+2}\, x^\gamma\, \delta^\mu_{[\alpha} \delta^\nu_\beta \partial_{\gamma]} \overset{\circ}{G}_{[\mu\nu]n}(x) \qquad (4.2.62) \\ &= \frac{1}{n+2} \Big((x\partial)\delta^\nu_{[\beta} - 2x^\nu \partial_{[\beta}\Big)\, \delta^\mu_{\alpha]} \overset{\circ}{G}_{[\mu\nu]n}(x)\,. \end{aligned}$$

with the (antisymmetric) tensorial harmonic polynomials of order n (see (4.1.40)):

$$\begin{aligned} \overset{\circ}{G}_{[\alpha\beta]n}(x) =\Big\{ & \delta^\mu_\alpha \delta^\nu_\beta + \frac{1}{(n+1)n} \Big(2x_{[\alpha} \delta^{[\mu}_{\beta]} \partial^{\nu]} (x\partial) - x^2 \partial_{[\alpha} \delta^{[\mu}_{\beta]} \partial^{\nu]}\Big) \\ & - \frac{2}{(n+2)(n+1)n}\, x_{[\alpha} \partial_{\beta]} x^{[\mu} \partial^{\nu]} \Big\} H^{(4)}_n\big(x^2|\Box\big) G_{[\mu\nu]n}(x). \end{aligned}$$

Resumming these local terms gives the nonlocal twist–4 operator as follows

$$G^{\mathrm{tw4(iii)}}_{[\alpha\beta]}(0,\kappa x) = \int_0^1 \mathrm{d}\lambda\, \lambda \delta^\mu_{[\alpha} \Big((x\partial)\delta^\nu_{\beta]} - 2x^\nu \partial_{\beta]}\Big)\, \overset{\circ}{G}_{[\mu\nu]}(0,\kappa\lambda x)\,, \qquad (4.2.63)$$

where, using the integral representation of the additional factor $1/n$ (the remaining factors of the denominator are taken together with $1/n!$ to get $1/(n+2)!$) and of the beta function, we introduced:

$$\begin{aligned} \overset{\circ}{G}_{[\alpha\beta]}(0,\kappa x) = G_{[\alpha\beta]}(0,\kappa x) &+ \sum_{k=1}^\infty \int_0^1 \frac{\mathrm{d}t}{t} \left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{k!(k-1)!} \left(\frac{1-t}{t}\right)^{k-1} G_{[\alpha\beta]}(0,\kappa t x) \\ + \int_0^1 \frac{\mathrm{d}\tau}{\tau} \Big\{ & \Big(2x_{[\alpha} \delta^{[\mu}_{\beta]} \partial^{\nu]} (x\partial) - x^2 \partial_{[\alpha} \delta^{[\mu}_{\beta]} \partial^{\nu]}\Big) \sum_{k=0}^\infty \int_0^1 \mathrm{d}t \left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{k!k!} \left(\frac{1-t}{t}\right)^k \\ & - 2x_{[\alpha} \partial_{\beta]} x^{[\mu} \partial^{\nu]} \sum_{k=0}^\infty \int_0^1 \mathrm{d}t\, t \left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{(k+1)!k!} \left(\frac{1-t}{t}\right)^{k+1} \Big\} G_{[\mu\nu]}(0,\kappa\tau t x). \qquad (4.2.64) \end{aligned}$$

Then, by construction, Eq. (4.2.63) defines a nonlocal operator of twist–4. As is easily seen by partial integration it fulfils the following relation

$$G^{\mathrm{tw4(iii)}}_{[\alpha\beta]}(0,\kappa x) = \overset{\circ}{G}_{[\alpha\beta]}(0,\kappa x) - G^{\mathrm{tw3}}_{[\alpha\beta]}(0,\kappa x) \qquad (4.2.65)$$

and, because of Eq. (4.2.34) and the properties of (4.2.64), it is a harmonic tensor operator:

$$\Box G^{\rm tw4(iii)}_{[\alpha\beta]}(0,\kappa x) = 0\ , \qquad \partial^\alpha G^{\rm tw4(iii)}_{[\alpha\beta]}(0,\kappa x) = 0\,. \tag{4.2.66}$$

Projection onto the light–cone

Let us now project onto the light–cone. After the same calculations as has been carried out for the operator $M^{\rm tw3}_{[\alpha\beta]}(0,\kappa\tilde{x})$ in [63] we obtain

$$\begin{aligned} G^{\rm tw4(iii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = &\int_0^1 {\rm d}\lambda\,\lambda\delta^\mu_{[\alpha}\left((x\partial)\delta^\nu_{\beta]} - 2x^\nu\partial_{\beta]}\right) G_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ &- G^{\rm tw5(iii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}), \end{aligned} \tag{4.2.67}$$

where the twist–5 part is determined by the trace terms, namely

$$\begin{aligned} G^{\rm tw5(iii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = -\int_0^1 {\rm d}\lambda\frac{1-\lambda^2}{\lambda}\Big\{&x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]} \\ &- x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box\Big\}G_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \end{aligned} \tag{4.2.68}$$

Obviously, there exist no vector (and scalar) operators of symmetry type (iii).

Determination of the complete antisymmetric light–cone tensor operator

This finishes the computation of twist contributions of the antisymmetric light–cone tensor operator. The complete *antisymmetric twist–4 light–cone tensor operator* obtains from Eq. (4.2.67) and the twist–4 trace terms of the twist–3 operator, Eq. (4.2.49):

$$\begin{aligned} G^{\rm tw4}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= G^{\rm tw4(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4(iii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\ &= \int_0^1 {\rm d}\lambda\Big\{\lambda\left((x\partial)\delta^\nu_{[\beta} - 2x^\nu\partial_{[\beta}\right)\delta^\mu_{\alpha]} + \frac{1-\lambda^2}{\lambda}\Big(x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]} \\ &\quad -x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box - x_{[\alpha}\partial_{\beta]}x^{[\mu}\partial^{\nu]}\Big)\Big\}G_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \end{aligned} \tag{4.2.69}$$

Furthermore, the complete *antisymmetric twist–5 light–cone tensor operator* obtains from Eqs. (4.2.50) and (4.2.68) as follows:

$$\begin{aligned} G^{\rm tw5}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= G^{\rm tw5(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw5(iii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\ &= \int_0^1 {\rm d}\lambda\frac{1-\lambda}{\lambda}\Big\{x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box - 2x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]}\Big\}G_{[\mu\nu]}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \end{aligned} \tag{4.2.70}$$

Together with the twist-3 part, Eq. (4.2.48), we finally obtain the complete decomposition of the *antisymmetric light–cone tensor operator* (compare Eq. (4.2.8)):

$$G_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = G^{\rm tw3}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw4}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\rm tw5}_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,. \tag{4.2.71}$$

4.2.4 Tensor operators of symmetry class (iv)

Construction of the nonlocal symmetric class-(iv) operator of definite twist

Now let us investigate the symmetric tensor operators of the symmetry class (iv) which are given by the following Young tableau (with normalizing factor γ_n):

$$\begin{array}{|c|c|c|c|c|}\hline \mu_1 & \mu_2 & \mu_3 & \quad\cdots\quad & \mu_n \\ \hline \alpha & \beta \\ \cline{1-2}\end{array} \;\hat{=}\; \frac{4(n-1)}{n+1} \underset{\alpha\mu_1}{\mathcal{A}} \underset{\beta\mu_2}{\mathcal{A}} \underset{\alpha\beta}{\mathcal{S}} \underset{\mu_1\ldots\mu_n}{\mathcal{S}} F_\alpha{}^\rho(0) D_{\mu_1}\ldots D_{\mu_n} F_{\beta\rho}(0) - \text{traces}\,.$$

These local tensor operators have twist–4 and transform according to the representation $\mathbf{T}(\frac{n+2}{2},\frac{n-2}{2}) \oplus \mathbf{T}(\frac{n-2}{2},\frac{n+2}{2})$. They are given by:

$$\begin{aligned} G^{\mathrm{tw4(iv)}}_{\alpha\beta\mu_1\ldots\mu_n} = \frac{1}{(n+1)n}\Big(& G_{(\alpha\beta)(\mu_1\ldots\mu_n)} - G_{(\alpha\mu_2)(\mu_1\beta\mu_3\ldots\mu_n)} - G_{(\mu_1\beta)(\alpha\mu_2\ldots\mu_n)} \\ & + G_{(\mu_1\mu_2)(\alpha\beta\mu_3\ldots\mu_n)}\Big) - \text{trace terms}\,, \end{aligned}$$

with $G_{(\alpha\beta)(\mu_1\ldots\mu_n)} \equiv F_{(\alpha|}{}^{\rho}(0) D_{(\mu_1}\ldots D_{\mu_n)} F_{|\beta)\rho}(0)$. If we multiply this expression by $x^{\mu_1}\ldots x^{\mu_n}$, we obtain:

$$\begin{aligned} G^{\mathrm{tw4(iv)}}_{(\alpha\beta)n}(x) &= x^{\mu_1}\ldots x^{\mu_n} G^{\mathrm{tw4(iv)}}_{\alpha\beta\mu_1\ldots\mu_n} \\ &= \frac{4}{(n+1)n}\, x^\rho x^\tau\, \delta^\mu_{[\alpha}\partial_{\rho]} \delta^\nu_{[\beta}\partial_{\tau]}\, \overset{\circ}{G}_{(\mu\nu)n}(x) \qquad (4.2.72) \\ &= \frac{1}{(n+1)n}\Big(\delta^\mu_\alpha\delta^\nu_\beta x^\rho (x\partial)\partial_\rho - 2x^\mu (x\partial)\delta^\nu_{(\alpha}\partial_{\beta)} + x^\mu x^\nu \partial_\alpha\partial_\beta\Big) \overset{\circ}{G}_{(\mu\nu)n}(x)\,. \end{aligned}$$

Here, $\overset{\circ}{G}_{(\mu\nu)n}(x)$ is the harmonic symmetric tensor polynomial of order n (cf. Eq. (4.1.49), Section 4.1):

$$\begin{aligned} \overset{\circ}{G}_{(\alpha\beta)n}(x) = \Big\{ & \delta^\mu_\alpha\delta^\nu_\beta + \frac{n}{(n+2)^2} g_{\alpha\beta} x^{(\nu}\partial^{\mu)} - \frac{1}{(n+2)(n+1)}\Big(2n x_{(\alpha}\delta^{(\nu}_{\beta)}\partial^{\mu)} - x^2 \delta^{(\nu}_{(\alpha}\partial_{\beta)}\partial^{\nu)}\Big) \\ & - \frac{1}{(n+2)^2(n+1)}\Big(2n x_{(\alpha}\partial_{\beta)} x^{(\nu}\partial^{\mu)} - x^2 \partial_\alpha\partial_\beta x^{(\nu}\partial^{\mu)}\Big) + \frac{(n+3)n}{(n+2)^2(n+1)^2}\Big(x_\alpha x_\beta - \frac{1}{2}x^2 g_{\alpha\beta}\Big)\partial^\mu\partial^\nu \\ & - \frac{1}{(n+2)^2(n+1)^2}\Big(2x^2 \partial_{(\alpha} x_{\beta)} + \frac{1}{4}x^4 \partial_\alpha\partial_\beta\Big)\partial^\mu\partial^\nu \Big\} \breve{G}_{(\mu\nu)n}(x)\,. \end{aligned}$$

Resumming all local operators of symmetry class (iv) and using the integral representation $((n+1)n)^{-1} = \int_0^1 \mathrm{d}\lambda \lambda^n (1-\lambda)/\lambda$ gives

$$G^{\mathrm{tw4(iv)}}_{(\alpha\beta)}(0,\kappa x) = \Big(\delta^\mu_\alpha\delta^\nu_\beta x^\rho (x\partial)\partial_\rho - 2x^\mu (x\partial)\delta^\nu_{(\beta}\partial_{\alpha)} + x^\mu x^\nu \partial_\alpha\partial_\beta\Big) \int_0^1 \mathrm{d}\lambda\, \frac{1-\lambda}{\lambda}\, \overset{\circ}{G}_{(\mu\nu)}(0,\kappa\lambda x), \qquad (4.2.73)$$

where $\overset{\circ}{G}_{(\mu\nu)}(0,\kappa x)$ is given by

$$
\begin{aligned}
\overset{\circ}{G}_{(\alpha\beta)}(0,\kappa x) &= \breve{G}_{(\alpha\beta)}(0,\kappa x) + \sum_{k=1}^{\infty}\int_0^1 \frac{\mathrm{d}t}{t}\Big(\frac{-x^2}{4}\Big)^k \frac{\Box^k}{k!(k-1)!}\Big(\frac{1-t}{t}\Big)^{k-1}\breve{G}_{(\alpha\beta)}(0,\kappa t x)\\
&-\big[2x_{(\alpha}\delta_{\beta)}^{(\mu}\partial^{\nu)}(x\partial) - x^2\partial_{(\alpha}\delta_{\beta)}^{(\mu}\partial^{\nu)}\big]\sum_{k=0}^{\infty}\int_0^1 \mathrm{d}t\,t\left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{k!(k+1)!}\left(\frac{1-t}{t}\right)^{k+1}\breve{G}_{(\mu\nu)}(0,\kappa t x)\\
&-\Big\{\big[2x_{(\alpha}\partial_{\beta)}x^{(\mu}\partial^{\nu)}(x\partial) - g_{\alpha\beta}x^{(\mu}\partial^{\nu)}(x\partial)\big(x\partial+1\big) - x^2\partial_\alpha\partial_\beta x^{(\mu}\partial^{\nu)}\big]\int_0^1 \mathrm{d}\lambda\,\lambda\\
&-\big[\big(x_\alpha x_\beta - \tfrac{x^2}{2}g_{\alpha\beta}\big)\partial^\mu\partial^\nu(x\partial)\big(x\partial+3\big) - \big(2x^2\partial_{(\alpha}x_{\beta)} + \tfrac{x^4}{4}\partial_\alpha\partial_\beta\big)\partial^\mu\partial^\nu\big]\int_0^1 \mathrm{d}\lambda\,(1-\lambda)\Big\}\\
&\qquad\times\sum_{k=0}^{\infty}\int_0^1 \mathrm{d}t\,t\left(\frac{-x^2}{4}\right)^k \frac{\Box^k}{(k+1)!k!}\left(\frac{1-t}{t}\right)^{k+1}\breve{G}_{(\mu\nu)}(0,\kappa\lambda t x). \qquad (4.2.74)
\end{aligned}
$$

Of course, Eq. (4.2.73) may be rewritten as follows

$$
G^{\mathrm{tw4(iv)}}_{(\alpha\beta)}(0,\kappa x) = \overset{\circ}{G}_{(\alpha\beta)}(0,\kappa x) - 2\int_0^1 \mathrm{d}\lambda\,\tfrac{1}{\lambda}\partial_{(\alpha}\overset{\circ}{G}{}^{+}_{\beta)}(0,\kappa\lambda x) + \int_0^1 \mathrm{d}\lambda\,\tfrac{1-\lambda}{\lambda}\partial_\alpha\partial_\beta\overset{\circ}{G}(0,\kappa\lambda x),
$$

which will be used in the further considerations. Contrary to any of the formerly obtained symmetric twist–4 tensor operators, this operator is the 'true tensor part', i.e., not being proportional to $g_{\alpha\beta}$ or x_α and x_β. Obviously, this twist–4 operator satisfies the conditions of tracelessness

$$
\Box G^{\mathrm{tw4(iv)}}_{(\alpha\beta)}(0,\kappa x) = 0, \quad \partial^\alpha G^{\mathrm{tw4(iv)}}_{(\alpha\beta)}(0,\kappa x) = 0, \quad g^{\alpha\beta}G^{\mathrm{tw4(iv)}}_{(\alpha\beta)}(0,\kappa x). \qquad (4.2.75)
$$

Projection onto the light–cone

The symmetric twist–4 tensor operator of symmetry type (iv) on the light–cone is obtained as follows

$$
\begin{aligned}
G^{\mathrm{tw4(iv)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& \Big(\delta^\mu_\alpha\delta^\nu_\beta x^\rho(x\partial)\partial_\rho - 2x^\mu(x\partial)\delta^\nu_{(\beta}\partial_{\alpha)} + x^\mu x^\nu\partial_\alpha\partial_\beta\Big)\\
&\times\int_0^1 \mathrm{d}\lambda\frac{1-\lambda}{\lambda}G_{(\mu\nu)}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{>(\mathrm{iv})}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}), \qquad (4.2.76)
\end{aligned}
$$

where the trace terms of higher twist are given by

$$
\begin{aligned}
G^{>(\mathrm{iv})}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) =& \int_0^1 \mathrm{d}\lambda\Big\{\Big(2\lambda x_{(\alpha}\delta^\nu_{\beta)}\partial^\mu - (1-\lambda)x_\alpha x_\beta\partial^\mu\partial^\nu\Big)G_{(\mu\nu)}(\kappa_1\lambda x,\kappa_2\lambda x)\\
&-\Big(\tfrac{1}{\lambda}g_{\alpha\beta}\partial^\mu + \tfrac{1-\lambda^2}{\lambda}x_{(\alpha}\delta^\mu_{\beta)}\Box - \tfrac{(1-\lambda)^2}{2\lambda}x_\alpha x_\beta\Box\partial^\mu\Big)G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\\
&+\Big(\tfrac{1-\lambda}{2\lambda}g_{\alpha\beta}\Box + \tfrac{(1-\lambda)^2}{2\lambda}x_{(\alpha}\partial_{\beta)}\Box - \tfrac{1}{2}\Big(\tfrac{1-\lambda^2}{2\lambda} + \ln\lambda\Big)x_\alpha x_\beta\Box^2\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\\
&+\Big(\tfrac{1}{2\lambda}g_{\alpha\beta}(x\partial) - \lambda x_{(\alpha}\partial_{\beta)} + \tfrac{1}{2}(1-\lambda)x_\alpha x_\beta\Box\Big)G^\rho{}_\rho(\kappa_1\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}. \qquad (4.2.77)
\end{aligned}
$$

The explicit twist decomposition of this operator by means of the Young tableaux (i) and (ii) gives

$$\begin{aligned}
G^{>(\mathrm{iv})}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & G^{\mathrm{tw4(iv)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(iv)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(iv)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw4(iv)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(iv)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(iv)e1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw4(iv)e2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(iv)f}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw4(iv)h}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw5(iv)d}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw5(iv)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw6(iv)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw6(iv)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw6(iv)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)e1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)e2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw6(iv)e3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)e4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)e5}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) \\
& + G^{\mathrm{tw6(iv)f}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)h}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) + G^{\mathrm{tw6(iv)k}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned} \tag{4.2.78}$$

with the following operators of well-defined twist

$$\begin{aligned}
G^{\mathrm{tw4(iv)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & g_{\alpha\beta}\Box\int_0^1 \mathrm{d}\lambda\frac{1-\lambda}{2\lambda}G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\
G^{\mathrm{tw4(iv)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & x_{(\alpha}\partial_{\beta)}\Box\int_0^1 \mathrm{d}\lambda\frac{(1-\lambda)^2}{2\lambda}G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{\mathrm{tw6(iv)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \\
G^{\mathrm{tw4(iv)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & x_{(\alpha}\partial_{\beta)}\Box\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda}+\frac{\ln\lambda}{\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\
& - G^{\mathrm{tw6(iv)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}$$

and

$$\begin{aligned}
G^{\mathrm{tw6(iv)a}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & -\frac{1}{2}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda}+\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\
G^{\mathrm{tw6(iv)b}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & -\frac{1}{2}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda}{\lambda}-\frac{1-\lambda^2}{4\lambda}+\frac{\ln\lambda}{2\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\
G^{\mathrm{tw6(iv)d1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & -\frac{1}{4}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda}+\frac{\ln\lambda}{\lambda}+\frac{\ln^2\lambda}{\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\
G^{\mathrm{tw6(iv)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & \frac{1}{2}x_\alpha x_\beta\Box^2\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda}+\frac{\ln\lambda}{\lambda}+\frac{\ln^2\lambda}{\lambda}\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}$$

being related to the scalar operator, as well as

$$\begin{aligned}
G^{\mathrm{tw4(iv)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & -g_{\alpha\beta}\partial^\mu\int_0^1 \mathrm{d}\lambda\frac{1}{\lambda}G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,, \\
G^{\mathrm{tw4(iv)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & -2x_{(\alpha}\partial_{\beta)}\partial^\mu\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda}+\frac{\ln\lambda}{\lambda}\Big)G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\
& - G^{\mathrm{tw6(iv)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,, \\
G^{\mathrm{tw4(iv)e1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & x_{(\alpha}\partial_{\beta)}\partial^\mu\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda}G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{\mathrm{tw6(iv)e1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}$$

and

$$
\begin{aligned}
G^{\mathrm{tw5(iv)d}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {}& x_{(\alpha}\big(\delta^{\mu}_{\beta)}(x\partial) - x^{\mu}\partial_{\beta)}\big)\Box \int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
& - G^{\mathrm{tw6(iv)d3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) - G^{\mathrm{tw6(iv)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw6(iv)c}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}x_\alpha x_\beta \Box\partial^{\mu}\int_0^1 \mathrm{d}\lambda\frac{(1-\lambda)^2}{\lambda} G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)d2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}x_\alpha x_\beta \Box\partial^{\mu}\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda} + \frac{\ln^2\lambda}{\lambda}\Big) G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)d4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\, x_\alpha x_\beta \Box\partial^{\mu}\int_0^1 \mathrm{d}\lambda\,\frac{\ln^2\lambda}{\lambda}\, G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)e1}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\,\frac{1}{2}x_\alpha x_\beta \Box\partial^{\mu}\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)e3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= x_\alpha x_\beta \Box\partial^{\mu}\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^{+}_{\mu}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the symmetric vector operator,

$$
\begin{aligned}
G^{\mathrm{tw5(iv)e}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {}& x_{(\alpha}\big(\delta^{\nu}_{\beta)}(x\partial) - x^{\nu}\partial_{\beta)}\big)\partial^{\mu}\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda}\, G_{(\mu\nu)}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
& - G^{\mathrm{tw6(iv)e3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) - G^{\mathrm{tw6(iv)e4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) - G^{\mathrm{tw6(iv)e5}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
G^{\mathrm{tw6(iv)e4}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {}& x_\alpha x_\beta \partial^{\mu}\partial^{\nu}\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda} G_{(\mu\nu)}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)f}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {}& -\, x_\alpha x_\beta \partial^{\mu}\partial^{\nu}\int_0^1 \mathrm{d}\lambda(1-\lambda) G_{(\mu\nu)}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the symmetric tensor operator and, furthermore,

$$
\begin{aligned}
G^{\mathrm{tw4(iv)f}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= g_{\alpha\beta}(x\partial)\int_0^1 \mathrm{d}\lambda\frac{1}{2\lambda} G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw4(iv)h}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\, x_{(\alpha}\partial_{\beta)}\int_0^1 \mathrm{d}\lambda\,\lambda G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{\mathrm{tw6(iv)h}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,\\
G^{\mathrm{tw4(iv)e2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\, x_{(\alpha}\partial_{\beta)}\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda} G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} - G^{\mathrm{tw6(iv)e2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,,
\end{aligned}
$$

and

$$
\begin{aligned}
G^{\mathrm{tw6(iv)h}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\,\frac{1}{4}x_\alpha x_\beta \Box\int_0^1 \mathrm{d}\lambda\frac{1-\lambda^2}{\lambda} G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)e2}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}x_\alpha x_\beta \Box\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)k}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \frac{1}{2}x_\alpha x_\beta \Box\int_0^1 \mathrm{d}\lambda\,(1-\lambda)\, G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\
G^{\mathrm{tw6(iv)e5}}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= -\, x_\alpha x_\beta \Box\int_0^1 \mathrm{d}\lambda\Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda}\Big) G^{\rho}{}_{\rho}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,,
\end{aligned}
$$

being related to the trace of the symmetric tensor operator.

Here, it should be remarked that, contrary to the former cases (i) – (iii), the trace terms (4.2.77) contain operators having the same twist, $\tau = 4$, as the primary operator of symmetry class (iv). However, these operators being multiplied by either $g_{\alpha\beta}$ or x_α and x_β are scalar or vector operators corresponding to symmetry type (i). Whereas the local operators resulting from the twist–4 tensor operator of symmetry class (iv) are contained in the tensor space $\mathbf{T}(\frac{n+2}{2}, \frac{n-2}{2}) \oplus \mathbf{T}(\frac{n-2}{2}, \frac{n+2}{2})$, the local operators of the twist–4 scalar and vector operators are totally symmetric traceless tensors and, therefore, contained in the tensor space $\mathbf{T}(\frac{n}{2}, \frac{n}{2})$.

Determination of the complete symmetric light–cone tensor operators

Now, having finished the decomposition of a general 2nd rank tensor operator, let us sum up the remaining symmetric tensor operators of twist greater than 3, which appear in the trace terms of the symmetric twist operators with Young symmetry (i), (ii) as well as (iv), to complete twist–4, twist–5 and twist–6 operators.

The 'scalar part' of the twist–4 light–ray tensor operator is given by

$$
\begin{aligned}
G^{\mathrm{tw4,s}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = -g_{\alpha\beta}\Big\{ & \tfrac{1}{2}\Box \int_0^1 \mathrm{d}\lambda\, \lambda(\ln\lambda) G(\kappa_1\lambda x, \kappa_2\lambda x) \\
& - (x\partial)\int_0^1 \mathrm{d}\lambda \frac{1}{2\lambda} G^{\rho}{}_{\rho}(\kappa_1\lambda x, \kappa_2\lambda x) + \partial^\mu \int_0^1 \mathrm{d}\lambda\, \lambda G^{+}_{\mu}(\kappa_1\lambda x, \kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}},
\end{aligned}
\tag{4.2.79}
$$

and the 'vector part' of the twist–4 light–ray tensor operator reads

$$
\begin{aligned}
& G^{\mathrm{tw4,v}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = -\tilde{x}_{(\alpha} \times \\
& \quad \Big\{\int_0^1 \mathrm{d}\lambda \Big\{\Big(\tfrac{1-\lambda^2}{2\lambda} + \lambda\ln\lambda\Big)\partial_{\beta)} + \tfrac{1}{4}x_{\beta)}\Big(\tfrac{1-\lambda^2}{\lambda} + \lambda\ln\lambda + \tfrac{\ln\lambda}{\lambda}\Big)\Box\Big\}\Box G(\kappa_1\lambda x, \kappa_2\lambda x) \\
& \quad + \int_0^1 \frac{\mathrm{d}\lambda}{\lambda}\Big\{\partial_{\beta)} + \tfrac{1}{2}x_{\beta)}(\ln\lambda)\Box\Big\} G^{\rho}{}_{\rho}(\kappa_1\lambda x, \kappa_2\lambda x) \\
& \quad - \int_0^1 \frac{\mathrm{d}\lambda}{\lambda}\Big\{2\big(1-\lambda^2\big)\partial_{\beta)} + \tfrac{1}{2}x_{\beta)}\big(1-\lambda^2 + 2\ln\lambda\big)\Box\Big\}\partial^\mu G^{+}_{\mu}(\kappa_1\lambda x, \kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}.
\end{aligned}
\tag{4.2.80}
$$

Furthermore, the contributions to the complete 'vector part' of the twist–5 light–ray tensor operator are constructed by means of the Young tableau (ii), and their local operators are antisymmetric[6] traceless tensors contained in the space $\mathbf{T}(\frac{n}{2}, \frac{n-2}{2}) \oplus \mathbf{T}(\frac{n-2}{2}, \frac{n}{2})$. This complete twist–5 tensor operator is given by

$$
\begin{aligned}
G^{\mathrm{tw5,v}}_{(\alpha\beta)}(\kappa_1\tilde{x}, \kappa_2\tilde{x}) = -\tilde{x}_{(\alpha}\Big\{ & \int_0^1 \mathrm{d}\lambda \Big\{\frac{(1-\lambda)^2}{2\lambda}\big(\delta^\nu_{\beta)}(x\partial) - x^\nu\partial_{\beta)}\big)\Box + \Big(\frac{1-\lambda}{\lambda} + \frac{\ln\lambda}{\lambda}\Big)x_{\beta)}\partial^\nu\Box \\
& - \Big(\frac{1-\lambda}{\lambda} - \frac{1-\lambda^2}{4\lambda} + \frac{1}{2\lambda}\ln\lambda\Big)x_{\beta)}\Box^2 x^\nu\Big\} G^{+}_{\nu}(\kappa_1\lambda x, \kappa_2\lambda x) \\
- \int_0^1 \mathrm{d}\lambda \Big\{ & \frac{1-\lambda^2}{\lambda}\big(\delta^\nu_{\beta)}(x\partial) - x^\nu\partial_{\beta)} - x_{\beta)}\partial^\nu\big)\partial^\mu - \Big(\frac{1-\lambda^2}{2\lambda} + \frac{\ln\lambda}{\lambda}\Big)x_{\beta)}\Box\big(\partial^\mu x^\nu - g^{\mu\nu}\big)\Big\} \\
& \times G_{(\mu\nu)}(\kappa_1\lambda x, \kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}.
\end{aligned}
\tag{4.2.81}
$$

---

[6] Note that antisymmetry is not in α and β but results from the differential operator!

The 'scalar part' of the twist–6 light–ray tensor operator has Young symmetry (i). Thus, the local twist–6 operators are totally symmetric traceless tensors lying in the space $\mathbf{T}(\frac{n-2}{2},\frac{n-2}{2})$. This nonlocal twist–6 operator is given by

$$
\begin{aligned}
G^{\rm tw6,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = -\tilde{x}_\alpha\tilde{x}_\beta\Big\{ & \frac{1}{2}\,\Box^2\int_0^1 \mathrm{d}\lambda\Big(\tfrac{1-\lambda}{\lambda}+\tfrac{1+\lambda}{2\lambda}\ln\lambda\Big)G(\kappa_1\lambda x,\kappa_2\lambda x)\\
&+\frac{1}{2}\,\Box\int_0^1 \mathrm{d}\lambda\,\Big(\tfrac{1-\lambda}{\lambda}+\tfrac{\ln\lambda}{\lambda}\Big)\Big(G^\rho{}_\rho(\kappa_1\lambda x,\kappa_2\lambda x)-2\partial^\mu G^+_\mu(\kappa_1\lambda x,\kappa_2\lambda x)\Big)\\
&-\partial^\mu\partial^\nu\int_0^1 \mathrm{d}\lambda\,\tfrac{1-\lambda}{\lambda}\,G_{(\mu\nu)}(\kappa_1\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}\,.
\end{aligned}
\tag{4.2.82}
$$

Now, we may pick up all the contributions to the symmetric (traceless) tensor operator. Together with the twist–2 part, Eq. (4.2.20), the twist–3 part, Eq. (4.2.51), and the twist–4 part, Eq. (4.2.76), we finally obtain the following complete decomposition of the *symmetric light–cone tensor operator* (compare Eq. (4.2.7)):

$$
\begin{aligned}
G_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = {} & G^{\rm tw2(i)}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\rm tw3(ii)}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\rm tw4(iv)}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
&+G^{\rm tw4,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\rm tw4,v}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\\
&+G^{\rm tw5,v}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G^{\rm tw6,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})\,.
\end{aligned}
\tag{4.2.83}
$$

Let us remark that by construction the terms of the first line are traceless, whereas the traces of the third line vanish because of antisymmetry and $\tilde{x}^2=0$ on the light–cone, respectively. However, the traces of the second line do not vanish but restore the trace of the tensor operator,

$$
g^{\alpha\beta}G_{\alpha\beta}(\kappa_1\tilde{x},\kappa_2\tilde{x}) = g^{\alpha\beta}G^{\rm tw4,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+g^{\alpha\beta}G^{\rm tw4,v}_{(\alpha\beta)}((\kappa_1\tilde{x},\kappa_2\tilde{x})). \tag{4.2.84}
$$

This may be proven explicitly by taking the trace of Eqs. (4.2.79) and (4.2.80), using $(x\partial)\cong\lambda\partial_\lambda$, performing partial integrations and observing the equality

$$
\big(g^{\alpha\beta}-2\partial^\alpha x^\beta+(1/2)\Box x^\alpha x^\beta\big)G_{(\alpha\beta)}(\kappa_1\lambda x,\kappa_2\lambda x)\big|_{\lambda=0}=0\,. \tag{4.2.85}
$$

Herewith, the twist decomposition of a generic bilocal 2nd rank light–ray tensor operator is completed. In the next Section we show how this general formalism may be applied also to tri– and multi–local light–ray operators.

4.2.5 Conclusions

Let us shortly summarize our results. We determined in a systematic way the various contributions of definite twist for a generic bilocal 2nd rank tensor operator $G_{\alpha\beta}(\kappa_1 x,\kappa_2 x)$ as well as its symmetric and antisymmetric parts $G_{(\alpha\beta)}(\kappa_1 x,\kappa_2 x)$ and $G_{[\alpha\beta]}(\kappa_1 x,\kappa_2 x)$, respectively. In addition, we determined the related vector and scalar operators which occur by truncating with x^β and $x^\alpha x^\beta$ or $g^{\alpha\beta}$, respectively. By projection onto the light–cone we obtained the decomposition of a generic bilocal light–ray tensor operator of 2nd rank – and its reductions to (anti)symmetric tensors as well as the related vector and scalar operators – into operators of definite twist. In order to make the main results quite obvious we summarize them in Table 4.1 and 4.2.[7] There, we indicate the different twist contributions to $G_{\alpha\beta}(\kappa_1\tilde{x},\kappa_2\tilde{x})=G_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})+G_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ by their symmetry type (i) up to (iv) and the number of the corresponding equations; in the second half of the Tables the contributions from the trace terms are shown.

Eq.	YT	$G_{(\alpha\beta)}(\kappa_1\tilde{x},\kappa_2\tilde{x})$	$G_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$
4.2.20	(i)	$\tau = 2$	–
4.2.51	(ii)	$\tau = 3$	–
4.2.48	(ii)	–	$\tau = 3$
4.2.67	(iii)	–	$\tau = 4$
4.2.76	(iv)	$\tau = 4$	–
4.2.79	(i)	$\tau = 4$	–
4.2.80	(i)	$\tau = 4$	–
4.2.69	(i)	–	$\tau = 4$
4.2.81	(ii)	$\tau = 5$	–
4.2.70	(ii)	–	$\tau = 5$
4.2.82	(i)	$\tau = 6$	–

Table 4.1: Twist decomposition of general tensor operators $G_{\alpha\beta} = G_{(\alpha\beta)} + G_{[\alpha\beta]}$.

Eq.	YT	$G_{(\alpha\bullet)}(\kappa_1\tilde{x},\kappa_2\tilde{x})$	$G_{[\alpha\bullet]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$
4.2.25	(i)	$\tau = 2$	–
4.2.53	(ii)	$\tau = 3$	–
4.2.54	(ii)	–	$\tau = 3$
4.2.59	(i)	$\tau = 4$	–
4.2.57	(i)	–	$\tau = 4$

Table 4.2: Twist decomposition of general vector operators $G_\alpha = G_{(\alpha\bullet)} + G_{[\alpha\bullet]}$.

The present study made also obvious how difficult an exact treatment of higher twist contributions will be. There is a complicate interplay between the higher twists resulting from different symmetry types and the corresponding trace terms. The results also suffer from the equations of motion (EOM) which are frequently used in the applications because they do not contribute between physical states [111]. The twist decomposition of these EOM operators

$$\begin{aligned} {}^{\rm EOM}\mathcal{O}_\Gamma(\kappa_1\tilde{x},\kappa_2\tilde{x}) &= \bar{\psi}(\kappa_1\tilde{x})\Gamma U(\kappa_1\tilde{x},\kappa_2\tilde{x})\Big[\mathrm{i}\overrightarrow{\not{D}} - m\Big]\psi(\kappa_2\tilde{x}) \\ &\quad - \bar{\psi}(\kappa_1\tilde{x})\Big[\mathrm{i}\overleftarrow{\not{D}} + m\Big]\Gamma U(\kappa_1\tilde{x},\kappa_2\tilde{x})\psi(\kappa_2\tilde{x}), \end{aligned}$$

is analogous to the corresponding quark operators $\mathcal{O}_\Gamma(\kappa_1\tilde{x},\kappa_2\tilde{x})$ which is given in [96, 63]. The only difference is that for the equation of motion operators the canonical dimension and the twist raises by one unit.

4.3 Extension to trilocal tensor operators

The procedure discussed in the previous Section for bilocal gluon operators may be extended to trilocal operators, too. These operators occur in the higher twist contributions to light–cone dominated processes and, additionally, as counterparts in the renormalization of such

---

[7] After substitution $\tau \to \tau - 1$ in all formulas for $G_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$ in Table 4.1 we get the corresponding twist decomposition for the bilocal operator $M_{[\alpha\beta]}(\kappa_1\tilde{x},\kappa_2\tilde{x})$. If we substitute $\mathrm{d}\lambda\,\lambda \to \mathrm{d}\lambda$ in the formulas for the vector operator $G_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})$ in Table 4.2, one obtains the twist decomposition for the operator $O_\alpha(\kappa_1\tilde{x},\kappa_2\tilde{x})$.

higher twist operators which have been obtained in Section 4.2. In the following we consider various trilocal operators of minimal twist–3 and twist–4. The local versions of such higher twist operators have been considered already in the early days of QCD in systematic studies of deep inelastic scattering [68] and on behalf of giving some parton interpretation for the distribution amplitudes [111, 53, 117, 76, 38]. Various studies of local twist–4 operators determined their anomalous dimension matrices and the behaviour of their structure functions [122, 109, 100, 101, 75]. The relevance of local twist–3 operators for the structure function g_2 of polarized deep inelastic scattering is also well-known [2, 117, 37, 113, 89, 90, 91, 92]. If written nonlocal all these operators are necessarily multilocal. However, nonlocal higher twist operators have been extensively used in the study of the structure function g_2 [9, 61, 62] and of the photon and vector meson wave functions [10, 11, 14, 15].

Here, our main interest is not to give an exhaustive study of the various scattering processes but to present the twist decomposition of some characteristic multilocal operators.

4.3.1 General trilocal tensor operators: Quark–antiquark–gluon, four–quark and four–gluon operators

This Subsection is devoted to the consideration of trilocal operators which share the same twist decomposition as the gluon operators $G_{\alpha\beta}$ (or their (anti)symmetric parts). We call them unconstrained because their truncation with $x^\alpha x^\beta$ does not vanish identically. In order to compactify notations we introduce the following abbreviations:

$$\Gamma^i = \{1, \gamma_5\}, \quad \Gamma^i_\alpha = \{\gamma_\alpha, \gamma_5\gamma_\alpha\}, \quad \Gamma^i_{\alpha\beta} = \{\sigma_{\alpha\beta}, \gamma_5\sigma_{\alpha\beta}\}, \quad F^i_{\mu\nu} = \{F_{\mu\nu}, \widetilde{F}_{\mu\nu}\}\,.$$

Quark–antiquark–gluon operators

First, we consider the following quark–antiquark–gluon operators,[8]

$$V^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \bar{\psi}(\kappa_1 x) U(\kappa_1 x, \tau x) \Gamma^{i\,\rho}_{\alpha} F^j_{\beta\rho}(\tau x) U(\tau x, \kappa_2 x) \psi(\kappa_2 x)\,, \tag{4.3.1}$$

and

$$W^{ij}_{[\alpha\beta]}(\kappa_1 x, \tau x, \kappa_2 x) = \bar{\psi}(\kappa_1 x) U(\kappa_1 x, \tau x) \Gamma^i F^j_{\alpha\beta}(\tau x) U(\tau x, \kappa_2 x) \psi(\kappa_2 x)\,, \tag{4.3.2}$$

whose minimal twist is $\tau_{\min} = 3$ and 4, respectively; they have the same symmetry as the gluon tensor operators $G_{\alpha\beta}$ and $G_{[\alpha\beta]}$, respectively. The trilocal operators $V^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x)$ are inalienable for a systematic study of the chiral-odd twist-3 distribution functions $h_L(x)$ and $e(x)$ [79, 94, 95, 13] which give information about the quark-gluon correlations in the nucleon.

In order to be able to apply the procedure of twist decomposition as well as the results of Section 4.2 we have to verify the local structure of these trilocal operators. Let us study, for example, the first of the above quark–gluon operators, $V^{11}_{\alpha\beta}$, in some detail and Taylor expand its three local fields around $y = 0$:

---

[8] For simplicity we restrict to the nonsinglet case and suppress flavour matrices.

$$
\begin{aligned}
&V^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \\
&\quad = \bar\psi(\kappa_1 x) U(\kappa_1 x, 0) U(0, \tau x) \sigma_\alpha{}^\rho F_{\beta\rho}(\tau x) U(\tau x, 0) U(0, \kappa_2 x) \psi(\kappa_2 x) \\
&\quad = \sum_{n_1, m, n_2 = 0}^{\infty} \frac{\kappa_1^{n_1} \tau^m \kappa_2^{n_2}}{n_1! m! n_2!} \Big[\bar\psi(y) \Big(\overleftarrow{D}_y x \Big)^{n_1} \Big] \sigma_\alpha{}^\rho \Big[(x D_z)^m F_{\beta\rho}(z) \Big] \Big[\Big(x \overrightarrow{D}_y \Big)^{n_2} \psi(y) \Big] \Big|_{y=z=0} \\
&\quad = \sum_{N=0}^{\infty} \frac{1}{N!} \bar\psi(0) \left(\sum_{n=0}^{N} \sum_{\ell=0}^{n} \binom{N}{n} \binom{n}{\ell} \Big(\kappa_1 \overleftarrow{D} x \Big)^{n-\ell} \sigma_\alpha{}^\rho \Big[(\tau x D)^{N-n} F_{\beta\rho}(0) \Big] \Big(\kappa_2 x \overrightarrow{D} \Big)^{\ell} \right) \psi(0) \, ,
\end{aligned}
$$

where the left and right derivatives are given by Eqs. (2.2.4), and

$$
D_\mu F_{\alpha\beta} = \partial_\mu F_{\alpha\beta} + \mathrm{i} g [A_\mu, F_{\alpha\beta}] \, . \tag{4.3.3}
$$

This leads to the following expansion into local tensor operators

$$
V^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \sum_{N=0}^{\infty} \frac{1}{N!} x^{\mu_1} \ldots x^{\mu_N} V^{11}_{\alpha\beta\mu_1 \ldots \mu_N}(\kappa_1, \tau, \kappa_2) \, , \tag{4.3.4}
$$

where

$$
V^{11}_{\alpha\beta\mu_1 \ldots \mu_N}(\kappa_1, \tau, \kappa_2) \equiv \bar\psi(0) \sigma_\alpha{}^\rho \, \mathbf{D}_{[\beta\rho]\mu_1 \ldots \mu_N}(\kappa_1, \tau, \kappa_2) \psi(0) \, , \tag{4.3.5}
$$

and the field strength–dependent generalized covariant derivative of Nth order is given by

$$
\begin{aligned}
&\mathbf{D}_{[\beta\rho]\mu_1 \ldots \mu_N}(\kappa_1, \tau, \kappa_2) = \\
&\quad = N! \sum_{n=0}^{N} \sum_{\ell=0}^{n} \frac{\kappa_1^{n-\ell}}{(n-\ell)!} \overleftarrow{D}_{\mu_1} \ldots \overleftarrow{D}_{\mu_{n-\ell}} \frac{\tau^{N-n}}{(N-n)!} \Big[D_{\mu_{n+1}} \ldots D_{\mu_N} F_{\beta\rho}(0) \Big] \frac{\kappa_2^\ell}{\ell!} \overrightarrow{D}_{\mu_{n-\ell+1}} \ldots \overrightarrow{D}_{\mu_n} .
\end{aligned} \tag{4.3.6}
$$

Obviously, if the field strength were not present this operation would reduce to the product $\overleftrightarrow{D}_{\mu_1}(\kappa_1, \kappa_2) \ldots \overleftrightarrow{D}_{\mu_N}(\kappa_1, \kappa_2)$ of generalized derivatives (2.2.4). Furthermore, for $\tau = 0$ we are left with a more simple expression which, however, cannot be reduced to the Nth power of some extended derivative because F is equipped with some matrix structure. Anyway, the local tensor operators (4.3.5) of rank $N+2$ with canonical dimension $d = N+5$, which are given as a sum of $\frac{1}{2}(N+1)(N+2)$ terms, have to be decomposed according to their geometric twist. In principle, this has to be done term-by-term. But, because of the linearity of that procedure we are not required to do this for any term explicitly.

Due to the same general local structure of $V^{ij}_{\alpha\beta}$ and $W^{ij}_{\alpha\beta}$ – which is governed by Taylor expansions completely analogous to Eq. (4.3.4) – their decomposition into terms of definite twist leads to the same expressions as we obtained in Section 4.2 but with $G_{\alpha\beta}$ and $G_{[\alpha\beta]}$ exchanged by $V^{ij}_{\alpha\beta}$ and $W^{ij}_{\alpha\beta}$, respectively. The only difference consists in a shift of any twist by one unit, $\tau \rightarrow \tau + 1$, in the various inputs of Table 4.1 and 4.2.

In contrast to the operators V and W the quark–antiquark–gluon operators

$$
\mathcal{O}^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = x^\rho \, \bar\psi(\kappa_1 x) U(\kappa_1 x, \tau x) \Gamma^i_\alpha F^j_{\beta\rho}(\tau x) U(\tau x, \kappa_2 x) \psi(\kappa_2 x) \tag{4.3.7}
$$

and

$$
\mathcal{O}'^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = x^\rho x^\sigma \bar\psi(\kappa_1 x) U(\kappa_1 x, \tau x) \Gamma^i_{\alpha\rho} F^j_{\beta\sigma}(\tau x) U(\tau x, \kappa_2 x) \psi(\kappa_2 x) \tag{4.3.8}
$$

are special, namely, despite of their arbitrary symmetry with respect to α and β they vanish identically if multiplied by x^β and x^α or x^β, respectively. Therefore, they show some peculiarities which will be treated in Subsection 4.2.

Four–quark and four–gluon operators

Let us now consider trilocal operators which are built up from four quark or four gluon fields. We denote them by ${}^{I}\mathcal{Q}^{ij}_{\alpha\beta}$ with $I = 0$ and $I = 2$, respectively (here, I counts the number of external x's of the operators):

$$
{}^{I=0}\mathcal{Q}^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \big(\,\bar\psi(\kappa_1 x)\Gamma^i_\alpha U(\kappa_1 x, \tau x)\psi(\tau x)\big)\big(\,\bar\psi(\tau x)U(\tau x, \kappa_2 x)\Gamma^j_\beta \psi(\kappa_2 x)\big) \tag{4.3.9}
$$

as well as

$$
\begin{aligned}
{}^{I=2}\mathcal{Q}^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) &= x^\rho x^\sigma\, {}^{I=2}\mathcal{Q}^{ij}_{\alpha\rho\sigma\beta}(\kappa_1 x, \tau x, \kappa_2 x) \qquad (4.3.10)\\
&= x^\rho x^\sigma \Big(F_{\alpha\mu}(\kappa_1 x)U(\kappa_1 x, \tau x)F^{i\mu}_{\rho}(\tau x)\Big)\Big(F^{j\nu}_{\sigma}(\tau x)U(\tau x, \kappa_2 x)F_{\beta\nu}(\kappa_2 x)\Big)\,.
\end{aligned}
$$

Both sets of operators have minimal twist $\tau_{\min} = 4$. Their local versions have been already considered by various authors, see, e.g., [75]. The explicit form of the local operators will be given below. Because there are no restrictions concerning the free indices α and β the Young patterns (i) – (iv) are involved. The twist decomposition may be performed along the lines of Section 4.2 and, as in the foregoing cases, the outcome will be almost the same as for $G_{\alpha\beta}$. Again, the difference is that now the value of twist raises by two units, $\tau \to \tau 2$, relative to the gluon operators. In addition, because of the change in the external x–factors (not being accompanied by λ) the measures of the λ–integrations have to be changed according to (see also Subsection 4.2)

$$
\mathrm{d}\lambda \longrightarrow \mathrm{d}\lambda\, \lambda^I\,. \tag{4.3.11}
$$

Open, up to now, is the structure of the local operators. Let us study in detail the first of the four–quark operators, ${}^{I=0}\mathcal{Q}^{11}_{\alpha\beta}$, whose Taylor expansion may be written in two equivalent ways:

$$
\begin{aligned}
&{}^{I=0}\mathcal{Q}^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \qquad (4.3.12)\\
&\quad= \big(\,\bar\psi(\kappa_1 x)\gamma_\alpha U(\kappa_1 x, 0)U(0, \tau x)\psi(\tau x)\big)\;\big(\bar\psi(\tau x)U(\tau x, 0)U(0, \kappa_2 x)\gamma_\beta \psi(\kappa_2 x)\big)\\
&\quad= \sum_{n_1, m, n_2 = 0}^{\infty} \frac{\kappa_1^{n_1}\tau^m\kappa_2^{n_2}}{n_1! m! n_2!}\Big[\bar\psi(y)\Big(\overleftarrow{D}_y x\Big)^{n_1}\Big]\gamma_\alpha\Big[(xD_z)^m\big(\psi(z)\bar\psi(z)\big)\Big]\gamma_\beta\Big[\Big(x\overrightarrow{D}_y\Big)^{n_2}\psi(y)\Big]\Big|_{y=z=0}\\
&\quad= \sum_{n_1=0}^{\infty}\frac{1}{n_1!}\Big(\bar\psi(y)\Big(x\overleftrightarrow{D}_y(\kappa_1, \tau)\Big)^{n_1}\gamma_\alpha\psi(y)\Big)\Big|_{y=0}\sum_{n_2=0}^{\infty}\frac{1}{n_2!}\Big(\bar\psi(z)\Big(x\overleftrightarrow{D}_z(\tau, \kappa_2)\Big)^{n_2}\gamma_\beta\psi(z)\Big)\Big|_{z=0},
\end{aligned}
$$

where the two possibilities also use different generalized covariant derivatives. The left, right and left-right derivatives are given by Eqs. (2.2.4), and, treating $\psi(0)\bar\psi(0)$ as a matrix in the group algebra, another form of the derivative obtains:

$$
\begin{aligned}
D_\mu\left(\psi(0)\bar\psi(0)\right) &= \partial_\mu\left(\psi(0)\bar\psi(0)\right) + \mathrm{i}g[A_\mu, \left(\psi(0)\bar\psi(0)\right)]\\
&= \left(\overrightarrow{D}_\mu\psi(0)\right)\bar\psi(0) + \psi(0)\left(\bar\psi(0)\overleftarrow{D}_\mu\right)\,.
\end{aligned}
$$

This leads to the following expansion into local tensor operators

$$
{}^{I=0}\mathcal{Q}^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \sum_{N=0}^{\infty}\frac{1}{N!}x^{\mu_1}\ldots x^{\mu_N}\, {}^{I=0}\mathcal{Q}^{ij}_{\alpha\beta\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2)\,, \tag{4.3.13}
$$

where

$$ {}^{I=0}\mathcal{Q}^{ij}_{\alpha\beta\mu_1\ldots\mu_N}(\kappa_1,\tau,\kappa_2) \equiv \bar\psi(0)\gamma_\alpha \mathbf{D}_{\mu_1\ldots\mu_N}(\kappa_1,\tau,\kappa_2)\gamma_\beta\psi(0)\,, \tag{4.3.14} $$

and the quark field–dependent generalized covariant derivative of Nth order is given by

$$ \begin{aligned} \mathbf{D}_{\mu_1\ldots\mu_N}(\kappa_1,\tau,\kappa_2) &= \sum_{n=0}^{N} \binom{N}{n} \overset{\leftrightarrow}{D}_{\mu_1}\ldots\overset{\leftrightarrow}{D}_{\mu_n}\psi(0)\times\bar\psi(0)\overset{\leftrightarrow}{D}_{\mu_{n+1}}\ldots\overset{\leftrightarrow}{D}_{\mu_N} \\ &= \sum_{n=0}^{N} N! \sum_{\ell=0}^{n} \overset{\leftarrow}{D}_{\mu_1}\ldots\overset{\leftarrow}{D}_{\mu_{n-\ell}} \frac{\kappa_1^{n-\ell}\tau^\ell}{(n{-}\ell)!\ell!} \vec{D}_{\mu_{n-\ell+1}}\ldots\vec{D}_{\mu_n}\psi(0) \\ &\qquad\times \sum_{k}^{N-n} \bar\psi(0)\overset{\leftarrow}{D}_{\mu_{n+1}}\ldots\overset{\leftarrow}{D}_{\mu_{n+k}} \frac{\tau^k\kappa_2^{N-n-k}}{k![N{-}n{-}k)!} \vec{D}_{\mu_{n+k+1}}\ldots\vec{D}_{\mu_N}\,. \end{aligned} \tag{4.3.15} $$

Analogous generalized covariant derivatives occur for the four–gluon operators. The only difference will be that instead of the quark fields corresponding gluon field strengths appear in (4.3.15), and that the derivatives are to be taken in the adjoint representation. Despite having a complicated structure this does not matter in the twist decomposition of the trilinear operators ${}^{I}\mathcal{Q}^{ij}_{\alpha\beta}(\kappa_1 x,\tau x,\kappa_2 x)$.

Let us finish this Subsection with a short remark concerning so-called *four-particle* operators which also have been considered in the literature for local operators. These operators have the general structure, cf. [111, 53]

$$ (\bar\psi U\psi)(\bar\psi U\psi), \quad (\bar\psi UFUFU\psi) \quad \text{or} \quad (FUFUFUF), $$

with any field being located at a different point $\kappa_i x, i=1,\ldots,4$. The expansion into local tensors is obtained in the same way as above, leading to a general expression of the following form:

$$ {}^{I}\mathcal{R}_{\alpha\beta}(\kappa_i x) = \sum_{N=0}^{\infty}\frac{1}{N!}x^{\mu_1}\ldots x^{\mu_N}x^\rho\ldots x^\sigma\,{}^{I}\mathcal{R}_{\alpha\beta\rho\ldots\sigma\mu_1\ldots\mu_N}(\kappa_i)\,, \tag{4.3.16} $$

where the terms with equal value of N are given as well-defined sums of local tensor operators. Any of these operators decompose in the same manner into irreducible representations of the Lorentz group. The latter result is immediately related to the fact that all indices $\mu_1\ldots\mu_N$ are to be symmetrized, i.e., lying in the first row of any relevant Young tableau! Different symmetry classifications solely depend on the distribution of the remaining indices $\alpha\beta\rho\ldots\sigma$ – the others being somehow truncated – to the various Young tableaux. As long as only up to two free indices $\alpha\beta$ are relevant the twist decomposition takes place according to the results of Section 4.2, eventually modified by the power λ^I which is related to the number of external x's in ${}^{I}\mathcal{R}$. In addition, the twist of the various components may be shifted by some (equal) amount.

4.3.2 Constrained trilocal operators: Shuryak-Vainshtein– and three–gluon operators

Now we consider the special quark–antiquark–gluon operator $\mathcal{O}^{ij}_{\alpha\beta}$, Eq. (4.3.7), and related three–gluon operators both having minimal twist–3; they will be denoted by ${}^{I}\mathcal{O}^{ij}_{\alpha\beta}(\kappa_1 x,\tau x,\kappa_2 x)$ with $I=0$ and $I=1$, respectively. The quark–antiquark–gluon operators are given by

$$ \begin{aligned} {}^{I=0}\mathcal{O}^{ij}_{\alpha\beta}(\kappa_1 x,\tau x,\kappa_2 x) &= x^\rho\,{}^{I=0}\mathcal{O}^{ij}_{\alpha\beta\rho}(\kappa_1 x,\tau x,\kappa_2 x) \\ &= x^\rho\,\bar\psi(\kappa_1 x)U(\kappa_1 x,\tau x)\Gamma^i_\alpha F^j_{\beta\rho}(\tau x)U(\tau x,\kappa_2 x)\psi(\kappa_2 x)\,, \end{aligned} \tag{4.3.17} $$

where again some possible flavour structure has been suppressed. These operators are related to the following generalizations of the so-called Shuryak-Vainshtein operators [117, 9],

$$S^{\pm}_{\beta}(\kappa_1 x, \tau x, \kappa_2 x) = x^{\alpha}\big({}^{I=1}\mathcal{O}^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) \pm \mathrm{i}\,{}^{I=1}\mathcal{O}^{22}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x)\big)x^{\alpha},$$

which, in the flavour singlet case, mix with the following three–gluon operators:

$$\begin{aligned} {}^{I=1}\mathcal{O}^{i}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) &= x^{\rho}x^{\sigma}\,{}^{I=1}\mathcal{O}^{i}_{\alpha\beta\rho\sigma}(\kappa_1 x, \tau x, \kappa_2 x) \\ &= x^{\rho}x^{\sigma}\,F^{a}_{\alpha\nu}(\kappa_1 x)U^{ab}(\kappa_1 x, \tau x)F^{i\,bc}_{\beta\rho}(\tau x)U^{cd}(\tau x, \kappa_2 x)F^{d\nu}_{\sigma}(\kappa_2 x)\,, \end{aligned} \tag{4.3.18}$$

where $F^{ab}_{\alpha\beta}(x) \equiv f^{acb}F^{c}_{\alpha\beta}(x)$ and the phase factors are taken in the adjoint representation.

Because of their construction these special trilocal operators have the property

$$x^{\beta}\,{}^{I}\mathcal{O}^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) \equiv 0\,. \tag{4.3.19}$$

They contain twist–3 up to twist–6 contributions which, because of the antisymmetry of the gluon field strength, have to be determined by means of the Young tableaux (ii) – (iv). Because of (4.3.19) the scalar operators $x^{\alpha}x^{\beta}\,{}^{I}\mathcal{O}^{ij}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x)$ vanish identically.

The Taylor expansion of the operator ${}^{I=0}\mathcal{O}^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x)$ reads

$${}^{I=0}\mathcal{O}^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \sum_{N=0}^{\infty}\frac{1}{N!}x^{\mu_1}\ldots x^{\mu_N}x^{\rho}\,{}^{I=0}\mathcal{O}^{11}_{\alpha\beta\rho\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2)\,, \tag{4.3.20}$$

with

$${}^{I=0}\mathcal{O}^{11}_{\alpha\beta\rho\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2) \equiv \bar{\psi}(0)\gamma_{\alpha}\mathbf{D}_{[\beta\rho]\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2)\psi(0)\,, \tag{4.3.21}$$

where the field strength–dependent derivative already has been given by (4.3.6). The Taylor expansion of the related three–gluon operator obtains as follows:

$${}^{I=1}\mathcal{O}^{11}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) = \sum_{N=0}^{\infty}\frac{1}{N!}x^{\mu_1}\ldots x^{\mu_N}x^{\rho}x^{\sigma}\;{}^{I=1}\mathcal{O}^{A}_{\alpha\beta\rho\sigma\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2) \tag{4.3.22}$$

where

$${}^{I=1}\mathcal{O}^{11}_{\alpha\beta\rho\sigma\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2) \equiv F^{a}_{\alpha\nu}(0)\mathbf{D}^{ab}_{[\beta\rho]\mu_1\ldots\mu_N}(\kappa_1, \tau, \kappa_2)F^{b\nu}_{\sigma}(0)\,, \tag{4.3.23}$$

with the same field strength dependent generalized covariant derivative (4.3.6), but now in the adjoint representation. The generalization arbitrary values i and j is obvious.

As it became obvious by the above considerations all the nonlocal three–particle operators ${}^{I}\mathcal{O}^{ij}_{\alpha\beta}$ have the same general local structure and, therefore, decompose according to the same symmetry patterns. They only differ in the rank of the local tensor operators which is due to the external powers of x. Again, this must be taken into account by a change of the integration measure according to[9]

$$\mathrm{d}\lambda \longrightarrow \mathrm{d}\lambda\,\lambda^{I+1}\,.$$

From now on we omit the indices i and j.

---

[9] Observe, that because of notational simplicity in the explicit expressions below we shifted the variable $I \to I-1$.

In order to determine the twist–3 light–cone operators ${}^I\mathcal{O}_{\alpha\beta}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x})$ for $I = 0$ and $I = 1$, let us consider the following Young tableaux of symmetry pattern (iiB) $\begin{array}{|c|c|c|c|c|}\hline \rho & \mu_1 & \cdots & \mu_N & \alpha \\ \hline \beta \\ \cline{1-1}\end{array}$ and $\begin{array}{|c|c|c|c|c|c|}\hline \rho & \sigma & \mu_1 & \cdots & \mu_N & \alpha \\ \hline \beta \\ \cline{1-1}\end{array}$, respectively:

$$\begin{aligned}{}^I\mathcal{O}^{\rm tw3}_{\alpha\beta}(\kappa_1 x, \tau x, \kappa_2 x) &= \frac{1}{2}\int_0^1 \mathrm{d}\lambda\,\lambda^I\Big[\big(1+\lambda^2\big)\delta^\mu_\beta\partial_\alpha+\big(1-\lambda^2\big)\delta^\mu_\alpha\partial_\beta\Big]{}^I\overset{\circ}{\mathcal{O}}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\\ &= {}^I\mathcal{O}^{\rm tw3}_{[\alpha\beta]}(\kappa_1 x, \tau x, \kappa_2 x) + {}^I\mathcal{O}^{\rm tw3}_{(\alpha\beta)}(\kappa_1 x, \tau x, \kappa_2 x)\,; \qquad (4.3.24)\end{aligned}$$

here, and in the following, we write $\mathcal{O}_\mu \equiv x^\rho\mathcal{O}_{\rho\mu}$. Let us remark that, first, because of property (4.3.19), only the above mentioned Young tableaux contribute and, second, the difference between the expressions (4.3.24) and (4.2.43) result also from that property, namely the truncation by x^ρ in the second term of the integrand, $-x^\mu\partial_\alpha\partial_\beta{}^I\overset{\circ}{\mathcal{O}}_\mu(\kappa_1 x, \tau x, \kappa_2 x) = 2\delta^\mu_{(\alpha}\partial_{\beta)}{}^I\overset{\circ}{\mathcal{O}}_\mu(\kappa_1 x, \tau x, \kappa_2 x)$, after partial integrations leads to the above expression (4.3.24). In the case $I = 0$ this operator is in agreement with the expression given by Balitsky and Braun (cf. Eq. (5.14) of [9]). However, their operator lacks to be really of twist–3 since it is not traceless; therefore it contains also twist–4, twist–5 as well as twist–6 operators resulting from the trace terms.

Now, let us to project onto the light–cone. As in the case of the gluon tensor only the first terms, $k = 0, 1, 2$, in the expansion of ${}^I\overset{\circ}{\mathcal{O}}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)$ are to be taken into account. For the antisymmetric part of the twist–3 operator one gets

$${}^I\mathcal{O}^{\rm tw3}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\,\lambda^{2+I}\partial_{[\alpha}{}^I\mathcal{O}_{\beta]}(\kappa_1\lambda\tilde{x}, \tau\lambda\tilde{x}, \kappa_2\lambda\tilde{x}) - {}^I\mathcal{O}^{>({\rm ii})}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) \qquad (4.3.25)$$

where

$$\begin{aligned}{}^I\mathcal{O}^{>({\rm ii})}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) &= \frac{1}{2}\int_0^1 \mathrm{d}\lambda\,\lambda^{1+I}(1-\lambda)\big(2x_{[\alpha}\partial_{\beta]}\partial^\mu - x_{[\alpha}\delta^\mu_{\beta]}\Box\big)\\ &\qquad\times{}^I\mathcal{O}_\mu(\kappa_1\lambda x, \tau\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}. \qquad (4.3.26)\end{aligned}$$

The decomposition of ${}^I\mathcal{O}^{>({\rm ii})}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x})$ arises as

$${}^I\mathcal{O}^{>({\rm ii})}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) = {}^I\mathcal{O}^{\rm tw4(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) + {}^I\mathcal{O}^{\rm tw5(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x})\,, \qquad (4.3.27)$$

with

$$\begin{aligned}{}^I\mathcal{O}^{\rm tw4(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) &= \tfrac{1}{2}x_{[\alpha}\partial_{\beta]}\partial^\mu\int_0^1 \mathrm{d}\lambda\,\lambda^I(1-\lambda^2)\,{}^I\mathcal{O}_\mu(\kappa_1\lambda x, \tau\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}\,,\\ {}^I\mathcal{O}^{\rm tw5(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) &= -\tfrac{1}{4}x_{[\alpha}\big(\delta^\mu_{\beta]}(x\partial) - x^\mu\partial_{\beta]}\big)\Box\\ &\qquad\times\int_0^1\mathrm{d}\lambda\,\lambda^I(1-\lambda)^2\,{}^I\mathcal{O}_\mu(\kappa_1\lambda x, \tau\lambda x, \kappa_2\lambda x)\big|_{x=\tilde{x}}\,.\end{aligned}$$

The symmetric part reads

$${}^I\mathcal{O}^{\rm tw3}_{(\alpha\beta)}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\,\lambda^I\partial_{(\alpha}{}^I\mathcal{O}_{\beta)}(\kappa_1\lambda\tilde{x}, \tau\lambda\tilde{x}, \kappa_2\lambda\tilde{x}) - {}^I\mathcal{O}^{>({\rm ii})}_{(\alpha\beta)}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x}) \qquad (4.3.28)$$

where

$$ {}^{I}\mathcal{O}^{>(\mathrm{ii})}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\, \lambda^I \Big\{ \frac{1}{2}(1-\lambda^2) g_{\alpha\beta}\partial^\mu + \frac{1}{2}(1-\lambda)\delta^\mu_{(\alpha} x_{\beta)}\Box \\ + \lambda(1-\lambda) x_{(\alpha}\partial_{\beta)}\partial^\mu - \frac{1}{4}(1-\lambda)^2 x_\alpha x_\beta \partial^\mu \Box \Big\} {}^{I}\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \tag{4.3.29} $$

Let us remark that the conditions of tracelessness for the light-cone operators are:

$$ g^{\alpha\beta}\,{}^{I}\mathcal{O}^{\mathrm{tw3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = 0\,, \qquad \begin{aligned} \mathrm{d}^\alpha\,{}^{I}\mathcal{O}^{\mathrm{tw3}}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) &= 0\,, \\ \mathrm{d}^\alpha\,{}^{I}\mathcal{O}^{\mathrm{tw3}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) &= 0\,. \end{aligned} $$

Now we can calculate the twist–3 vector operator from the twist–3 tensor operator. The twist–3 light–cone vector operator reads

$$ {}^{I}\mathcal{O}^{\mathrm{tw3}}_{\alpha}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\, \lambda^{2+I} \Big[\delta^\mu_\alpha(x\partial) - x^\mu\partial_\alpha - x_\alpha\partial^\mu\Big] {}^{I}\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ = {}^{I}\mathcal{O}_\alpha(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) - \tilde{x}_\alpha \int_0^1 \mathrm{d}\lambda\, \lambda^{2+I}\partial^\mu\, {}^{I}\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}, \tag{4.3.30} $$

and the twist–4 vector operator which is contained in the trace terms of the twist–3 vector operator

$$ {}^{I}\mathcal{O}^{\mathrm{tw4(ii)}}_{\alpha}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \tilde{x}_\alpha \int_0^1 \mathrm{d}\lambda\, \lambda^{2+I}\partial^\mu\, {}^{I}\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}\,. \tag{4.3.31} $$

The determination of the antisymmetric twist–4 operator having symmetry type (iii) obtains from Young tableaux $\begin{array}{|c|c|c|c|}\hline \rho & \mu_1 & \cdots & \mu_N \\ \hline \beta \\ \cline{1-1} \alpha \\ \cline{1-1} \end{array}$ and $\begin{array}{|c|c|c|c|c|}\hline \rho & \sigma & \mu_1 & \cdots & \mu_N \\ \hline \beta \\ \cline{1-1} \alpha \\ \cline{1-1} \end{array}$, respectively. Again, making use of property (4.3.19), the result is

$$ {}^{I}\mathcal{O}^{\mathrm{tw4(iii)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \int_0^1 \mathrm{d}\lambda\lambda^{2+I}\delta^\mu_{[\alpha}\Big((x\partial)\delta^\nu_{\beta]} - 2x^\nu\partial_{\beta]}\Big){}^{I}\mathcal{O}_{[\mu\nu]}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ - {}^{I}\mathcal{O}^{\mathrm{tw5(iii)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}), \tag{4.3.32} $$

twist–5 part is determined by the trace namely with

$$ {}^{I}\mathcal{O}^{\mathrm{tw5(iii)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa\tilde{x}) = -\int_0^1 \mathrm{d}\lambda\, \lambda^I (1-\lambda^2)\Big\{ x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]} \\ - x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box\Big\}{}^{I}\mathcal{O}_{[\mu\nu]}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \tag{4.3.33} $$

The antisymmetric twist–5 tensor operator is given by:

$$ {}^{I}\mathcal{O}^{\mathrm{tw5}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa\tilde{x}) = {}^{I}\mathcal{O}^{\mathrm{tw5(ii)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa\tilde{x}) + {}^{I}\mathcal{O}^{\mathrm{tw5(iii)}}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa\tilde{x}) \\ = -\frac{1}{4}x_{[\alpha}\big(\delta^\mu_{\beta]}(x\partial) - x^\mu\partial_{\beta]}\big)\Box\int_0^1 \mathrm{d}\lambda\, \lambda^I (1-\lambda)^2\, {}^{I}\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ - \int_0^1 \mathrm{d}\lambda\, \lambda^I(1-\lambda^2)\Big\{ x_{[\alpha}\big(\delta^{[\mu}_{\beta]}(x\partial) - x^{[\mu}\partial_{\beta]}\big)\partial^{\nu]} \\ - x_{[\alpha}\delta^{[\mu}_{\beta]}x^{\nu]}\Box\Big\}{}^{I}\mathcal{O}_{[\mu\nu]}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}}. \tag{4.3.34} $$

Finally, we obtain the complete decomposition of the antisymmetric tensor operator:

$$\begin{aligned} {}^I\mathcal{O}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) &= {}^I\mathcal{O}^{\rm tw3(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) + {}^I\mathcal{O}^{\rm tw4(iii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) \\ &\quad + {}^I\mathcal{O}^{\rm tw4(ii)}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) + {}^I\mathcal{O}^{\rm tw5}_{[\alpha\beta]}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}). \end{aligned} \tag{4.3.35}$$

The determination of the symmetric twist–4 operator having symmetry type (iv) obtains from Young tableaux $\begin{array}{|c|c|c|c|c|}\hline \rho & \mu_1 & \mu_2 & \cdots & \mu_N \\ \hline \beta & \alpha \\ \cline{1-2} \end{array}$ and $\begin{array}{|c|c|c|c|c|}\hline \rho & \sigma & \mu_1 & \cdots & \mu_N \\ \hline \beta & \alpha \\ \cline{1-2} \end{array}$, respectively. Making use of property (4.3.19) the result is

$$\begin{aligned} &{}^I\mathcal{O}^{\rm tw4(iv)}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \\ &\quad \int_0^1 \mathrm{d}\lambda\, \lambda^I(1-\lambda)\Big(\delta^\mu_\alpha\delta^\nu_\beta x^\rho(x\partial)\partial_\rho - 2x^\mu(x\partial)\delta^\nu_{(\beta}\partial_{\alpha)}\Big){}^I\mathcal{O}_{(\mu\nu)}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big|_{x=\tilde{x}} \\ &\quad -{}^I\mathcal{O}^{>{\rm (iv)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})\,, \end{aligned} \tag{4.3.36}$$

where

$$\begin{aligned} &{}^I\mathcal{O}^{>{\rm (iv)}}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \\ &\int_0^1 \mathrm{d}\lambda\Big\{\Big(2\lambda^{2+I}x_{(\alpha}\delta^\nu_{\beta)}\partial^\mu - \lambda^{1+I}(1-\lambda)x_\alpha x_\beta\partial^\mu\partial^\nu\Big){}^I\mathcal{O}_{(\mu\nu)}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x) \\ &-\frac{1}{2}\Big(\lambda^I g_{\alpha\beta}\partial^\mu + \lambda^I(1-\lambda^2)x_{(\alpha}\delta^\mu_{\beta)}\Box - \frac{1}{2}\lambda^I(1-\lambda)^2 x_\alpha x_\beta\Box\partial^\mu\Big){}^I\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x) \\ &+\Big(\frac{1}{2}\lambda^I g_{\alpha\beta}(x\partial) - \lambda^{2+I}x_{(\alpha}\partial_{\beta)} + \frac{1}{2}\lambda^{1+I}(1-\lambda)x_\alpha x_\beta\Box\Big){}^I\mathcal{O}^\rho{}_\rho(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}. \end{aligned} \tag{4.3.37}$$

Now, we use the property (4.3.19) and sum up the symmetric *higher twist* operators, which appear in the trace terms of the symmetric twist operators with Young symmetry (ii) and (iv), to a complete twist–4, twist–5 and twist–6 operator. The complete twist–4 operators are constructed by means of the Young tableau (i). The 'scalar part' of the twist–4 tensor operator is given by

$$\begin{aligned} {}^I\mathcal{O}^{\rm tw4,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = \frac{1}{2}g_{\alpha\beta}\Big\{&(x\partial)\int_0^1 \mathrm{d}\lambda\,\lambda^I\,{}^I\mathcal{O}^\rho{}_\rho(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x) \\ &- \partial^\mu\int_0^1 \mathrm{d}\lambda\,\lambda^{2+I}\,{}^I\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}} \end{aligned} \tag{4.3.38}$$

and the twist–4 'vector part' reads

$$\begin{aligned} {}^I\mathcal{O}^{\rm tw4,v}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) &= -\tilde{x}_{(\alpha}\Big\{\int_0^1 \mathrm{d}\lambda\,\lambda^I\,\Big\{\partial_{\beta)} + \frac{1}{2}x_{\beta)}(\ln\lambda)\Box\Big\}{}^I\mathcal{O}^\rho{}_\rho(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x) \\ &- \int_0^1 \mathrm{d}\lambda\,\lambda^I\,\Big\{\big(1-\lambda^2\big)\partial_{\beta)} + \frac{1}{4}x_{\beta)}\big(1-\lambda^2+2\ln\lambda\big)\Box\Big\}\partial^\mu\,{}^I\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}. \end{aligned} \tag{4.3.39}$$

Furthermore, the complete twist–5 vector operator is again constructed by means of the Young tableau (ii). The ‘vector part’ of the twist–5 tensor operator is given by

$$
\begin{aligned}
&{}^I\mathcal{O}^{\rm tw5,v}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) = -\tilde{x}_{(\alpha}\times \qquad\qquad (4.3.40)\\
&\Big\{\frac{1}{4}\int_0^1 \mathrm{d}\lambda\,\lambda^I\Big(\big(1-\lambda\big)^2\big(\delta^\nu_{\beta)}(x\partial)-x^\nu\partial_{\beta)}\big)\Box+2\big(1-\lambda+\ln\lambda\big)x_{\beta)}\partial^\nu\Box\Big)\,{}^I\mathcal{O}_\nu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\\
&-\int_0^1 \mathrm{d}\lambda\,\lambda^I\Big\{\big(1-\lambda^2\big)\big(\delta^\nu_{\beta)}(x\partial)-x^\nu\partial_{\beta)}-x_{\beta)}\partial^\nu\big)\partial^\mu-\Big(\frac{1}{2}\big(1-\lambda^2\big)+\ln\lambda\Big)x_{\beta)}\Box\big(\partial^\mu x^\nu-g^{\mu\nu}\big)\Big\}\\
&\qquad\qquad\qquad\qquad\times\,{}^I\mathcal{O}_{(\mu\nu)}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}.
\end{aligned}
$$

The full twist–6 scalar operator has Young symmetry (i). The ‘scalar part’ of the twist–6 tensor operator is given by

$$
\begin{aligned}
&{}^I\mathcal{O}^{\rm tw6,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})\\
&\quad=\tilde{x}_\alpha\tilde{x}_\beta\Big\{\frac{1}{2}\Box\int_0^1 \mathrm{d}\lambda\,\lambda^I\big(1-\lambda+\ln\lambda\big)\big(\partial^\mu\,{}^I\mathcal{O}_\mu(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)-{}^I\mathcal{O}^\rho{}_\rho(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\big)\\
&\qquad\qquad+\partial^\mu\partial^\nu\int_0^1 \mathrm{d}\lambda\,\lambda^I(1-\lambda)\,{}^I\mathcal{O}_{(\mu\nu)}(\kappa_1\lambda x,\tau\lambda x,\kappa_2\lambda x)\Big\}\Big|_{x=\tilde{x}}. \qquad (4.3.41)
\end{aligned}
$$

Thus, together with the twist–3 part, Eq. (4.3.28), and the twist–4 part, Eq. (4.3.36), we finally obtain the complete decomposition of the symmetric tensor operator:

$$
\begin{aligned}
{}^I\mathcal{O}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}) &= {}^I\mathcal{O}^{\rm tw3(ii)}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})+{}^I\mathcal{O}^{\rm tw4(iv)}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})\\
&+{}^I\mathcal{O}^{\rm tw4,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})+{}^I\mathcal{O}^{\rm tw4,v}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})+{}^I\mathcal{O}^{\rm tw5,v}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x})\\
&+{}^I\mathcal{O}^{\rm tw6,s}_{(\alpha\beta)}(\kappa_1\tilde{x},\tau\tilde{x},\kappa_2\tilde{x}). \qquad (4.3.42)
\end{aligned}
$$

4.3.3 Conclusions

In this Section we have discussed the twist decomposition of trilocal light–ray operators. The only difference is that the respective values of twist increase by a definite amount which is equal for any given operator and possibly, depending on the number I of external x–factors, by a change in the integration measure, $\mathrm{d}\lambda \to \mathrm{d}\lambda\ \lambda^I$. The applicability of our procedure for bilocal operators rests upon the fact that the general form of the Taylor expansion (around $y=0$) for any of the nonlocal operators is the same: It is given by an infinite sum of terms of Nth order being multiplied by $x^{\mu_1}x^{\mu_2}\dots x^{\mu_N}$ where each term consists of a finite sum of local operators having exactly the same tensor structure. In principle, this works also for multi-local operators of higher order.

However, there are special trilocal operators which do not immediately fit into the general scheme of Section 4.2. Its twist decomposition is obtained by restricting the expressions of the generic case to the special conditions under consideration. The results are classified in Table 4.3 and 4.4, again by indicating their symmetry type and the number of the corresponding equation. In the same manner one should proceed for other kinds of special multilocal operators like that of Eq. (4.3.8).

The various tensor operators which we considered here are nonlocal generalizations of local operators partly considered earlier in the literature for twist–3 and twist–4. There, as far as possible an explicit twist decomposition has been circumvented by truncating *any* of the

Eq.	YT	type	${}^{I}\mathcal{O}_{\alpha\beta}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x})$
4.3.28	(ii)	sym	$\tau = 3$
4.3.25	(ii)	asym	$\tau = 3$
4.3.32	(iii)	asym	$\tau = 4$
4.3.36	(iv)	sym	$\tau = 4$
4.3.38	(i)	sym	$\tau = 4$
4.3.39	(i)	sym	$\tau = 4$
4.3.28	(i)	asym	$\tau = 4$
4.3.40	(ii)	sym	$\tau = 5$
4.3.34	(ii)	asym	$\tau = 5$
4.3.41	(i)	sym	$\tau = 6$

Table 4.3: Twist decomposition of special tensor operators ${}^{I}\mathcal{O}_{\alpha\beta}$ with the property $x^{\beta}\, {}^{I}\mathcal{O}_{\alpha\beta} \equiv 0$

Eq.	YT	${}^{I}\mathcal{O}_{\alpha}(\kappa_1\tilde{x}, \tau\tilde{x}, \kappa_2\tilde{x})$
4.3.30	(ii)	$\tau = 3$
4.3.31	(i)	$\tau = 4$

Table 4.4: Twist decomposition of special vector operators ${}^{I}\mathcal{O}_{\alpha}$ with the property $x^{\alpha}\, {}^{I}\mathcal{O}_{\alpha} \equiv 0$

(constant) totally symmetrized tensor indices of the local operators by some light–like vector n^{μ}, i.e., by reducing to the scalar case. To the best of our knowledge the only work where the twist decomposition of (scalar) bilocal operators has been considered is that of Balitsky and Braun [9]. However, their work is based on the external field formalism and does not have an obvious group theoretical systematics.

In principle, our procedure may be extended also to tensors of arbitrary high rank. Despite being defined by an obvious algorithm, its application to the next step, the twist decomposition of a generic 3rd rank tensor operator, will be very cumbersome. First of all, it would be necessary to determine the projection operators onto all traceless 3rd rank tensorial harmonic polynomials and, secondly, also additional Young patterns had to be taken into account. Fortunately, such kind of nonlocal operators – at least in the near future – may not be of physical relevance. Therefore, any further study in that direction seems to be reasonable only from a group theoretical point of view.

4.4 Harmonic operators of definite geometric twist in terms of Gegenbauer polynomials and Bessel functions

4.4.1 Totally symmetric harmonic operators of any geometric twist

Let us now discuss the twist decomposition of a scalar operator of degree n into all its twist parts by means of the polynomial technique. In this way, the scalar operators of degree n are the generating polynomials of symmetric tensors of rank n. In this case, the infinite twist decomposition of non-local (scalar) operators into the infinite tower of harmonic operators of definite twist is possible. The corresponding group theoretical background has been formulated and proven by Bargmann and Todorov [19]. In order to demonstrate the efficiency of the polynomial technique and because of their physical relevance we apply it to the complete twist decomposition of the scalar operators $N(x,-x)$, $M(x,-x)$ and $O(x,-x)$ having minimal twist $\tau=3$ and $\tau=2$, respectively, as well as to the related vector and tensor operators.

Scalar harmonic operators without external operations

To begin with the simplest case let us study the local operators $N_n(x)$ of degree n which are generating polynomials of symmetric tensors of degree n. The general formula for the decomposition of the local scalar operator $N_n(x)$ which can be considered as a homogeneous polynomial of degree n into its harmonic polynomials of definite twist $\tau=3+2j$, $j=0,1,\ldots,$ is given by [19]

$$N_n(x)=\sum_{j=0}^{[\frac{n}{2}]}\frac{(n+1-2j)!}{4^j j!(n+1-j)!}\,x^{2j}\,N^{\mathrm{tw}(3+2j)}_{n-2j}(x), \tag{4.4.1}$$

where the harmonic operators of definite twist are defined by (see also Ref. [121], Eq. 9.3.2(3))

$$N^{\mathrm{tw}(3+2j)}_{n-2j}(x)=\sum_{k=0}^{[\frac{n-2j}{2}]}\frac{(-1)^k(n-2j-k)!}{4^k k!(n-2j)!}\,x^{2k}\Box^k\left(\Box^j N_n(x)\right). \tag{4.4.2}$$

They satisfy the condition of tracelessness

$$\Box N^{\mathrm{tw}(3+2j)}_{n-2j}(x)=0.$$

Therefore, the polynomials of Eq. (4.4.2) span the space of homogeneous harmonic polynomials of degree $n-2j$.

After the substitution $N_n(x)=\int \mathrm{d}^4p\,N(p)(xp)^n$ into (4.4.2), and using

$$\Box^{k+j}\,(xp)^n=\frac{n!}{(n-2k-2j)!}(p^2)^{k+j}(xp)^{n-2k-2j},$$

we rewrite these harmonic operators in terms of Gegenbauer polynomials as follows:[10]

$$N^{\mathrm{tw}(3+2j)}_{n-2j}(x)=\frac{n!}{(n-2j)!}\int \mathrm{d}^4p\,N(p)\,p^{2j}\left(\frac{1}{2}\sqrt{p^2x^2}\right)^{n-2j}C^1_{n-2j}\left(\frac{px}{\sqrt{p^2x^2}}\right). \tag{4.4.3}$$

From the group theoretical point of view, Eq. (4.4.3) is another form of the harmonic extension (4.4.2). Here, the $p-$integration introduces a superposition of the Fourier transforms $N(p)$ times p^{2j} with coefficients being (two-sided) harmonic polynomials,

$$h^1_{n-2j}(p|x)=\left(\frac{1}{2}\sqrt{p^2x^2}\right)^{n-2j}C^1_{n-2j}\big(px/\sqrt{p^2x^2}\big) \tag{4.4.4}$$

with

$$\Box_x\,h^1_{n-2j}(p|x)=0=\Box_p\,h^1_{n-2j}(p|x),\qquad h^1_{n-2j}(p|x)=h^1_{n-2j}(x|p). \tag{4.4.5}$$

The resummation of the local expressions $N_n(x)$ according to $N(x,-x)=\sum_{n=0}^{\infty}(\mathrm{i}^n/n!)N_n(x)$ is obtained by using the well-known integral representation of Euler's beta function,

$$B(n,m)=\frac{\Gamma(n)\Gamma(m)}{\Gamma(n+m)}=\int_0^1\mathrm{d}t\,t^{n-1}(1-t)^{m-1}. \tag{4.4.6}$$

---

[10] Let me note that the Chebyshev polynomials of the second kind, $U_n(z)$, used in Refs. [16, 23] are defined in terms of Gegenbauer polynomials as follows: $U_n(z)=C^1_n(z)$ (see [121]).

Then the (infinite) twist decomposition of the nonlocal scalar operator $N(x,-x)$ into nonlocal harmonic operators $N^{\mathrm{tw}(3+2j)}(x,-x)$ of twist $\tau = 3+2j$, $j = 0,1,2,\ldots$, reads

$$N(x,-x) = N^{\mathrm{tw3}}(x,-x) + \sum_{j=1}^{\infty} \frac{x^{2j}}{4^j j!(j-1)!} \int_0^1 \mathrm{d}t\, t\,(1-t)^{j-1}\, N^{\mathrm{tw}(3+2j)}(tx,-tx), \qquad (4.4.7)$$

with

$$\begin{aligned} N^{\mathrm{tw}(3+2j)}(x,-x) = \sqrt{\pi} \int \mathrm{d}^4 p\, N(p)\, (-p^2)^j\, (1 + p\partial_p) \left(\sqrt{(px)^2 - p^2x^2}\right)^{-1/2} & \\ \times J_{1/2}\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right) \mathrm{e}^{\mathrm{i}px/2}. & \qquad (4.4.8) \end{aligned}$$

Let us remark that, for notational simplicity, in Eq. (4.4.7) we wrote the nonlocal harmonic operators for $j \geq 1$ without including the integration over t which is due to the normalization of the local harmonic operators, cf. Eqs. (4.4.1) and (4.4.3). The normalization of the nonlocal harmonic operators may be read off from the terms in front of the $t-$integral in Eq. (4.4.7). Let us point also to the remarkable fact that the property of harmonicity is independent of j; this obtains immediately from the fact that the following expressions are (two-sided) harmonic functions,

$$\mathcal{H}_1(p|x) = \sqrt{\pi} \left(\sqrt{(px)^2 - p^2x^2}\right)^{-1/2} J_{1/2}\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right) \mathrm{e}^{\mathrm{i}px/2} \qquad (4.4.9)$$

with

$$\Box_x\, \mathcal{H}_1(p|x) = 0 = \Box_p\, \mathcal{H}_1(p|x), \qquad \mathcal{H}_1(p|x) = \mathcal{H}_1(x|p). \qquad (4.4.10)$$

Furthermore, we observe that the factor $(1 + p\partial/\partial p)$ in Eq. (4.4.8) may be changed into $(1 + x\partial/\partial x)$ after which it can be taken outside the $p-$integration and, in Eq. (4.4.7), could be changed into $(1 + t\partial/\partial t)$; if desired, this could be used for a partial integration.

The resulting expression (4.4.8) can be rewritten in a form which, in the case $j = 0$, has been already introduced by [11, 15]. Namely, using

$$z^{-n-1/2} J_{n+1/2}(z) = (-1)^n \sqrt{\frac{2}{\pi}} \Big(\frac{1}{z}\frac{\mathrm{d}}{\mathrm{d}z}\Big)^n \Big(\frac{\sin z}{z}\Big), \qquad (4.4.11)$$

we obtain

$$\begin{aligned} N^{\mathrm{tw}(3+2j)}(x,-x) &= \int \mathrm{d}^4p\, N(p)\, (-p^2)^j\, \frac{2}{\sqrt{(px)^2 - p^2x^2}}\, p\partial_p \left(\sin\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right) \mathrm{e}^{\mathrm{i}px/2}\right) \\ &= \int \mathrm{d}^4p\, N(p)\, (-p^2)^j\, \frac{1}{2\sqrt{(px)^2 - p^2x^2}} \bigg(\left(px + \sqrt{(px)^2 - p^2x^2}\right) \mathrm{e}^{(\mathrm{i}/2)\left(px + \sqrt{(px)^2 - p^2x^2}\right)} \\ &\qquad - \left(px - \sqrt{(px)^2 - p^2x^2}\right) \mathrm{e}^{(\mathrm{i}/2)\left(px - \sqrt{(px)^2 - p^2x^2}\right)} \bigg) \qquad (4.4.12) \\ &= \int \mathrm{d}^4p\, N(p)\, (-p^2)^j \Big\{ \cos\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right) + \frac{\mathrm{i}(px)}{\sqrt{(px)^2 - p^2x^2}} \sin\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right)\Big\} \mathrm{e}^{\mathrm{i}px/2}. \end{aligned}$$

If we use the abbreviation, which was introduced by Balitsky and Braun (see Eq. (5.15) in Ref. [11]),

$$\left[\mathrm{e}^{\mathrm{i}px}\right]_{\mathrm{lt}} = \Big\{ \cos\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right) + \frac{\mathrm{i}(px)}{\sqrt{(px)^2 - p^2x^2}} \sin\left(\frac{1}{2}\sqrt{(px)^2 - p^2x^2}\right)\Big\} \mathrm{e}^{\mathrm{i}px/2}, \quad (4.4.13)$$

we may rewrite Eq. (4.4.12) according to

$$N^{\mathrm{tw}(3+2j)}(x,-x)=\int \mathrm{d}^4p\, N(p)\,(-p^2)^j\ \left[\mathrm{e}^{\mathrm{i}px}\right]_{\mathrm{lt}}\,. \tag{4.4.14}$$

Analogous results may be obtained also in the more complicated cases which will be considered below.

Scalar harmonic operators with external operations

Now, let us consider the case of the operators $O(x,-x)=x^\alpha O_\alpha(x,-x)$, where an additional power of x occurs through contraction of the vector operator with x^α. Therefore, the formulas (4.4.1) and (4.4.2) are to be written down for the scalar operator $O_{n+1}(x)=\int \mathrm{d}^4p\, x_\mu O^\mu(p)(xp)^n$, i.e., by replacing n by $n+1$ within these expressions for the decomposition into harmonic operators of any twist $\tau=2+2j$. Then the decomposition of the scalar local operators $O_{n+1}(x)$ reads:

$$O_{n+1}(x)=\sum_{j=0}^{[\frac{n+1}{2}]}\frac{(n+2-2j)!}{4^j j!(n+2-j)!}\,x^{2j}\,O^{\mathrm{tw}(2+2j)}_{n+1-2j}(x), \tag{4.4.15}$$

where the local harmonic operators of twist $\tau=2+2j$ are given by

$$O^{\mathrm{tw}(2+2j)}_{n+1-2j}(x)=\sum_{k=0}^{[\frac{n+1-2j}{2}]}\frac{(-1)^k(n+1-2j-k)!}{4^k k!(n+1-2j)!}\,x^{2k}\Box^k\left(\Box^j O_{n+1}(x)\right). \tag{4.4.16}$$

After performing the differentiations $\Box^{k+j}O_{n+1}(x)$ we can rewrite these harmonic operators in terms of Gegenbauer polynomials as follows:

$$\begin{aligned} O^{\mathrm{tw}(2+2j)}_{n+1-2j}(x)=\frac{n!}{(n+1-2j)!}\int \mathrm{d}^4p\, O^\mu(p)\,p^{2j}\Bigg\{x_\mu\left(\frac{1}{2}\sqrt{p^2x^2}\right)^{n-2j}C^2_{n-2j}\!\left(\frac{px}{\sqrt{p^2x^2}}\right) \qquad (4.4.17)\\ -\frac{1}{2}p_\mu x^2\left(\frac{1}{2}\sqrt{p^2x^2}\right)^{n-1-2j}C^2_{n-1-2j}\!\left(\frac{px}{\sqrt{p^2x^2}}\right)+2j\,\frac{p_\mu}{p^2}\left(\frac{1}{2}\sqrt{p^2x^2}\right)^{n+1-2j}C^1_{n+1-2j}\!\left(\frac{px}{\sqrt{p^2x^2}}\right)\Bigg\}, \end{aligned}$$

where the additional terms proportional to p_μ occur because of the appearance of derivatives with respect to Gamma structure $(x\gamma)$. The change in the order of the Gegenbauer polynomials and, consequently, in the order of the Bessel functions and the accompanying factors $\sqrt{(px)^2-p^2x^2}$ occurring below, is also due to that fact. Of course, for $j=0$ we re-obtain the expression (3.3.14)

Resumming with respect to n ($\geq 2j\pm 1$ or $2j$, respectively) and, again, using the representation (4.4.6) of Euler's beta function we obtain the following (infinite) twist decomposition

$$O(x,-x)=O^{\mathrm{tw}2}(x,-x)+\sum_{j=1}^{\infty}\frac{x^{2j}}{4^j j!(j-1)!}\int_0^1 \mathrm{d}t\,t\,(1-t)^{j-1}\,O^{\mathrm{tw}(2+2j)}(tx,-tx), \tag{4.4.18}$$

with

$$\begin{aligned} O^{\mathrm{tw}(2+2j)}(x,-x)=\sqrt{\pi}\int \mathrm{d}^4p\, O^\mu(p)\,(-p^2)^j\Bigg\{\left(x_\mu(2+p\partial_p)-\frac{1}{2}\,\mathrm{i}p_\mu x^2\right) \qquad (4.4.19)\\ \times(3+p\partial_p)\left(\sqrt{(px)^2-p^2x^2}\right)^{-3/2}J_{3/2}\left(\frac{1}{2}\sqrt{(px)^2-p^2x^2}\right)\\ -2j\,\frac{\mathrm{i}p_\mu}{p^2}\,(1+p\partial_p)\left(\sqrt{(px)^2-p^2x^2}\right)^{-1/2}J_{1/2}\left(\frac{1}{2}\sqrt{(px)^2-p^2x^2}\right)\Bigg\}\,\mathrm{e}^{\mathrm{i}px/2}. \end{aligned}$$

Again, by convention, the resummed nonlocal harmonic operators $O^{\mathrm{tw}(2+2j)}(tx,-tx)$ of twist $\tau = 2+2j$, $j = 0, 1, 2, \cdots$, are obtained by changing any $x \to tx$. Also here analogous comments are in order as for the simpler case of the operator $N(x,-x)$. Harmonicity of the nonlocal operators (4.4.19), $\Box O^{\mathrm{tw}(2+2j)}(x,-x) = 0$, holds by construction. Obviously, for $j = 0$ we recover the expression (3.3.16) .

Finally, we consider the scalar operator $M(x,-x) = x^\nu \partial^\mu M_{[\mu\nu]}(x,-x)$ resulting from the skew tensor operator $M_{[\mu\nu]}(x,-x)$. This operator is governed by totally symmetric tensors since the skew tensors $M_{[\mu\nu]n}$ having symmetry type $[n+2] = (n+1,1)$ are composed with another skew tensor $x^{[\nu}\partial^{\mu]}$ having symmetry type $[2] = (1,1)$ and the Clebsch–Gordan series of their direct product consists only of one symmetry type, $[n] = (n)$.

The local operators are decomposed according to formulas (4.4.1) and (4.4.2). However, because of the special 'external' structure the resulting expressions simplify considerably. Namely, after performing the differentiations

$$\Box^{k+j} \partial^\mu \sigma_{\mu\nu} x^\nu (xp)^n = \frac{n!}{(n-1-2k-2j)!} \big(p^\mu \sigma_{\mu\nu} x^\nu\big) p^{2j} (p^2)^k (px)^{n-1-2k-2j},$$

we arrive at the following harmonic operators in terms of Gegenbauer polynomials:

$$M^{\mathrm{tw}(3+2j)}_{n-2j}(x) = \frac{n!}{(n-2j)!} \int \mathrm{d}^4 p \, M_{[\mu\nu]}(p) \, p^\mu x^\nu \, p^{2j} \left(\frac{1}{2}\sqrt{p^2 x^2}\right)^{n-1-2j} C^2_{n-1-2j}\left(\frac{px}{\sqrt{p^2 x^2}}\right). \tag{4.4.20}$$

Now, resumming with respect to $n \,(\geq 2j+1)$ we finally get the infinite twist decomposition:

$$M(x,-x) = M^{\mathrm{tw}3}(x,-x) + \sum_{j=1}^{\infty} \frac{x^{2j}}{4^j j!(j-1)!} \int_0^1 \mathrm{d}t \, t \, (1-t)^{j-1} \, M^{\mathrm{tw}(3+2j)}(tx,-tx), \tag{4.4.21}$$

with

$$\begin{aligned} M^{\mathrm{tw}(3+2j)}(x,-x) &= \sqrt{\pi} \int \mathrm{d}^4 p \, M_{[\mu\nu]}(p) \, (-p^2)^j \, \mathrm{i} p^\mu x^\nu \\ &\quad \times (2 + p\partial_p)\,(3 + p\partial_p) \left(\sqrt{(px)^2 - p^2 x^2}\right)^{-3/2} J_{3/2}\left(\frac{1}{2}\sqrt{(px)^2 - p^2 x^2}\right) \mathrm{e}^{\mathrm{i}px/2}. \end{aligned} \tag{4.4.22}$$

Again, harmonicity of the nonlocal operators $M^{\mathrm{tw}(3+2j)}(x,-x)$ of twist $\tau = 3+2j$, $j = 0, 1, 2, \cdots$, is fulfilled without taking care of j.

Harmonic vector and tensor operators related to totally symmetric local operators

The prescribed procedure may be also used for the twist decomposition of any totally symmetric (tensor) operator with an arbitrary number of free tensor indices which, in the framework of the polynomial technique, are recovered by straightforward differentiation with respect to x of the corresponding scalar operators.

The simplest of these operators, whose tower of infinite twist part starts with $\tau = 2$ and which contains the leading contributions to virtual Compton scattering is the operator $O^{\mathsf{S}}_\alpha(x,-x)$. Its symmetry type S is characterized by the Young frames $(n+1)$, $1 \leq n \leq \infty$ (cf. also Ref. [63] where we denoted that symmetry type by (i)). The local expressions, $O^{\mathsf{S}}_{\alpha n}(x)$, are obtained from the local scalar operators (4.4.15) of degree $n+1$ by applying $(1/(n+1))\partial_\alpha$. Eventually, we get

$$\begin{aligned}\partial_\alpha O_{n+1}(x) = &\sum_{j-1=0}^{[\frac{n+1}{2}]} \frac{(n+2-2j)!}{4^j (j-1)!(n+2-j)!}\, 2x_\alpha\, x^{2(j-1)}\, O^{\mathrm{tw}(2+2j)}_{n+1-2j}(x) \\ &+ \sum_{j=0}^{[\frac{n+1}{2}]} \frac{(n+2-2j)!}{4^j j!(n+2-j)!}\, x^{2j} \left(\partial_\alpha O^{\mathrm{tw}(2+2j)}_{n+1-2j}(x)\right).\end{aligned}$$

and the representation (4.4.6) of the beta function, both sums may be rewritten as

$$\sum_{j=0}^{[\frac{n+1}{2}]} \frac{(n-2j)!}{4^j j!(n+1-j)!}\, x^{2j}\, O^{\mathrm{tw}(4+2j)}_{n-1-2j}(x) = \sum_{j=0}^{[\frac{n-1}{2}]} \frac{x^{2j}}{4^j (j!)^2} \int_0^1 \mathrm{d}t\, t\, (1-t)^j\, O^{\mathrm{tw}(4+2j)}_{n-1-2j}(tx),$$

$$\begin{aligned}\sum_{j=0}^{[\frac{n+1}{2}]} \frac{(n+2-2j)!}{4^j j!(n+2-j)!}\, x^{2j} \partial_\alpha O^{\mathrm{tw}(2+2j)}_{n+1-2j}(x) &= \partial_\alpha O^{\mathrm{tw}\ 2}_{n+1}(x) \\ &+ \sum_{j=1}^{[\frac{n+1}{2}]} \frac{x^{2j}}{4^j j!(j-1)!} \int_0^1 \mathrm{d}t\, t\, (1-t)^{j-1} \left(\partial_\alpha O^{\mathrm{tw}(2+2j)}_{n+1-2j}(tx)\right),\end{aligned}$$

respectively, where $O^{\mathrm{tw}(4+2j)}_{n-1-2j}(x)$ is given by Eq. (4.4.17) with $j \to j+1$. From this it becomes obvious that, beginning with twist-4, one obtains two different contributions of the same twist, namely a vector and a scalar part.

Now, putting together these terms and resumming over n, thereby representing the normalizing coefficient $1/(n+1)$ as $\int_0^1 \mathrm{d}\lambda\, \lambda^n$, we obtain

$$\begin{aligned}O^{\mathsf{S}}_\alpha(x,-x) = \int_0^1 \mathrm{d}\lambda \Big\{ &\left(\partial_\alpha O^{\mathrm{tw}2}\right)(\lambda x, -\lambda x) \\ &+ \sum_{j=1}^{\infty} \frac{x^{2j}\lambda^{2j}}{4^j j!(j-1)!} \int_0^1 \mathrm{d}t\, t\, (1-t)^{j-1} \left(\partial_\alpha O^{\mathrm{tw}(2+2j)}\right)(\lambda t x, -\lambda t x) \\ &- \frac{1}{2} x_\alpha \sum_{j=0}^{\infty} \frac{x^{2j}\lambda^{2j+1}}{4^j (j!)^2} \int_0^1 \mathrm{d}t\, t\, (1-t)^j\, O^{\mathrm{tw}(4+2j)}(\lambda t x, -\lambda t x) \Big\}, \end{aligned} \tag{4.4.23}$$

where $O^{\mathrm{tw}(4+2j)}(x,-x) \equiv O^{\mathrm{tw}(2+2(j+1))}(x,-x)$ and $O^{\mathrm{tw}(2+2j)}(x,-x)$ are given by Eq. (4.4.19). Both the integrals over λ and t are due to normalizations and, therefore, it depends on the personal taste if they are to be included into the definition of the harmonic operators of twist τ or if they are considered as part of the twist decomposition. It is interesting to note that in Eq. (4.4.23) only the twist-2 vector and the twist-4 scalar operator survives on the light-cone. All other higher twist operators for $j \geq 1$ are cancelled due to the factor x^2.

In the same manner we are able to define the twist decomposition of (completely symmetric) tensor operators which result from the expression

$$O^{\mathsf{S}}_{(\alpha_1\alpha_2\cdots\alpha_r)n}(x) = \frac{\partial_{\alpha_1}}{(n+1)} \cdots \frac{\partial_{\alpha_r}}{(n+r)} \sum_{j=0}^{[\frac{n+r}{2}]} \frac{(n+1+r-2j)!}{4^j j!(n+1+r-j)!}\, x^{2j}\, O^{\mathrm{tw}(2+2j)}_{n+r-2j}(x)\ .$$

The general procedure is obvious from the consideration of the vector operator. Completely analogous to that case with only one derivative we would obtain a finite number of different operators of a given twist τ.

4.4.2 Harmonic quark-antiquark operators of geometric twist 2 and 3

In this Section we determine the harmonic quark-antiquark operators up to twist $\tau = 2$ and 3 which are relevant for the generalization of the parton distributions of definite (geometric) twist [66] to the double distributions appearing in the parametrization in the non-forward matrix elements of the corresponding non-local operators off the light-cone. According to the framework of polynomial technique, we start with the local operators.

Local harmonic operators

Now we discuss the local harmonic operators of geometric twist-2 and twist-3 for both the chiral-even vector operators,

$$O_{\alpha n}(x) = \bar{\psi}(0)\gamma_\alpha(\mathrm{i}x\overset{\leftrightarrow}{D})^n\psi(0), \tag{4.4.24}$$

$$O_{5\alpha n}(x) = \bar{\psi}(0)\gamma_\alpha\gamma_5(\mathrm{i}x\overset{\leftrightarrow}{D})^n\psi(0), \tag{4.4.25}$$

and the chiral-odd scalar and skew tensor operators,

$$N_n(x) = \bar{\psi}(0)(\mathrm{i}x\overset{\leftrightarrow}{D})^n\psi(0), \tag{4.4.26}$$

$$M_{[\alpha\beta]n}(x) = \bar{\psi}(0)\sigma_{\alpha\beta}(\mathrm{i}x\overset{\leftrightarrow}{D})^n\psi(0), \tag{4.4.27}$$

together with their related scalar and vector operators

$$O_{n+1}(x) = x^\alpha O_{\alpha n}(x), \qquad M_{\mu n+1}(x) = x^\nu M_{[\mu\nu]n}(x), \qquad M_n(x) = \partial^\mu M_{\mu n+1}(x). \tag{4.4.28}$$

Now we consider the decomposition the vector operators $O_{\alpha n}(x)$ into its harmonic twist-2 and twist-3 part. The irreducible vector operators $O^{\mathrm{tw2}}_{\alpha n}(x)$ of Lorentz type $(\frac{n+1}{2}, \frac{n+1}{2})$ have symmetry types $[n+1] = (n+1)$ and can be taken over from the previous Section. On the other hand, the twist-3 vector operators transform according to $(\frac{n+1}{2}, \frac{n-1}{2}) \oplus (\frac{n-1}{2}, \frac{n+1}{2})$ and have symmetry types $[n+1] = (n, 1)$. They may be summed up to the lowest term of another infinite tower of non-local operators of twist $\tau = 3, 4, \ldots$. The conditions of tracelessness for harmonic vector operators are as follows:

$$\Box O^{\mathrm{tw}\,\tau}_{\alpha n}(x) = 0, \qquad \partial^\alpha O^{\mathrm{tw}\,\tau}_{\alpha n}(x) = 0 \qquad \text{for} \qquad \tau = 2, 3. \tag{4.4.29}$$

Eventually, the scalar twist-2 operator is given by (see Eqs. (4.1.7) and (4.1.8) with $h = 2$)

$$O^{\mathrm{tw2}}_{n+1}(x) = \sum_{k=0}^{[\frac{n+1}{2}]} \frac{(-1)^k(n+1-k)!}{4^k k!(n+1)!}\, x^{2k}\, \Box^k O_{n+1}(x).$$

by means of a partial differentiation, ∂_α, together with the normalization factor $1/(n+1)$ as follows (see Eq. (4.1.10)):

$$\begin{aligned} O^{\mathrm{tw2}}_{\alpha n}(x) &= \frac{1}{n+1}\,\partial_\alpha O^{\mathrm{tw2}}_{n+1}(x) \\ &= \frac{1}{n+1}\sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k(n-k)!}{4^k k!(n+1)!}\, x^{2k}\, \mathcal{D}_\alpha(k)\, \Box^k O_{n+1}(x). \end{aligned} \tag{4.4.30}$$

Here, we introduced the operation

$$\mathcal{D}_\alpha(k) = (k + 1 + x\partial)\partial_\alpha - \tfrac{1}{2}\, x_\alpha \Box \tag{4.4.31}$$

which is a generalization of the inner derivative off the light-cone and its extension to arbitrary values of $k \geq 0$ (see Section 4.1).

The twist-3 vector operator which satisfies the conditions (4.4.29) is[11] (see Eq. (4.1.26) with $h = 2$)

$$\begin{aligned} O^{\rm tw3}_{\alpha n}(x) &= \frac{2}{n+1}\, x^\beta \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]} - \frac{1}{n+1}\, x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (n-k)!}{4^k k! n!}\, x^{2k}\, \Box^k O_{\mu n}(x), \qquad (4.4.32)\\ &= \frac{2}{n+1} \sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k (n-1-k)!}{4^k k! n!}\, x^{2k} x^\beta \Big\{ \delta^\mu_{[\alpha}\mathcal{D}_{\beta]}(k) - \frac{1}{n+1}\, x_{[\alpha}\partial_{\beta]}\mathcal{D}^\mu(k) \Big\} \Box^k O_{\mu n}(x). \end{aligned}$$

The proper scalar twist-3 operator is of Lorentz type $(\frac{n}{2}, \frac{n}{2})$ and reads

$$N^{\rm tw3}_n(x) = \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (n-k)!}{4^k k! n!}\, x^{2k} \Box^k N_n(x). \tag{4.4.33}$$

Here I give the decomposition of the skew tensor operator $M_{[\alpha\beta]n}(x)$ into twist-2 and twist-3. The twist-2 tensor operator $M^{\rm tw2}_{[\alpha\beta]n}(x)$ transforms according to $(\frac{n+2}{2}, \frac{n}{2}) \oplus (\frac{n}{2}, \frac{n+2}{2})$, whereas the twist-3 operator which is obtained from the trace terms of $M^{\rm tw2}_{[\alpha\beta]n}(x)$ is of Lorentz type $(\frac{n}{2}, \frac{n}{2})$. They have symmetry types $[n+2] = (n+1, 1)$ and $[n] = (n)$, respectively, and are given by (see Eq. (4.1.26))

$$\begin{aligned} M^{\rm tw2}_{[\alpha\beta]n}(x) &= \frac{2}{n+2} \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]} - \frac{1}{n+2}\, x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \sum_{k=0}^{[\frac{n+1}{2}]} \frac{(-1)^k (n+1-k)!}{4^k k! (n+1)!}\, x^{2k}\, \Box^k M_{\mu n+1}(x)\\ &= \frac{2}{n+2} \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (n-k)!}{4^k k! (n+1)!}\, x^{2k} \Big\{ \delta^\mu_{[\alpha}\mathcal{D}_{\beta]}(k) - \frac{1}{n+2}\, x_{[\alpha}\partial_{\beta]}\mathcal{D}^\mu(k) \Big\} \Box^k M_{\mu n+1}(x), \end{aligned} \tag{4.4.34}$$

and

$$M^{\rm tw3}_{[\alpha\beta]n}(x) = \frac{2}{(n+2)}\, x_{[\alpha} \widehat{M}^{\rm tw3}_{\beta]n-1}(x), \tag{4.4.35}$$

with

$$\begin{aligned} \widehat{M}^{\rm tw3}_{\beta n-1}(x) &= \frac{1}{n}\, \partial_\beta \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (n-k)!}{4^k k! n!}\, x^{2k} \Box^k M_n(x)\\ &= \frac{1}{n} \sum_{k=0}^{[\frac{n-1}{2}]} \frac{(-1)^k (n-1-k)!}{4^k k! n!}\, x^{2k}\, \mathcal{D}_\beta(k)\, \Box^k M_n(x) \end{aligned} \tag{4.4.36}$$

---

[11] We use the convention $A_{[\mu\nu]} \equiv \frac{1}{2}\,(A_{\mu\nu} - A_{\nu\mu})$.

Besides the twist-3 operator (4.4.35) resulting from the trace terms of the twist-2 operator (4.4.34) there exists also a genuine twist-3 operator having Lorentz type $\left(\frac{n}{2}, \frac{n}{2}\right)$ and being governed by the symmetry type $[n+2] = (n,1,1)$. It is given by the following expression[12] (see Eq. (4.1.41) with $h=2$)

$$\begin{aligned}\widetilde{M}^{\mathrm{tw3}}_{[\alpha\beta]n}(x) &= \frac{3}{n+2}\, x^\gamma \left\{ \delta^\mu_{[\alpha}\delta^\nu_\beta \partial_{\gamma]} - \frac{2}{n}\, \delta^\mu_{[\alpha} x_\beta \partial_{\gamma]} \partial^\nu \right\} \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (n-k)!}{4^k k! n!}\, x^{2k}\, \Box^k M_{[\mu\nu]n}(x) \qquad (4.4.37)\\ &= \frac{3}{n+2} \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (n-1-k)!}{4^k k! n!}\, x^{2k}\, x^\gamma \left\{ \delta^\mu_{[\alpha}\delta^\nu_\beta \mathcal{D}_{\gamma]}(k) - \frac{2}{n}\, \delta^\mu_{[\alpha} x_\beta \partial_{\gamma]} \mathcal{D}^\nu(k) \right\} \Box^k M_{[\mu\nu]n}(x).\end{aligned}$$

By construction, Eqs. (4.4.34), (4.4.36) and (4.4.37) are (traceless) harmonic polynomials obeying

$$\Box M^{\mathrm{tw}\,\tau}_{[\alpha\beta]n}(x) = 0, \qquad \partial^\alpha M^{\mathrm{tw}\,\tau}_{[\alpha\beta]n}(x) = 0 = \partial^\beta M^{\mathrm{tw}\,\tau}_{[\alpha\beta]n}(x), \qquad \tau = 2, \tilde{3}, \qquad (4.4.38)$$

and

$$\Box \widehat{M}^{\mathrm{tw3}}_{\alpha n-1}(x) = 0, \qquad \partial^\alpha \widehat{M}^{\mathrm{tw3}}_{\alpha n-1}(x) = 0. \qquad (4.4.39)$$

Let us point to the structural similarities of the operators (4.4.30) and (4.4.36) on the one hand and (4.4.32) and (4.4.34) on the other hand. – Furthermore, we observe that when organizing the various expressions such that everywhere the partial derivatives appear to the right of the kth power of x^2 the differential operations ensuring the symmetry type are transformed according to $\partial_\alpha \to \mathcal{D}_\alpha(k)$ whereas the generalized (first order) inner derivative $x_{[\alpha}\partial_{\beta]}$ remains unchanged because it commutes with x^2. Namely, a short look on our results, Eqs. (4.4.30), (4.4.32), (4.4.33), (4.4.34), (4.4.35) and (4.4.37), shows that the expressions on-cone are obtained by restricting the sums to those terms with $k=0$ and replacing $x \to \tilde{x}$, $\partial_\alpha \to \partial/\partial\tilde{x}^\alpha$. Let us mention that this is nothing else but another form of the harmonic extension which is well-known in group theory [19, 48, 49]. The harmonic extension gives an one-to-one relation between homogeneous polynomials on the light-cone and the corresponding harmonic polynomials off-cone. It is important to note that the unique harmonic extension must not destroy the type of the Lorentz representation.

According to Eqs. (4.4.30), (4.4.32), (4.4.33), (4.4.34), (4.4.35) and (4.4.37) we may introduce the orthogonal projection operators onto the subspaces with definite spin and, therefore, with definite geometric twist $\tau = 2, 3$ as follows:

$$\begin{aligned} O^{(\tau)}_{(5)\alpha n}(x) &= \mathcal{P}^{(\tau)\,\mu}_{\alpha n} O_{(5)\mu n}(x), &(4.4.40)\\ N^{(\tau)}_n(x) &= \mathcal{P}^{(\tau)}_n N_n(x), &(4.4.41)\\ M^{(\tau)}_{[\alpha\beta]n}(x) &= \mathcal{P}^{(\tau)\,[\mu\nu]}_{[\alpha\beta]n} M_{[\mu\nu]n}(x). &(4.4.42)\end{aligned}$$

The corresponding twist projectors onto the light-cone are discussed in more detail in Section 4.1 (see also Ref. [66]).

Resummed harmonic operators with twist $\tau = 2,\, 3$

Now we are able to rewrite the local harmonic operators of definite twist $\tau = 2, 3$ in terms of Gegenbauer polynomials in the x-space analogous to the scalar harmonic operators. In a second step we resum the infinite series (for n) of Gegenbauer polynomials to the related Bessel functions thereby obtaining nonlocal harmonic operators of (the same) definite twist.

---

[12] We use the convention $A_{\alpha\beta\gamma} \equiv \frac{1}{3}(A_{\alpha[\beta\gamma]} + A_{\beta[\gamma\alpha]} + A_{\gamma[\alpha\beta]})$.

For notational simplicity we introduce the following abbreviations for the homogeneous polynomials and their related functions:

$$h^\nu_n(p|x) = \left(\frac{1}{2}\sqrt{p^2x^2}\right)^n C^\nu_n\!\left(\frac{px}{\sqrt{p^2x^2}}\right), \tag{4.4.43}$$

$$\mathcal{H}_\nu(p|x) = \sqrt{\pi}\left(\sqrt{(px)^2 - p^2x^2}\right)^{1/2-\nu} J_{\nu-1/2}\left(\frac{1}{2}\sqrt{(px)^2-p^2x^2}\right)\mathrm{e}^{\mathrm{i}px/2}. \tag{4.4.44}$$

Using the following functional relations of the Gegenbauer and Bessel functions [112],

$$\begin{aligned} mC^\nu_m(z) &= 2\nu\left(zC^{\nu+1}_{m-1}(z) - C^{\nu+1}_{m-2}(z)\right), \\ \frac{\mathrm{d}}{\mathrm{d}z}\left(z^{-\lambda}J_\lambda(z)\right) &= -z^{-\lambda}J_{\lambda+1}(z), \\ z\frac{\mathrm{d}}{\mathrm{d}z}J_\lambda(z) &= -\lambda J_\lambda(z) + zJ_{\lambda-1}(z), \end{aligned}$$

respectively, one obtains the following results for the partial derivations of these functions:

$$\partial_\alpha h^\nu_n(p|x) = \nu\left(p_\alpha h^{\nu+1}_{n-1}(p|x) - \frac{1}{2}p^2x_\alpha h^{\nu+1}_{n-2}(p|x)\right), \tag{4.4.45}$$

$$\partial_\alpha\mathcal{H}_\nu(p|x) = \left(\mathrm{i}p_\alpha(2\nu+1+x\partial) - \frac{1}{2}(\mathrm{i}p)^2x_\alpha\right)\mathcal{H}_{\nu+1}(p|x). \tag{4.4.46}$$

The local twist-2 vector operator[13] is obtained through derivation with respect to x^α of the local twist-2 scalar operator as follows, cf., Eq. (4.4.30),

$$\begin{aligned} O^{\mathrm{tw2}}_{\alpha n}(x) &= \frac{1}{(n+1)^2}\,\partial_\alpha\int\mathrm{d}^4p\,\big(\bar\psi\gamma_\mu\psi\big)(p)\left\{x^\mu\,h^2_n(p|x) - \frac{1}{2}\,p^\mu x^2\,h^2_{n-1}(p|x)\right\} \\ &= \frac{1}{(n+1)^2}\int\mathrm{d}^4p\,\big(\bar\psi\gamma_\mu\psi\big)(p)\Big\{\delta^\mu_\alpha\,h^2_n(p|x) - p^\mu x_\alpha\,h^2_{n-1}(p|x) \\ &\qquad + 2x^\mu p_\alpha\,h^3_{n-1}(p|x) - \big(x^\mu x_\alpha p^2 + p^\mu p_\alpha x^2\big)\,h^3_{n-2}(p|x) + \frac{1}{2}\,p^\mu x_\alpha x^2p^2\,h^3_{n-3}(p|x)\Big\}, \end{aligned} \tag{4.4.47}$$

and the corresponding resummed nonlocal twist-2 vector operator reads

$$\begin{aligned} O^{\mathrm{tw2}}_\alpha(x,-x) &= \partial_\alpha\int_0^1\mathrm{d}t\int\mathrm{d}^4p\,\big(\bar\psi\gamma_\mu\psi\big)(p)\left\{x^\mu\,(2+p\partial_p) - \frac{1}{2}\,\mathrm{i}\,tp^\mu\,x^2\right\}(3+p\partial_p)\;\mathcal{H}_2(p|tx) \\ &= \int\mathrm{d}^4p\,\big(\bar\psi\gamma_\mu\psi\big)(p)\,(2+p\partial_p)\int_0^1\mathrm{d}t\Big\{\Big[(3+p\partial_p)\,\delta^\mu_\alpha - \mathrm{i}tp^\mu x_\alpha\Big]\mathcal{H}_2(p|tx) \\ &\quad + \Big[(3+p\partial_p)\Big((4+p\partial_p)\,\mathrm{i}tp_\alpha x^\mu - \frac{1}{2}(\mathrm{i}t)^2\big(p^2x^\mu x_\alpha + x^2p^\mu p_\alpha\big)\Big) + \frac{1}{4}(\mathrm{i}t)^3p^\mu p^2x_\alpha x^2\Big]\mathcal{H}_3(p|tx)\Big\}. \end{aligned} \tag{4.4.48}$$

Here, the equality $1/(n+1) = \int_0^1\mathrm{d}t\,t^n$ has been used; analogous equalities will be used in the following.

---

[13] The corresponding local tensor operator of 2nd rank is given by: $O^{\mathrm{tw2}}_{\alpha\beta n-1}(x) = 1/n\,\partial_\beta O^{\mathrm{tw2}}_{\alpha n}(x)$.

The resummed local twist-3 vector operator reads, cf., Eq. (4.4.32):

$$\begin{aligned}
O^{\mathrm{tw3}}_{\alpha n}(x) &= \frac{2}{n+1}\, x^\beta \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]} - \frac{1}{n+1}\, x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \int \mathrm{d}^4p\, \big(\bar\psi\gamma_\mu\psi\big)(p)\, h^1_n(p|x) \\
&= \frac{1}{(n+1)^2} \int \mathrm{d}^4p\, \big(\bar\psi\gamma_\mu\psi\big)(p) x^\beta \Big\{ 2(n+1)\delta^\mu_{[\alpha}p_{\beta]}\, h^2_{n-1}(p|x) + (n+2)\, x_{[\alpha}\delta^\mu_{\beta]}p^2\, h^2_{n-2}(p|x) \\
&\qquad\qquad - 4x_{[\alpha}p_{\beta]}p^\mu\, h^3_{n-2}(p|x) + 2x_{[\alpha}p_{\beta]}x^\mu p^2\, h^3_{n-3}(p|x) \Big\},
\end{aligned} \tag{4.4.49}$$

and the bilocal twist-3 vector operator is given by

$$\begin{aligned}
O^{\mathrm{tw3}}_{\alpha}(x,-x) &= 2 \int \mathrm{d}^4p\, \big(\bar\psi\gamma_\mu\psi\big)(p)\; x^\beta \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]}\,(1+p\partial_p) - x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \int_0^1 \mathrm{d}t\, \mathcal{H}_1(p|tx) \\
&= \int \mathrm{d}^4p\, \big(\bar\psi\gamma^\mu\psi\big)(p) \int_0^1 \mathrm{d}t\; (2+p\partial_p)\; x^\beta \Big\{ \Big[2\,\mathrm{i}t\, \delta^\mu_{[\alpha}p_{\beta]}(2+p\partial_p)\; - (\mathrm{i}t)^2\delta^\mu_{[\alpha}x_{\beta]}p^2 \Big] \mathcal{H}_2(p|tx) \\
&\qquad\qquad - \mathrm{i}t\, x_{[\alpha}p_{\beta]} \Big[2\,\mathrm{i}t\, p^\mu\,(5+p\partial_p) - (\mathrm{i}t)^2 x^\mu p^2 \Big]\, \mathcal{H}_3(p|tx) \Big\}.
\end{aligned} \tag{4.4.50}$$

The local twist-3 chiral-odd scalar operator will be given only for completeness, cf., Eq. (4.4.3):

$$N^{\mathrm{tw3}}_n(x) = \int \mathrm{d}^4p\, \big(\bar\psi\psi\big)(p)\, h^1_n(p|x), \tag{4.4.51}$$

and for the corresponding bilocal one, one gets

$$N^{\mathrm{tw3}}(x,-x) = \int \mathrm{d}^4p\, \big(\bar\psi\psi\big)(p)\,(1+p\partial_p)\; \mathcal{H}_1(p|x). \tag{4.4.52}$$

The local resummed twist-2 chiral-odd skew tensor operator is given by, cf., Eq. (4.4.34),

$$\begin{aligned}
M^{\mathrm{tw2}}_{[\alpha\beta]n}(x) =& \frac{2}{(n+2)(n+1)} \Big\{ \delta^\mu_{[\alpha}\partial_{\beta]} - \frac{1}{n+2}\, x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \\
&\qquad \times \int \mathrm{d}^4p\, \big(\bar\psi\sigma_{\mu\nu}\psi\big)(p) \Big\{ x^\nu\, h^2_n(p|x) - \frac{1}{2}\, p^\nu\, x^2\, h^2_{n-1}(p|x) \Big\},
\end{aligned} \tag{4.4.53}$$

and the related bilocal twist-2 skew tensor operator reads

$$\begin{aligned}
M^{\mathrm{tw2}}_{[\alpha\beta]}(x,-x) = 2 \int \mathrm{d}^4p\, \big(\bar\psi\sigma_{\mu\nu}\psi\big)(p) \int_0^1 \mathrm{d}t\, t \Big\{ (2+p\partial_p)\, \delta^\mu_{[\alpha}\partial_{\beta]} - x_{[\alpha}\partial_{\beta]}\partial^\mu \Big\} \\
\times \Big[x^\nu\,(3+p\partial_p) - \frac{1}{2}\,\mathrm{i}\, t p^\nu x^2 \Big] \mathcal{H}_2(p|tx).
\end{aligned} \tag{4.4.54}$$

For the related, resummed local twist-3 skew tensor operator resulting from the trace terms we obtain

$$M^{\mathrm{tw3}}_{[\alpha\beta]n}(x) = \frac{2}{(n+2)n}\, x_{[\alpha}\partial_{\beta]} \int \mathrm{d}^4p\, \big(\bar\psi\sigma_{\mu\nu}\psi\big)(p)\, p^\mu x^\nu\, h^2_{n-1}(p|x), \tag{4.4.55}$$

and get the following twist-3 bilocal skew tensor

$$M^{\mathrm{tw3}}_{[\alpha\beta]}(x,-x) = 2x_{[\alpha}\partial_{\beta]} \int \mathrm{d}^4p\, \big(\bar\psi\sigma_{\mu\nu}\psi\big)(p)\, \mathrm{i}\, p^\mu x^\nu (2+p\partial_p) \int_0^1 \mathrm{d}t\, \mathcal{H}_2(p|tx). \tag{4.4.56}$$

In addition, the independent local twist-3 skew tensor having symmetry type $[n{+}2] = (n,1,1)$ with Lorentz structure $\left(\frac{n}{2},\frac{n}{2}\right)$ reads

$$\widetilde{M}^{\rm tw3}_{[\alpha\beta]n}(x) = \frac{3}{n+2}\, x^\gamma \Big\{ \delta^\mu_{[\alpha}\delta^\nu_\beta\partial_{\gamma]} - \frac{2}{n}\,\delta^\mu_{[\alpha}x_\beta\partial_{\gamma]}\partial^\nu \Big\} \int {\rm d}^4p\, \big(\bar\psi\sigma_{\mu\nu}\psi\big)(p)\, h^1_n(p|x) \tag{4.4.57}$$

and leads to the following expression for the twist-3 bilocal skew tensor

$$\widetilde{M}^{\rm tw3}_{[\alpha\beta]}(x,-x) = 3x^\gamma \Big\{ \delta^\mu_{[\alpha}\delta^\nu_\beta\partial_{\gamma]}x\partial - 2\,\delta^\mu_{[\alpha}x_\beta\partial_{\gamma]}\partial^\nu \Big\} \int {\rm d}^4p\, \big(\bar\psi\sigma_{\mu\nu}\psi\big)(p)\,(1+p\partial_p) \\ \times \int_0^1 {\rm d}t\, \frac{1-t^2}{2t}\, \mathcal{H}_1(p|tx). \tag{4.4.58}$$

Let me note that we can use the formulas (4.4.47) and (4.4.49) to calculate the target mass corrections of the Compton amplitude in the q-space. In order to do this one has to use the operators $O^{\rm tw2}_{(5)\alpha n}(q)$ and $O^{\rm tw3}_{(5)\alpha n}(q)$ which are multiplied by the coefficient functions in the q-space. In the forward case one finds the target mass corrections for the Bjorken sum rules and the Nachtmann moments (see Refs. [18, 124, 104]). Additionally, one is able to calculate the power corrections for the non-forward case, too.

Is also possible to rewrite all harmonic operators in terms of Gegenbauer polynomials and Bessel functions for $2h$-dimensional spacetime.

Chapter 5

Conclusions and Outlook

A unique group theoretical procedure was introduced which allows a decomposition of nonlocal tensor-valued light-ray operators and off-cone operators into (tensorial) operators and harmonic operators, respectively, with definite geometric twist. This decomposition is completely of group theoretical origin and is equivalent to the decomposition of local operators with respect to irreducible tensor representations of the Lorentz group. The explicite twist decomposition in $1+3$-spacetime dimensions has been given for all physically relevant bilocal quark and gluon tensor operators which appear, e.g., in the virtual Compton scattering in the generalized Bjorken region as well as for multilocal operators containing many quark and gluon fields, e.g., Shuryak-Vainshtein type operators, four-quark and four gluon operators, and three-gluon operators. In the case of light-ray operators the twist projectors are given in terms of interior differential operators, which were introduced by Bargmann and Todorov [19]. By the help of these interior differential operators the conditions of tracelessness can be easily formulated and satisfied. In the case of off-cone operators the local operators are given in terms of Gegenbauer polynomials and the nonlocal ones are expressed in terms of Bessel functions. There are one-to-one relations between the light-ray and off-cone operators with definite geometric twist. The light-ray operators are obtained through the projection of the off-cone operators onto the light-cone. In the other direction, the off-cone operators can be understood as the harmonic extension of the light-ray operators. Because we discussed the relations for arbitrary $2h$-spacetime dimensions one can also construct twist operators in higher or lower dimensions in straightforward manner. Perhaps this fact can be useful in order to renormalize the QCD operators by the help of dimensional regularization. Another interesting fact, from the mathematical point of view, may be the generalization of the twist decomposition for more complicated symmetry types. Perhaps such (higher) twist operators will be needed from the phenomenological point of view in future. Additionally, one can use the polynomial technique to construct a general mathematical theory of $SO(2h)$ or $SO(1,2h-1)$ irreducible tensor polynomials which do not exist in the mathematical literature. Then one will be able to compute the infinite twist decomposition of any multilocal tensor operator.

We calculated the forward and vacuum-to-vector meson matrix elements of bilocal light-ray operators with definite geometric twist. In this doing we were able to classify the corresponding quark distribution and vector meson distribution amplitudes with respect to definite geometric twist in a unique manner. Concerning higher twists we introduced new parton distribution functions and ρ-meson distribution amplitudes, as wells as their moments, having definite geometric twist. We were able to derive one-to-one relations between these new distribution amplitudes and the conventional ones with dynamical twist, which were earlier introduced by Jaffe and Ji [79] and by Ball *et al.* [14]. Thereby, Wandzura-Wilczek type relations and Burkhardt-Cottingham type sum rules for the dynamical distribution amplitudes, which should be tested experimentally, have been obtained for higher twists.

By calculating the (general) non-forward matrix elements of off-cone operators with definite geometric twist we introduced for the first time all geometric twist-2 and twist-3 double distribution amplitudes which are used from the phenomenological point of view. In the case of operators whose moments are related to totally symmetric tensors we were able to determine the complete twist decomposition up to infinite geometric twist. Using these bilocal operators we gave the classification of the (pseudo) scalar meson distribution amplitudes for all twists. An important result of our approach is that we were able to separate all kinematical factors like x^2, xP, and xS from the the distribution amplitudes and to resum the power corrections to all orders. This is based on the group-theoretical content of the procedure of trace subtraction from the local operators with definite symmetry type. Because the distribution amplitudes of definite geometric twist are already determined by the nonlocal matrix elements on the light-cone their renormalization properties are determined by the light-cone operators only. The nonlocal matrix elements off-cone contain supplementary kinematical structures which correspond to the structure of the harmonic extension of the matrix element on-cone.

Additionally, by means of these harmonic QCD operators one is able to calculate target and quark mass corrections in the Compton amplitude of non-forward scattering processes, like deeply virtual Compton scattering and diffractive scattering. Hopefully, this will be useful for predictions of target mass effects in physical cross sections of these processes.

Acknowledgments

The author is grateful to B. Geyer, P. Ball, J. Blümlein, V.K. Dobrev, J. Eilers, D. Müller, S. Neumeier, D. Robaschik and C. Weiss for various stimulating and useful discussions. The author thanks J. Eilers for the possibility to compare the decomposition of the 2nd rank tensor operator with his result which was very useful to find some mistakes. In addition, he gratefully acknowledges the Graduate College “Quantum field theory” at the Center for Theoretical Studies of Leipzig University for financial support.

Appendix A

Symmetry classes and tensor representations

In this Appendix we collect some facts about tensor representations of the classical matrix groups G and their relation to the symmetric group S_n which are of relevance in our consideration (see, e.g., [20, 72]). Given any irreducible matrix representation,

$$G \ni g \mapsto \hat{g} \in \mathcal{B}(V): \qquad \hat{g} e_i = e_j \, a^j{}_i, \tag{A.1}$$

on some vector space V (of dimension $\dim V$) with base $\{e_i,\, i = 1, 2, \ldots, \dim V\}$. The direct products of these representations act on the *tensor space* $T^n\, V$ which is build as tensor product of n copies of the vector space V:

$$T^n\, V \;\; = \;\; (V \otimes \ldots \otimes V)\,.$$

The elements of the tensor space $T^n\, V$ are the n–times contravariant *tensors* $\mathbf{t} = t^{i_1 \ldots i_n} e_{i_1 \ldots i_n}$, where $t^{i_1 \ldots i_n}$ are the *components* of $\mathbf{t}$ in the base $e_{i_1 \ldots i_n} \equiv \big(e_{i_1} \otimes \ldots \otimes e_{i_n}\big)$, $i_k \in [1, \ldots, \dim V]$. [1] The components of the tensors of rank n transform under G according to

$$(t')^{i_1 \ldots i_n} \equiv t^{i'_1 \ldots i'_n} = a^{i'_1}{}_{i_1} \ldots a^{i'_n}{}_{i_n} t^{i_1 \ldots i_n}. \tag{A.2}$$

In the following we shall comment on the connection between the irreducible representations of the general linear group $GL(N, \mathbb{C})$ (together with its subgroups) and of the symmetric group S_n on these tensor spaces. Let us, at the moment consider only the groups $G = GL(N, \mathbb{C}), SL(N, \mathbb{C}), SU(N)$. Given any (irreducible) representation of G on V, then its n–fold tensor product defines a reducible representation $A^{(n)}$ which, together with the reducible representation $\hat{\pi}$ of S_n, is determined by:

$$\begin{aligned} \hat{A}^{(n)}(g)\mathbf{t} \equiv \;& \hat{A}^{(n)}(g)(t^{i_1 \ldots i_n} e_{i_1 \ldots i_n}) &:=\;& t^{i_1 \ldots i_n} (\hat{g}e)_{i_1} \otimes \ldots \otimes (\hat{g}e)_{i_n}, \quad \forall g \in G, \\ \hat{\pi}\mathbf{t} \equiv \;& \hat{\pi}(t^{i_1 \ldots i_n} e_{i_1 \ldots i_n}) &:=\;& t^{\pi(i_1 \ldots i_n)} e_{i_1} \otimes \ldots \otimes e_{i_n}, \quad \forall \pi \in S_n; \end{aligned}$$

therefore the action of both groups on the tensors $\mathbf{t}$ commutes [2]

$$\big[\hat{A}^{(n)}(g), \hat{\pi}\big]\mathbf{t} = 0. \tag{A.3}$$

---

[1] Here, neither covariant tensors which act on direct products of the dual space V^* nor mixed ones are considered since, for the case under consideration, the indices may be raised or lowered by the metric tensor.

[2] This results from the fact that the product of the matrix elements a_i^j in Eq. (A.2) is *bisymmetric* under S_n.

Of course, the same conclusion holds for the elements $\sum_\pi \alpha(\pi)\hat{\pi}, \alpha(\pi) \in \mathbb{R}$, of the group algebra $\mathcal{R}[S_n]$ of the symmetric group. This algebra, considered as a vector space, carries the *regular representation* of S_n which is known to be fully reducible.

The irreducible representations $\Delta^{[m]}$ of the symmetric group are uniquely determined by the idempotent (normalized) Young operators

$$\mathcal{Y}_{[m]} = \frac{f_{[m]}}{n!}\mathcal{P}\mathcal{Q} \quad \text{with} \quad \mathcal{P} = \sum_{p\in H_{[m]}} p, \quad \mathcal{Q} = \sum_{q\in V_{[m]}} \delta_q \, q, \tag{A.4}$$

$$\mathcal{Y}_{[m]}\mathcal{Y}_{[m']} = \delta_{[m][m']}\mathcal{Y}_{[m]}, \tag{A.5}$$

which are related to corresponding Young tableaux being denoted by $[m]$. If

$$\underline{m} = (m_1, m_2, \ldots m_r) \quad \text{with} \quad m_1 \geq m_2 \geq \ldots \geq m_r, \quad \sum_{i=1}^{r} m_i = n, \tag{A.6}$$

defines a Young pattern (Fig. A.1) then a *Young tableau* $[m]$ is obtained by putting in (without repetition) the indices $i_1, \ldots i_n$ – corresponding to different "places" within the direct product – and $H_{[m]}$ and $V_{[m]}$ denotes their horizontal and vertical permutations with respect to $[m]$. A *standard tableau* is obtained when the indices $i_1, \ldots, i_n$ are ordered lexicographically. There are

$$f_{[m]} = n! \, \frac{\prod_{i<j}(l_i - l_j)}{\prod_{i=1}^{r} l_i!} \quad \text{with} \quad l_i = m_i + r - i, \quad \sum_{[m]} f_{[m]}^2 = n!, \tag{A.7}$$

different standard tableaux which correspond to $f_{[m]}$ different, but equivalent, irreducible representations of S_n whose dimension is given also by $f_{[m]}$. The (normalized) Young operators $\mathcal{Y}_{[m]}$ according to (A.5) project onto mutually orthogonal irreducible left ideals of $\mathcal{R}[S_n]$.



Figure A.1: Young pattern $\underline{m}$

In this way the tensor product of n spaces V carries the regular representation of the group S_n and decomposes into $\sum_{[m]} f_{[m]}$ irreducible subspaces containing tensors of symmetry class $[m]$. However, the tensors actually are specified by taking for the indices $i_1, \ldots, i_n$ arbitrary values from the range $[1, \ldots, \dim V]$. Therefore, with respect to the group G the standard tableau is defined by putting into the Young pattern the *values of the indices* i_k such that they are non–decreasing from left to right and increasing from top to bottom! Because of Eq. (A.3) by Schur's Lemma it follows that the representation $\hat{A}^{(n)}(g)$ of G is reducible and it decomposes into as many irreducible representations as there are (in general reducible) representations of S_n, i.e., any tensor representation of the group G is characterized by a standard tableau,

$$\hat{A}^{(n)}(g) = \bigoplus_{[m]} R^{[m]}(g) \otimes E_{l_{[m]}}, \quad \hat{\pi} = \bigoplus_{[m]} E_{r_{[m]}} \otimes \Delta^{[m]}(\pi), \tag{A.8}$$

where $E_{l_{[m]}}$ is the unit matrix; more explicitly we have

$$\hat{A}^{(n)}(g)_{[m]\kappa\alpha,[m']\kappa'\alpha'} \;=\; \delta_{[m][m']}\delta_{\alpha\alpha'}R^{[m]}_{\kappa\kappa'}(g), \tag{A.9}$$

where $\kappa = 1,\ldots,r_{[m]}$ counts different equivalent irreducible representations of S_n and $\alpha = 1,\ldots,l_{[m]} \leq f_{[m]}$ counts the basis elements of these representations. The representations $R^{[m]}(g)$ of G in $T^n\,V$ are characterized by equivalent Young operators $\widehat{\mathcal{Y}}_{[m]} = \mathcal{QP}$: [3]

$$T^n\,V \;=\; \underset{[m]}{\oplus}\,\underset{\kappa}{\oplus}\widehat{\mathcal{Y}}_{[m]\kappa}(T^n\,V). \tag{A.10}$$

The invariant subspaces projected out by $\widehat{\mathcal{Y}}_{[m]\kappa}(T^n\,V)$ are irreducible with respect to $GL(N,\mathbb{C})$, $SL(N,\mathbb{C})$ and $SU(N)$. After restriction onto these subgroups some of the irreducible representations become equivalent ones. However, for $O(N,\mathbb{C}), SO(N,\mathbb{C})$ and $Sp(2\nu)$ these representations, in general, are not irreducible and decompose further.

Concerning the orthogonal groups from the subspaces of the above introduced symmetry classes only the completely antisymmetric ones (for $n \leq N$) remain irreducible. The reason is that, because of the very definition of the orthogonal groups, $\delta_{ij}{a^i}_k{a^j}_l \;=\; \delta_{kl},\ \forall a \in O(N)$, the operation of taking the trace of a tensor commutes with the orthogonal transformations of that tensor:

$$\mathrm{tr}\,\mathbf{T}' \;=\; \delta_{ij}T'^{ij} \;=\; \delta_{ij}{a^i}_k{a^j}_l T^{kl} \;=\; \mathrm{tr}\,\mathbf{T}. \tag{A.11}$$

Again, by Schur's Lemma, irreducible subspaces of $O(N)$ are spanned by *traceless* tensors having definite symmetry class. This decomposition is obtained as follows:

$$T^{i_1 i_2 \ldots i_n}_{[m]} \;=\; \overset{\circ}{T}{}^{i_1 i_2 \ldots i_n}_{[m]} + \sum_{1\leq r,s\leq n} \delta^{i_r i_s}\, T^{i_1\ldots i_{r-1} i_{r+1}\ldots i_{s-1} i_{s+1}\ldots i_n}_{[m-2]}. \tag{A.12}$$

The tensors which appear under the sum have degree $n-2$ and a Young pattern $[m-2] \subset [m]$ obtained by removing two boxes from the (right) border without destroying the property (B.2) and (B.3) to be a pattern. They may be decomposed again into traceless ones plus some remainder, and so on. Therefore, a traceless tensor is obtained from the original one by successively subtracting the traces.

---

[3] Note, that the symmetrizations $\mathcal{P}$ and the antisymmetrizations $\mathcal{Q}$ with respect to $[m]$ are interchanged. This corresponds, in the terminology of quantum mechanics, to "quantum number permutations" instead of the "place permutations" above. And, contrary to $\mathcal{Y}_{[m]}$ which defines a left ideal in $\mathcal{R}$, $\widehat{\mathcal{Y}}_{[m]}$ defines a right ideal.

Appendix B

Irreducible tensor representations of the Lorentz group

In this Appendix the group theoretical background will be given which is used for the construction of the nonlocal operators whose local parts – which are obtained by Taylor expansion – transform according to irreducible tensor representations under the Lorentz group, i.e., have definite spin and therefore also well defined twist. These tensor representations are characterized by a specific symmetry class with respect to the symmetric group and are determined by a few types of Young tableaux.

The Lie algebra of the Lorentz group $\mathcal{L}_+^\uparrow$ is characterized by the generators of the three spatial rotations $\boldsymbol{M}$ and the three boosts $\boldsymbol{N}$; from it the two (complex) linear combinations $\boldsymbol{M}_\pm$ may be build which define two independent $SO(3)$ groups:

$$[M_\pm^i, M_\pm^j] = \mathrm{i}\epsilon^{ijk} M_\pm^k, \qquad [M_+^i, M_-^j] \; = \; 0 \quad \text{with} \quad \boldsymbol{M}_\pm = \boldsymbol{M} \pm \mathrm{i}\boldsymbol{N}.$$

Therefore, the (Lie algebra of the) complex Lorentz group is isomorphic to the (Lie algebra of the) direct product $SO(3,\mathbb{C}) \otimes SO(3,\mathbb{C}) \simeq SO(4,\mathbb{C})$. This characterization makes use of the fact that the irreducible (finite) representations

$$\mathcal{D}^{(j_+,j_-)}(\boldsymbol{\varphi},\boldsymbol{\vartheta}) \;=\; \mathcal{D}^{(j_+)}(\boldsymbol{\varphi} - \mathrm{i}\boldsymbol{\vartheta}) \otimes \mathcal{D}^{(j_-)}(\boldsymbol{\varphi} + \mathrm{i}\boldsymbol{\vartheta}) \tag{B.1}$$

of the restricted orthochronous Lorentz group $\mathcal{L}_+^\uparrow \simeq SO(1,3;\mathbb{R})$, where $\boldsymbol{\varphi}$ and $\boldsymbol{\vartheta}$ are the angle of rotation and the rapidity of the boost transformations, are determined by two numbers $(j_+, j_-), j_\pm = 0, 1/2, 1, \cdots$, which define the spin $j = j_+ + j_-$ of the representation.

Let us now state some general results concerning finite dimensional representations of the (complex) orthogonal groups (see, e.g., Chapters 8 and 10 of Ref. [20]):

(1) The group $SO(N,\mathbb{C})$ has two series of complex–analytic[1] irreducible representations. Every representation of the first (resp. the second) series determines and is in turn determined by a highest weight $\underline{m} = (m_1, m_2, \ldots, m_\nu)$ whose components m_i are integers (resp. half–odd integers) and satisfy the conditions

$$m_1 \geq m_2 \geq \ldots \geq m_{\nu-1} \geq |m_\nu| \quad \text{for} \quad N = 2\nu, \tag{B.2}$$

$$m_1 \geq m_2 \geq \ldots \geq m_{\nu-1} \geq m_\nu \geq 0 \quad \text{for} \quad N = 2\nu + 1. \tag{B.3}$$

The first series determines the tensor representations, whereas the second series determines the spinor representations.

---

[1] A representation is complex–analytic if it depends analytically on the group parameters.

(2) A tensor representation of $SO(N;\mathbb{C})$, for either $N = 2\nu$ or $N = 2\nu + 1$, being determined by the highest weight $\underline{m} = (m_1, m_2, \ldots, m_\nu)$, m_i integer, is equivalent to another tensor representation $T_{i_1 i_2 \ldots i_n}$, $n = \sum m_i$, which is realized in the space of traceless tensors whose symmetry class is characterized by a Young pattern defined by the partition $[m] = (m_1, m_2, \ldots, m_\nu)$.

(3) Furthermore, if $\mathcal{T}_{G_c}$ is a complex–analytic representation of a (complex) semi-simple Lie group G_c and $\mathcal{T}_{G_r}$ is the restriction of $\mathcal{T}_{G_c}$ to a real form G_r of G_c, then, $\mathcal{T}_{G_c}$ is irreducible (fully reducible) iff $\mathcal{T}_{G_r}$ is irreducible (fully reducible). As a consequence, any irreducible representation of the complex group remains irreducible if restricted to a real subgroup and, on the other hand, from any irreducible representation of the real group by analytic continuation in the group parameters an irreducible representation of its complexification is obtained.

Therefore, in order to investigate irreducible tensor representations of the Lorentz group we may consider equally well irreducible tensor representations of the complex group $SO(4,\mathbb{C})$. Even more, any irreducible representation of the orthogonal group $SO(4)$ by analytic continuation induces an irreducible representation of $SO(4,\mathbb{C})$, which by restriction to their real subgroup $\mathcal{L}_+^\uparrow$ subduces an irreducible representation of the Lorentz group. Since tensor representations of the orthogonal group are uniquely determined by the symmetry class $[m]$ of their tensors we may study the irreducible tensor representations of the Lorentz group through a study of the corresponding Young tableaux which determine the irreducible representations of the symmetric group S_n of permutations.

This characterization of tensor representations through their symmetry class holds for any of the classical matrix groups, $GL(N,\mathbb{C})$ and their various subgroups (see also Appendix A). However, for $O(N,\mathbb{C})$ and $SO(N,\mathbb{C})$ these representations, in general, are not irreducible. The reason is that taking the trace of a tensor commutes with the orthogonal transformations. Therefore, by Schur's Lemma, irreducible subspaces of $SO(N,\mathbb{C})$ are spanned by *traceless* tensors having definite symmetry class.

From the requirement of tracelessness the following restrictions obtain (see, e.g., [72]): The only Young tableaux being relevant are those whose first two columns are restricted to have length $\mu_1 + \mu_2 \leq N$ (see (i) – (iv) below). Two representations R and R' whose first columns are related by $\mu_1' = N - \mu_1$, where $\mu_1 \leq N/2$, are called associated; if $\mu_1 = \mu_1' = N/2$ this representation is called selfassociated. After restriction to the subgroup $SO(N) \subset O(N)$ associated representations are equivalent, whereas selfassociated representations decompose into two nonequivalent irreducible representations.

Let us illustrate this for the Lorentz group by the representations which will be of interest in the following. First of all, vector and axial vector representations V_μ and $A_\mu = \epsilon_{\mu\nu\kappa\lambda} A^{\nu\kappa\lambda}$, respectively, are associated ones, and antisymmetric tensor representations, $A_{\mu\nu} = T_{\mu\nu} - T_{\nu\mu}$, are selfassociated which, if restricted to $SO(1,3)$ decompose into the (anti-)selfdual tensors $A^{\pm}_{\mu\nu} = \frac{1}{2}(A_{\mu\nu} \mp \frac{1}{2}\,\mathrm{i}\epsilon_{\mu\nu\kappa\lambda} A^{\kappa\lambda})$. Since later on only representations of the orthochronous Lorentz group containing the parity operation are considered, this distinction is of no relevance. Therefore, any tensor of second order, $T_{\mu\nu}$, may be decomposed according to

$$T_{\mu\nu} \;=\; \underbrace{\tfrac{1}{2}\left(T_{\mu\nu} + T_{\nu\mu}\right) - \tfrac{1}{4} g_{\mu\nu}\, T_\rho{}^\rho}_{S_{\mu\nu}} + \underbrace{\tfrac{1}{2}\left(T_{\mu\nu} - T_{\nu\mu}\right)}_{A_{\mu\nu}} + \tfrac{1}{4}\, g_{\mu\nu} T_\rho{}^\rho . \tag{B.4}$$

Let us denote by $\mathbf{T}(j_+, j_-)$, $j_+ + j_-$ integer, the space of tensors which carry an irreducible representation $\mathcal{D}^{(j_+,j_-)}$ of the Lorentz group. Then from Eq. (B.4) we read off: $S_{\mu\nu} \in \mathbf{T}(1,1)$ is a symmetric traceless tensor, $A_{\mu\nu} = A^{+}_{\mu\nu} + A^{-}_{\mu\nu} \in \mathbf{T}(1,0) \oplus \mathbf{T}(0,1)$ are the selfdual and the anti-selfdual antisymmetric tensors, and $T_\rho{}^\rho \in \mathbf{T}(0,0)$ corresponds to the trivial representation defined through the unit tensor. This decomposition corresponds to the Clebsch-Gordan decomposition of the direct product of two vector representations:

$$\left(\tfrac{1}{2},\tfrac{1}{2}\right) \otimes \left(\tfrac{1}{2},\tfrac{1}{2}\right) \;=\; (1,1) \oplus \Big((1,0) \oplus (0,1)\Big) \oplus (0,0). \tag{B.5}$$

Now we consider the decomposition of the space of tensors of rank n into irreducible representation spaces of $SO(1,3;\mathbb{R})$. The different symmetry classes are strongly restricted by the requirement that only such Young patterns $[m]$ are allowed for which the sum of the first two columns is lower or equal to four. Therefore, only the following Young patterns correspond to nontrivial irreducible representations by traceless tensors:

i.  $j = n, n-2, n-4, \dots$

ii.  $j = n-1, n-2, n-3, \dots$

iii.  $j = n-2, n-3, \dots$

iv.  $j = n-2, n-3, \dots$

In addition, for $n = 4$, also the completely antisymmetric tensor of rank 4 which is proportional to $\epsilon_{\mu\nu\kappa\lambda}$, and therefore equivalent to the trivial representation is allowed.

For the cases (i) – (iv) the minimal spin j – depending on n being either even or odd – will be zero or one. In the case of symmetry type (iv) we have given only one special Young pattern; in principle the length of the second row may contain up to $m_2 = [\frac{n}{2}]$ boxes, and then the maximal spin is given by $j = n - m_2$. — The representations corresponding to symmetry class (i) are associated to representations of the symmetry class (iii) with $n+2$ boxes. The representations corresponding to symmetry class (ii) and (iv) are selfassociated. The symmetry class (ii) contains two non-equivalent parts being related to $(\frac{n}{2}, \frac{n}{2}-1)$ and $(\frac{n}{2}-1, \frac{n}{2})$; the symmetry class (iv) contains three nonequivalent parts related to $(\frac{n}{2}, \frac{n}{2}-2)$, $(\frac{n}{2}-1, \frac{n}{2}-1)$ and $(\frac{n}{2}-2, \frac{n}{2})$; and so on. Any tensor whose symmetry class does not coincide with one of the above classes vanishes identically due to the requirement of tracelessness.

There are two possible ways to construct the non-vanishing tensors. Either one symmetrizes the indices according to the corresponding (standard) Young tableaux and afterwards subtracts the traces, or one starts from tensors being already traceless and finally symmetrizes because this does not destroy the tracelessness. For practical reasons the latter procedure seems to be preferable and will be used in the construction of irreducible light–cone operators of definite twist.

Appendix C

The interior derivative on the light-cone and light-cone operators with definite geometric twist

In this Appendix we would like to discuss some properties of the so-called internal derivatives on the light–cone.

Let us now introduce the notion of *interior differential operators* on the complex light-cone

$$K_{2h}(\mathbb{C}) = \Big\{ z \in \mathbb{C}^{2h};\ z^2 = z_1^2 + \ldots + z_{2h}^2 = 0 \Big\}, \tag{C.1}$$

and on the real light-cone

$$K_{2h}(\mathbb{R}) = \Big\{ \tilde{x} \in \mathbb{M}^{2h};\ \tilde{x}^2 = \tilde{x}_1^2 - \tilde{x}_2^2 - \ldots - \tilde{x}_{2h}^2 = 0 \Big\}, \tag{C.2}$$

respectively. The $2h$-dimensional Minkowski space $\mathbb{M}^{2h}$ has the signature $(+-\cdots-)$.

For the first time, an interior differential operator has been used in order to characterize (irreducible) symmetric tensor representations U^n of $SO(2h)$ on the complex light-cone. In this turn one may consider the *graded algebra*

$$P \equiv P(K_{2h}^n) = \bigoplus_{n=0}^{\infty} K_{2h}^n, \tag{C.3}$$

where K_{2h}^n is the space of homogeneous polynomials $T_n(\tilde{x})$ of degree n on the cone [19, 48]. The reducible representation

$$U(\Lambda) f(z) = f(\Lambda^{-1} z), \qquad \Lambda \in SO(2h), \quad f \in P, \tag{C.4}$$

of $SO(2h)$ generates all symmetric tensor representations U^n

$$U = \bigoplus_{n=0}^{\infty} U^n, \tag{C.5}$$

which are given by the restriction of U to K_{2h}^n [19].

The real light-cone provides a convenient realization of the carrier space for the symmetric tensor representations of the group $SO(1, 2h-1)$. In the following we consider the real light-cone.

A differential operator Q is said to be an *interior* differential operator iff

$$Q\big(x^2 T_n(x)\big)\big|_{x^2\equiv\tilde{x}^2=0} = 0. \tag{C.6}$$

For example, the generators of dilation $X = h - 1 + \tilde{x}\tilde{\partial}$ and rotations $X_{\mu\nu} = \tilde{x}_\nu\tilde{\partial}_\mu - \tilde{x}_\mu\tilde{\partial}_\nu$ are first order interior differential operators on the light-cone. Obviously, they leave the space K^n_{2h} *invariant*. The interior operators on P form a (complex) algebra under addition and multiplication [19].

A further interior differential operator d_μ of second order may be introduced by the following requirements:
(a) it should be a *lowering operator*, i.e. mapping a homogeneous polynomial of degree n (of $\tilde{x}$) into a homogeneous polynomial of degree $n-1$, $K^n_{2h} \to K^{n-1}_{2h}$;
(b) it should behave as a vector under rotations,

$$[X_{\mu\nu}, \mathrm{d}_\lambda] = \delta_{\mu\lambda}\mathrm{d}_\nu - \delta_{\nu\lambda}\mathrm{d}_\mu; \tag{C.7}$$

(c) it should be the lowest order differential operator satisfying (a) and (b).
Choosing the normalization of this interior derivative so that $2\mathrm{d}_\mu$ is the generator of *special conformal transformations* of *massless* 0-helicity representations of the conformal Lie algebra $\mathfrak{so}(2,2h)$, the interior derivative is given as

$$\mathrm{d}_\mu f(\tilde{x}) \equiv \Big\{\big(h - 1 + x\partial\big)\partial_\mu - \tfrac{1}{2}x_\mu\Box\Big\} f(x)\big|_{x=\tilde{x}}, \tag{C.8}$$

with the following properties

$$\mathrm{d}^2 = 0\,, \quad [\mathrm{d}_\mu, \mathrm{d}_\nu] = 0 \quad \text{and} \quad \mathrm{d}_\mu\tilde{x}^2 = \tilde{x}^2\big(\mathrm{d}_\mu + 2\tilde{\partial}_\mu\big)\,. \tag{C.9}$$

Obviously, d_μ is a second order interior differential operator which together with $\tilde{x}_\mu$, X and $X_{\mu\nu}$ satisfies the conformal *Lie algebra* $\mathfrak{so}(2,2h)$ in $\tilde{x}$-space; especially there hold the commutator relations

$$[\mathrm{d}_\mu, \tilde{x}_\nu] = \delta_{\mu\nu}X + X_{\mu\nu}, \qquad [\mathrm{d}_\mu, X] = \mathrm{d}_\mu. \tag{C.10}$$

In that terminology $\tilde{x}$ is a *raising operator* which plays the role of "momentum" in the conformal algebra:

$$[\tilde{x}_\mu, \tilde{x}_\nu] = 0, \qquad \tilde{x}^2 = 0. \tag{C.11}$$

Dobrev *et al.* [46, 48, 85, 45] used the interior derivative in conformal OPE in order to construct local composite tensor operators which transform under an elementary representations of the conformal group $SO(2,2h)$. They formulated the condition for conformal invariance by means of the interior derivative on the light-cone.

For the first time, the application of the interior derivative in order to satisfy the conditions of tracelessness on the light-cone, and for antisymmetric tensors was given by Dobrev and Ganchev [49].

Additionally, the interior derivative has been used to formulate a conformal "Lorentz condition" in Conformal Quantum Electrodynamics [120, 55] and in manifestly $O(2,4)$-covariant formalism for the photon-Weyl graviton system [56].

Finally, I will show how one can use the interior derivatives for the decomposition of the local light-cone operator (see also Refs. [49, 67, 66]),

$$O_{\alpha n}(y,\tilde{x}) = \bar{\psi}(y)\gamma_\alpha(\tilde{x}\overleftrightarrow{D})^n\psi(y), \tag{C.12}$$

into all its twist parts. The operator (C.12) has the canonical dimension $d_{O_{\alpha n}} = 2h + n - 1$. We may uniquely decompose this light-cone operator according to

$$O_{\alpha n}(y, \tilde{x}) = O^{\mathrm{tw}(2h-2)}_{\alpha n}(y, \tilde{x}) + O^{\mathrm{tw}(2h-1)}_{\alpha n}(y, \tilde{x}) + \tilde{x}_\alpha\, O^{\mathrm{tw}(2h)}_{n-1}(y, \tilde{x}). \tag{C.13}$$

Using the Eq. (4.1.12) we get the operator of maximal spin, $j = n + 1$, and minimal twist $\tau = 2h - 2$

$$O^{\mathrm{tw}(2h-2)}_{\alpha n}(y, \tilde{x}) = \frac{1}{(n+1)(h+n-1)}\, \mathrm{d}_\alpha O_{n+1}(y, \tilde{x}). \tag{C.14}$$

The operator of twist $\tau = 2h - 1$ and spin $j = n$ is obtained from Eq. (4.1.28) as follows

$$\begin{aligned} O^{\mathrm{tw}(2h-1)}_{\alpha n}(y, \tilde{x}) &= \frac{2}{(n+1)(h+n-2)}\, \tilde{x}^\beta \left\{ \delta^\mu_{[\alpha} \mathrm{d}_{\beta]} + \frac{1}{2h+n-3}\, X_{[\alpha\beta]} \mathrm{d}^\mu \right\} O_{\mu n}(y, \tilde{x}), \\ &= \frac{1}{(n+1)(h+n-2)} \left\{ \delta^\mu_\alpha\, (\tilde{x}\mathrm{d}) - \tilde{x}^\mu \mathrm{d}_\alpha - \frac{n-1}{2h+n-3}\, \tilde{x}_\alpha \mathrm{d}^\mu \right\} O_{\mu n}(y, \tilde{x}). \end{aligned} \tag{C.15}$$

From the trace terms of Eqs. (C.14) and (C.15) we construct the operator with twist $\tau = 2h$ and spin $j = n - 1$ according to

$$O^{\mathrm{tw}(2h)}_{n-1}(y, \tilde{x}) = \frac{1}{(2h+n-3)(h+n-1)}\, \mathrm{d}^\mu O_{\mu n}(y, \tilde{x}). \tag{C.16}$$

The light-cone operators of geometric twist satisfy the following conditions of tracelessness on the light-cone:

$$\mathrm{d}^\alpha\, O^{\mathrm{tw}(2h-2)}_{\alpha n}(y, \tilde{x}) = 0, \tag{C.17}$$

$$\mathrm{d}^\alpha\, O^{\mathrm{tw}(2h-1)}_{\alpha n}(y, \tilde{x}) = 0. \tag{C.18}$$

The light-cone operator (C.16) is traceless by construction.

Appendix D

Target mass corrections for a scalar theory in $2h$-dimensional spacetime

The purpose of this Appendix is the investigation of the off-forward scattering amplitude with kinematical target mass corrections from the minimal twist contributions in $2h$-dimensional Euclidean spacetime ($2h \geq 3$). For simplicity we consider a scalar field theory. We will illustrate this with the simplest non-forward scattering process: the (scalar) meson production by two virtual photons

$$\gamma^*(q_1) + \gamma^*(q_2) \to M(P). \tag{D.1}$$

We denote the momenta of the two incoming photons by q_1 and q_2, and the momentum of the outgoing scalar meson by P. The non-forward scattering of this process may be written as

$$T(P, q) = \int \mathrm{d}^{2h}x\, \mathrm{e}^{\mathrm{i}qx}\, \langle 0|T(J(x)J(-x))|P\rangle, \qquad q \equiv \frac{1}{2}\big(q_2 - q_1\big), \tag{D.2}$$

with the scalar current $J(x) = \phi(x)\phi(x)$.

Using the Wick theorem and Born approximation we obtain:

$$T(J(x)J(-x)) \approx D_h(x)\big(N(x, -x) + N(-x, x)\big), \tag{D.3}$$

where $D_h(x)$ is the canonical Euclidean propagator for a scalar field in $2h$-dimensional spacetime [45]

$$D_h(x) = \frac{\Gamma(h-1)}{4\pi^h [x^2]^{h-1}}. \tag{D.4}$$

Now, we expand the nonlocal operator in an infinite tower of local ones[1]

$$N(x, -x) = \sum_{n=0}^{\infty} \frac{\mathrm{i}^n}{n!} \phi(0)(\mathrm{i}x\overleftrightarrow{D})^n \phi(0) \tag{D.5}$$

We perform the Fourier transformation by the help of the following formula[2]

$$\int \mathrm{d}^{2h}x\, \mathrm{e}^{\mathrm{i}qx} \frac{x_{\mu_1} \ldots x_{\mu_n}}{[x^2]^k} = (-\mathrm{i})^n \pi^h 2^{2h-2k+n} \frac{\Gamma(h-k+n)}{\Gamma(k)} \frac{q_{\mu_1} \ldots q_{\mu_n}}{(q^2)^{h-k+n}} + O(g_{\mu_i \mu_j}), \tag{D.6}$$

---

[1] Alternatively, we could use nonlocal operators. Then the nonlocal leading twist operator is given in terms of $\mathcal{H}_{h-1}$.

[2] More strictly, we have: $[x^2] \to [x^2 - \mathrm{i}0]$ and $(q^2) \to (q^2 + \mathrm{i}0)$.

in particular with

$$\int \mathrm{d}^{2h}x \, \frac{\mathrm{e}^{\mathrm{i}qx}}{[x^2]^{h-1}} \langle 0|N_n(x)|P\rangle = (-\mathrm{i})^n \pi^h 2^{2+n} \frac{\Gamma(1+n)}{\Gamma(h-1)} \frac{1}{(q^2)^{1+n}} \langle 0|N_n(q)|P\rangle, \tag{D.7}$$

where we neglected all terms involving $\Box_q$, since they vanish for leading twist. The leading twist operator has $\tau = 2(h-1)$. It is in the q-space a harmonic polynomial of degree n in q which satisfies $\Box_q N_n^{\mathrm{l.t.}}(q) = 0$. Eventually, we use the harmonic projector in the q-space and obtain for the corresponding matrix element

$$\begin{aligned} \langle 0|N_n^{\mathrm{l.t.}}(q)|P\rangle &= \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (h+n-k-2)!}{4^k k! (h+n-2)!} \, q^{2k} \Box_q^k (qP)^n f_n \\ &= \frac{n!}{(h+n-2)!} \sum_{k=0}^{[\frac{n}{2}]} \frac{(-1)^k (h+n-k-2)!}{4^k k! (n-2k)!} \, q^{2k} P^{2k} (qP)^{n-2k} f_n \\ &= \frac{n!}{(h-1)_n} \left(\frac{1}{2} \sqrt{q^2 P^2} \right)^n C_n^{h-1} \left(\frac{qP}{\sqrt{q^2 P^2}} \right) f_n, \end{aligned} \tag{D.8}$$

where f_n is the reduced matrix element. The rewriting from the first to the third line in Eq. (D.8) is nothing else but another form of a harmonic polynomial which is related to the irreducible representation of the group $SO(2h)$. Accordingly, the scattering amplitude in terms of local operators appears as an expansion with respect to Gegenbauer polynomials:

$$T(P,q) = \sum_{n=0}^{\infty} \frac{\Gamma(n+1)\Gamma(h-1)}{\Gamma(n+h-1)} \frac{1}{q^2} \left(\sqrt{\frac{P^2}{q^2}} \right)^n C_n^{h-1} \left(\frac{qP}{\sqrt{q^2 P^2}} \right) (1 + (-1)^n) f_n, \tag{D.9}$$

which is very close to Nachtmann's [107] target mass corrections in inclusive processes for a scalar field theory ($h = 2$). Using the integral representation of Euler's beta functions for the Gamma functions in Eq. (D.9)

$$\begin{aligned} \frac{\Gamma(n+1)\Gamma(h-1)}{\Gamma(n+h-1)} &= \frac{(n+h-1)!}{(n+h-2)!} B(n+1, h-1) \\ &= (h-1+n) \int_0^1 \mathrm{d}t \, t^n (1-t)^{h-2}, \end{aligned} \tag{D.10}$$

and replace $nt^n \to t\partial_t t^n$, we get the following expression for the scattering amplitude

$$\begin{aligned} T(P,q) = \int_{-1}^{1} \mathrm{d}\xi \, f(\xi) \int_0^1 \mathrm{d}t \, (1-t)^{h-2} \big(h-1+t\partial_t\big) \frac{1}{q^2} \\ \times \sum_{n=0}^{\infty} \left(t\xi \sqrt{\frac{P^2}{q^2}} \right)^n C_n^{h-1} \left(\frac{qP}{\sqrt{q^2 P^2}} \right) (1 + (-1)^n). \end{aligned} \tag{D.11}$$

The resummation in Eq. (D.9) can be done by making use of the generating function for the Gegenbauer polynomials

$$\begin{aligned} \sum_{n=0}^{\infty} a^n C_n^{h-1}(b) &= (1 - 2ab + a^2)^{-(h-1)} \\ &= \frac{(q^2)^{h-1}}{(q - t\xi P)^{2(h-1)}}. \end{aligned} \tag{D.12}$$

Here $a = t\xi\sqrt{P^2/q^2}$ and $b = qP/\sqrt{q^2P^2}$. Finally, we obtain the scattering amplitude with all mass-corrections on twist $\tau = 2(h-1)$ level

$$T(P,q) = \int_{-1}^{1} \mathrm{d}\xi\, f(\xi) \int_{0}^{1} \mathrm{d}t\, (1-t)^{h-2} (q^2)^{h-2} \big(h-1+t\partial_t\big) \left(\frac{1}{(q+t\xi P)^{2(h-1)}} + \frac{1}{(q-t\xi P)^{2(h-1)}} \right). \tag{D.13}$$

If we carry out the derivative in Eq. (D.13), we obtain

$$\begin{aligned} T(P,q) = \int_{-1}^{1} \mathrm{d}\xi\, f(\xi) \int_{0}^{1} \mathrm{d}t\, (1-t)^{h-2} (q^2)^{h-2} \Big\{ \big(h-1+2t^2\big) \left(\frac{1}{(q+t\xi P)^{2(h-1)}} + \frac{1}{(q-t\xi P)^{2(h-1)}} \right) \\ - 2t^2(h-1) \left(\frac{t\xi\, qP + t^2\xi^2P^2}{(q+t\xi P)^{2h}} - \frac{t\xi\, qP - t^2\xi^2P^2}{(q-t\xi P)^{2h}} \right) \Big\}. \end{aligned} \tag{D.14}$$

Thus, we are able to express the scattering amplitude in terms of a DA $f^{(\tau)}(\xi)$ with twist $\tau = 2(h-1)$ and a mass-dependent coefficient function.

For $h = 2$ we perform the partial integration with respect to t in Eq. (D.13) and obtain the simple expression for the scattering amplitude

$$T(P,q) = \int_{-1}^{1} \mathrm{d}\xi\, f(\xi) \left(\frac{1}{(q+\xi P)^2} + \frac{1}{(q-\xi P)^2} \right). \tag{D.15}$$

From this point of view, one may introduce the following kinematical DA:

$$\mathcal{F}(\xi,q,P) = f(\xi) \left(\frac{1}{(q+\xi P)^2} + \frac{1}{(q-\xi P)^2} \right), \tag{D.16}$$

which contains the power corrections of the DA $f(\xi)$ on twist $\tau = 2$ level. If we expand the scattering amplitude (D.15) in powers of P^2/q^2, we finally get the simple formula:

$$T(P,q) = \frac{2}{q^2} \sum_{n=0}^{\infty} (n+1) \frac{P^n}{q^n}\, f_n, \qquad n - \text{even}. \tag{D.17}$$

Bibliography

[1] K. Abe et al. (E143), Phys. Rev. Lett. **76** (1996) 587.

[2] M.A. Ahmed and G.G. Ross, Phys. Lett. B **56** (1975) 385; Nucl. Phys. B **111** (1976) 441.

[3] A. Ali, V.M. Braun, and H. Simma, Z. Phys. C **63** (1994) 437.

[4] I.V. Anikin, B. Pire, and O.V. Teryaev, Phys. Rev. D **62** (2000) 071501.

[5] I.V. Anikin and O.V. Teryaev, *Wandzura-Wilczek approximation from generalized rotational invariance*, `hep-ph/0102209`.

[6] S.A. Anikin and O.I. Zavialov, Ann. Phys. (N.Y.) **116** (1978) 135.

[7] X. Artru and M. Mekhfi, Z. Phys. C **45** (1990) 669.

[8] I.I. Balitsky, Phys. Lett. B **124** (1983) 230.

[9] I.I. Balitsky and V.M. Braun, Nucl. Phys. B **311** (1988/89) 541.

[10] I.I. Balitsky, V.M. Braun, and A.V. Kolesnichenko, Nucl. Phys. B **312** (1989) 509.

[11] I.I. Balitsky and V.M. Braun, Nucl. Phys. B **361** (1991) 93.

[12] I.I. Balitsky and A.V. Radyushkin, Phys. Lett. B **413** (1997) 114.

[13] P. Ball and V.M. Braun, Phys. Rev. D **54** (1996) 2182.

[14] P. Ball, V.M. Braun, Y. Koike, and K. Tanaka, Nucl. Phys. B **529** (1998) 323.

[15] P. Ball and V.M. Braun, Nucl. Phys. B **543** (1999) 201.

[16] P. Ball, JHEP **9901** (1999) 010.

[17] P. Ball and M. Lazar, Phys. Lett. B **515** (2001) 131.

[18] V. Baluni and E. Eichten, Phys. Rev. Lett. **37** (1976) 1181; Phys. Rev. D **14** (1976) 3045.

[19] V. Bargmann and I.T. Todorov, J. Math. Phys. **18** (1977) 1141.

[20] A.O. Barut and R. Raczka, Theory of Group Representations and Applications PWN – Polish Scientific Publishers, Warszawa, 1977.

[21] H. Bateman and A. Erdelyi, Higher Transcendental Functions, Vol. **2**, New York 1953.

[22] A.V. Belitsky and D. Müller, Nucl. Phys. B **589** (2000) 611.

[23] A.V. Belitsky and D. Müller, Phys. Lett. B **507** (2001) 173.

[24] J. Blümlein, B. Geyer, and D. Robaschik, Phys. Lett. B **406** (1997) 161.

[25] J. Blümlein and N. Kochelev, Nucl. Phys. B **498** (1997) 285.

[26] J. Blümlein and A. Tkabladze, Nucl. Phys. B **553** (1999) 427.

[27] J. Blümlein, B. Geyer, and D. Robaschik, Nucl. Phys. B **560** (1999) 283.

[28] J. Blümlein and D. Robaschik, Nucl. Phys. B **581** (2000) 449.

[29] J. Blümlein, Eur. Phys. J. C **20** (2001) 683.

[30] M. Bordag and D. Robaschik, Nucl. Phys. B **169** (1980) 445.

[31] R.A. Brandt and G. Preparata, Nucl. Phys. B **27** (1971) 541.

[32] V.M. Braun and I.E. Filyanov, Z. Phys. C **44** (1989) 157.

[33] V.M. Braun and I.E. Filyanov, Z. Phys. C **48** (1990) 239.

[34] Th. Braunschweig, B. Geyer, and D. Robaschik, Ann. Phys. (Leipzig) **44** (1987) 407.

[35] Th. Braunschweig, B. Geyer, J. Hořejši, and D. Robaschik, Z. Phys. C **33** (1987) 275.

[36] S.J. Brodsky, Y. Frishman, G.P. Lepage, and C. Sachrajda, Phys. Lett. B **91** (1980) 239.

[37] A.P. Bukhvostov, E.A. Kuraev, and L.N. Lipatov, Zh. Eksp. Teor. Fiz. **87** (1984) 37.

[38] A.P. Bukhvostov, G.V. Frolov, E.A. Kuraev, and L.N. Lipatov, Nucl. Phys. B **258** (1985) 601.

[39] A.P. Bukhvostov and G.V. Frolov, Yad. Fiz. **45** (1987) 1136; [Sov. J. Nucl. Phys. **45** (1987) 704].

[40] H. Burkhardt and W.N. Cottingham, Ann. Phys. (N.Y.) **56** (1970) 453.

[41] M. Burkardt, Phys. Rev. D **52** (1995) 3841.

[42] V.L Chernyak and A.R. Zhitnitsky, Phys. Rep. **112** (1984) 173.

[43] J.C. Collins and D.E. Soper, Nucl. Phys. B **194** (1982) 445.

[44] N.S. Craigie and J. Stern, Nucl. Phys. B **216** (1983) 209.

[45] N.S. Craigie, V.K. Dobrev, and I.T. Todorov, Ann. Phys. (N.Y.) **159** (1985) 411.

[46] V.K. Dobrev, V.B. Petkova, S.G. Petrova, and I.T. Todorov, Phys. Rev. D **13** (1976) 887.

[47] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova, and I.T. Todorov, Rep. Math. Phys. **9** (1976) 219.

[48] V.K. Dobrev, G. Mack, V.B. Petkova, S.G. Petrova, and I.T. Todorov, Harmonic Analysis of the n-Dimensional Lorentz Group and its Applications to Conformal Quantum Field Theory, Lecture Notes in Physics, No. **63** (Springer, 1977).

[49] V.K. Dobrev and A.Ch. Ganchev, *Conformal operators from spinor fields: Antisymmetric tensor case*, Dubna Report E2-82-881 (1982).

[50] A.V. Efremov and A.V Radyushkin, Phys. Lett. B **94** (1980) 245.

[51] A.V. Efremov, O.V. Teryaev, and E. Leader, Phys. Rev. D **55** (1997) 4307.

[52] B. Ehrnsperger, L. Mankiewicz, and A. Schäfer, Phys. Lett. B **323** (1994) 439.

[53] R.K. Ellis, W. Furmanski, and R. Petronzio, Nucl. Phys. B **207** (1982) 1; Nucl. Phys. B **212** (1983) 29.

[54] S. Ferrara, R. Gatto, and A.F. Grillo, Nucl. Phys. B **34** (1971) 349.

[55] P. Furlan, V.B. Petkova, G.M. Sotkov, and I.T. Todorov, Riv. Nuovo Cimento **8** (1985) 1.

[56] P. Furlan, V.B. Petkova, and G.M. Sotkov, J. Math. Phys. **28** (1987) 251.

[57] H. Georgi and H.D. Politzer, Phys. Rev. D **9** (1974) 416.

[58] H. Georgi and H.D. Politzer, Phys. Rev. D **14** (1976) 1829.

[59] B. Geyer, D. Robaschik, and E. Wieczorek, Fortschr. Phys. **27** (1979) 75.

[60] B. Geyer, D. Robaschik, M. Bordag, and J. Hořejši, Z. Phys. C **26** (1985) 591.

[61] B. Geyer, D. Müller, and D. Robaschik, Nucl. Phys. B (Proc. Suppl.) **51 C** (1996) 106.

[62] B. Geyer, D. Müller, and D. Robaschik, *The evolution of non-singlet twist-3 parton distribution functions*, DESY 96-239, Proc. of Third Meeting on Prospects in Nucleon-Nucleon Spin Physics at HERA, Dubna, 1996, `hep-ph/9611452`.

[63] B. Geyer, M. Lazar, and D. Robaschik, Nucl. Phys. B **559** (1999) 339.

[64] B. Geyer and M. Lazar, Nucl. Phys. B **581** (2000) 341.

[65] B. Geyer, M. Lazar, and D. Robaschik, Nucl. Phys. B **618** (2001) 99.

[66] B. Geyer and M. Lazar, Phys. Rev. D **63** (2001) 094003.

[67] B. Geyer and M. Lazar, *Twist decomposition of nonlocal light-ray operators and harmonic tensor functions*, Proceedings of the International Conference "Quantum Theory and Symmetries", edited by H.-D. Doebner, V.K. Dobrev, J.-D. Hennig and W. Lücke (World Scientific, Singapore, 2000) p. 362; `hep-th/9911024`.

[68] S. Gottlieb, Nucl. Phys. B **139** (1978) 125.

[69] D.J. Gross and S.B. Treiman, Phys. Rev. D **4** (1971) 1059.

[70] D.J. Gross and F. Wilczek, Phys. Rev. D **9** (1974) 980.

[71] A.H. Guth and D.E. Soper, Phys. Rev. D **12** (1975) 1143.

[72] M. Hamermesh, Group theory and its application to physical problems, Addison-Wesley, London (1962).

[73] P. Hoodbhoy and X. Ji, Phys. Rev. D **50** (1994) 4429.

[74] G. Itzykson and J.-B. Zuber, Quantum Field Theory, Mc Graw Hill, New York, 1980.

[75] R.L. Jaffe and M. Soldate, Phys. Lett. B **105** (1981) 467; Phys. Rev. D **26** (1982) 49.

[76] R.L. Jaffe, Nucl. Phys. B **229** (1983) 205.

[77] R.L. Jaffe and X. Ji, Phys. Rev. D **43** (1991) 724.

[78] R.L. Jaffe and X. Ji, Phys. Rev. Lett. **67** (1991) 552.

[79] R.L. Jaffe and X. Ji, Nucl. Phys. B **375** (1992) 527.

[80] R.L. Jaffe, *Spin, twist and hadron structure in deep inelastic processes*, Erice Lectures, `hep-ph/9602236`.

[81] X. Ji, Phys. Lett. B **289** (1992) 137.

[82] X. Ji, Nucl. Phys. B **402** (1993) 217.

[83] X. Ji, Phys. Rev. Lett., **78** (1997) 610, Phys. Rev. D. **55** (1997) 7114, J. Phys. G **24** (1998) 1181.

[84] N.I. Karchev, Nucl. Phys. B **211** (1983) 55.

[85] N. Karchev and I.T. Todorov, *Conformal Composite Fields and Operator Product Expansions*, ZiF-Preprint, Bielefeld (1983).

[86] H. Kawamura and T. Uematsu, Phys. Lett. B **343** (1995) 346.

[87] N. Kivel, M.V. Polyakov, A. Schäfer, and O.V. Teryaev, Phys. Lett. B **497** (2001) 73.

[88] J. Kodaira, S. Matsuda, K. Sasaki, and T. Uematsu, Nucl. Phys. B **159** (1979) 99.

[89] J. Kodaira, Y. Yasui, and T. Uematsu, Phys. Lett. B **344** (1995) 348.

[90] J. Kodaira, S. Matsuda, T. Uematsu, and K. Sasaki, Phys. Lett. B **345** (1995) 527.

[91] J. Kodaira, Y. Yasui, K. Tanaka, and T. Uematsu, Phys. Lett. B **387** (1996) 855.

[92] J. Kodaira and K. Tanaka, Prog. Theor. Phys **101** (1999) 191.

[93] K. Kogut and D.E. Soper, Phys. Rev. D **1** (1970) 2901.

[94] Y. Koike and K. Tanaka, Phys. Rev. D **51** (1995) 6125.

[95] Y. Koike and N. Nishiyama, Phys. Rev. D **55** (1997) 3068.

[96] M. Lazar, *Konstruktion und Anwendung irreduzibler nichtlokaler Lichtkegeloperatoren in der Quantenchromodynamik*, Diplomarbeit, Leipzig 1998.

[97] M. Lazar, Phys. Lett. B **497** (2001) 62; Phys. Lett. B **506** (2001) 385 (Erratum).

[98] M. Lazar, JHEP **0105** (2001) 029.

[99] G.P. Lepage and S.J. Brodsky, Phys. Lett. B **87** (1979) 359.

[100] S.P. Luttrell, S. Wada, and B.R. Webber, Nucl. Phys. B **188** (1981) 219.

[101] S.P. Luttrell and S. Wada, Nucl. Phys. B **197** (1982) 290.

[102] Yu.M. Makeenko, Sov. J. Nucl. Phys. **33** (1981) 440.

[103] A.V. Manohar, Phys. Rev. Lett. **65** (1990) 2511, Phys. Rev. Lett. **66** (1991) 289.

[104] S. Matsuda and T. Uematsu, Nucl. Phys. B **168** (1980) 181.

[105] M. Maul, B. Ehrnsperger, E. Stein, and A. Schäfer, Z. Phys. A **356** (1997) 443.

[106] D. Müller, D. Robaschik, B. Geyer, F.-M. Dittes, and J. Hořejši, Fortsch. Phys. **42** (1994) 101.

[107] O. Nachtmann, Nucl. Phys. B **63** (1973) 273; Nucl. Phys. B **78** (1974) 455.

[108] Th. Ohrndorf, Nucl. Phys. **198** (1982) 26.

[109] M. Okawa, Nucl. Phys. B **172** (1980) 481; Phys. B **187** (1981) 71.

[110] A. Piccione and C. Ridolfi, Nucl. Phys. B **513** (1998) 301.

[111] H.D. Politzer, Nucl. Phys. B **172** (1980) 349.

[112] P. Prudnikov, Yu.A. Brychov, and O.I. Marichev, Integrals and Series, Vol. **2**, Special Functions, Gordon & Breach, New York 1988.

[113] P.G. Ratcliffe, Nucl. Phys. B **264** (1986) 493.

[114] A.V. Radyushkin, Phys. Lett. B **380** (1996) 417; Phys. Rev. D **56** (1997) 5524; Phys. Rev. D **59** (1999) 014030; Phys. Lett. B **449** (1999) 81;

[115] A.V. Radyushkin and C. Weiss, Phys. Lett. B **493** (2000) 332; Phys. Rev. D **63** (2001) 114012.

[116] M.A. Shifman and M.I. Vysotsky, Nucl. Phys. B **186** (1981) 475.

[117] E.V. Shuryak and A.I. Vainshtein, Nucl. Phys. B **199** (1982) 951; Nucl. Phys. B **201** (1982) 141.

[118] G. Stoll, *Light-cone wave functions of mesons*, `hep-ph/9812432`.

[119] R.D. Tangerman and P.J. Mulders, *Polarized twist three distributions* g_2 *and* h_2 *and the role of intrinsic transvers momentum*, `hep-ph/9408305`.

[120] I.T. Todorov, *Local Field Representations of the Conformal Group and their Applications*, in: Mathematics and Physics, Lectures on Recent Results, Vol.**1**, L. Streit, Ed., Word Scientific, Singapore (1985).

[121] N.Ya. Vilenkin, Special functions and the theory of group representations, Moscow 1965 (in Russian), Ch. IX; translated in: Translations of Mathematical Monographs Vol. 22, AMS, Providence 1968,
N.Ya. Vilenkin and A.U. Klimyk, Representations of Lie groups and Special Functions, Vol. 2, Kluwer, Dordrecht 1993.

[122] S. Wada, Progr. Theor. Phys. **62** (1979) 475; Nucl. Phys. B **202** (1983) 201.

[123] S. Wandzura and F. Wilczek, Phys. Lett. B **72** (1977) 195.

[124] S. Wandzura, Nucl. Phys. B **122** (1977) 412.

[125] K.G. Wilson, Phys. Rev. **179** (1969) 1499.

[126] O.I. Zavialov, Renormalized Feynman Diagrams, Nauka, Moscow 1979 (in Russian); (extended) English edition: Renormalized Quantum Field Theory, Kluwer, Dortrecht 1990.